

## Partial Purification and Characterization of Chitin Deacetylase Produced by *Bacillus thermoleovorans* LW-4-11

Aris Toharisman<sup>1</sup> & Maggy T. Suhartono<sup>2</sup>

<sup>1</sup>Indonesian Sugar Research Institute

Jl. Pahlawan 25 Pasuruan 67126, East Java-INDONESIA, Email: atoharis@telkom.net

<sup>2</sup>Research Center for Biotechnology IPB, Jln. Puspa, Kotak Pos 1 Darmaga Campus,  
Bogor Agricultural University, Bogor 16180 – INDONESIA

### ABSTRACT

**Purifikasi Parsial dan Karakterisasi Enzim Kitin Deasetilase yang Dihasilkan *Bacillus thermoleovorans* LW-4-11.** Kitin deasetilase yang dihasilkan dari bakteri *Bacillus thermoleovorans* LW-4-11 telah dimurnikan menggunakan amonium sulfat 70% diikuti dengan perlakuan panas (suhu 70°C selama 1 jam). Kemurnian enzim yang dipisahkan dari medium fermentasi meningkat 4.28 kali dengan aktivitas spesifik sekitar 43.7 mU/mg. Enzim memiliki suhu dan pH optimum masing-masing 80°C dan 6.0 dalam substrat *O*-hydroxyethylated chitin (glycol chitin). Kitindeasetilase ini relatif tahan panas dengan waktu-paruh sekitar 30 menit pada suhu 80 °C. Enzim dihambat oleh ion Li<sup>+</sup>, Zn<sup>2+</sup>, Mn<sup>2+</sup>, Co<sup>2+</sup> and Ni<sup>+</sup> pada konsentrasi 1 mM, tetapi diaktifkan oleh EDTA 1 mM.

**Kata kunci:** purifikasi parsial, karakterisasi, kitin deasetilase, *B. thermoleovorans* LW-4-11

### INTRODUCTION

Chitin, a homopolymer of b-(1-4)-linked *N*-acetyl-D-glucosamine, is one of the most abundant, easily obtained, and renewable natural polymers, second only to cellulose. It is commonly found in fungi, marine vertebrates, and insects (Sandford 1989; Patil *et al.* 2000). Chitin is an insoluble material and its industrial use is still limited. Chitosan, a partially deacetylated form of chitin, is water soluble and has a large number and a wide variety of important applications (Muzzarelli 1996; Somashekar & Joseph 1996; Shahidi *et al.* 1999; Tsigos *et al.* 2000).

Chitosan is produced by the thermochemical deacetylation of chitin which leads to heterogeneous end-product owing to the severity of the treatment (Chang *et al.* 1997; Kolodziejska *et al.* 2000). The enzymatic conversion of chitin to chitosan using chitin deacetylase (CDA) provides an attractive alternative to presently employed chemical process. CDA catalyses the conversion of chitin to chitosan by the deacetylation of *N*-acetyl-D-glucosamine residues in chitin oligosaccharide (Tsigos *et al.* 2000; Jaworska & Konieczna 2006).



CDA's have been identified and characterized from several extracts of fungi (Gao *et al.* 1995; Deising & Siegrist 1995; Alfonso *et al.* 1995; Tokuyasu *et al.* 1996; Christodoulidou *et al.* 1999). However, CDA's from bacteria have been rarely reported (Toharisman *et al.* 2001). This paper describes the isolation, characterization, and partial purification of thermostable CDA from *B. thermoleovorans* LW-4-11.

## MATERIALS AND METHODS

Chitin, glycol chitosan and glucosamine were purchased from Sigma Chemicals. Glycol chitin was prepared from glycol chitosan with reacylation using acetic anhydride. Reagents for protein determination and sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis were purchased from Bio-Rad. All other chemicals were the highest grade available.

### Microorganisms and Cultivation

*B. thermoleovorans* LW-4-11, a chitinase producing bacterium, was used for the production of CDA. The strain was isolated from Langoan hot spring water in North Sulawesi by the method of Srinivasan (2000) on minimal medium containing of ( $\text{g l}^{-1}$ ): Bacto yeast extract, 10;  $(\text{NH}_4)_2\text{SO}_4$ , 4;  $\text{KH}_2\text{PO}_4$ , 0.15 g; and 100 mg chitin (100 mesh). For enzyme production, bacterium was cultured on Thermus medium (Takayanagi *et al.*, 1991) containing  $(\text{NH}_4)_2\text{SO}_4$ , 0.7%;  $\text{K}_2\text{HPO}_4$ , 0.1%; NaCl, 0.1%;  $\text{MgSO}_4$

$7\text{H}_2\text{O}$ , 0.01%, Bacto yeast extract, 0.2%, Bacto trypton, 0.1% and colloidal chitin, 1% and incubated at 70 °C for 3 days.

### Preparation of Enzyme

The bacterial cells were separated from culture broth by centrifugation at 10 000 x g for 15 minutes. The supernatant was brought to 70% saturation with ammonium sulphate at 4 °C and allowed to settle down overnight. The precipitate was recovered by centrifugation (10 000 x g, 15 minutes) and the pellet formed was solubilized in 20 mM Tris-HCl buffer, pH 7.0. The solution was dialyzed overnight against the same buffer at 4 °C and then incubated at 70 °C for 1 h to remove other heat-unstable proteins.

### Chitin Deacetylase Assay

Chitin deacetylase activity was estimated using glycol chitin as a substrate (Tokuyasu *et al.* 1996). The assay mixture contained 0.15% glycol chitin dissolved in 20 mM tetraborate/HCl buffer (pH 8.5). Reaction was initiated by the addition of 200 ml (crude extract) or 50 ml (ammonium sulphate precipitated) enzyme solution to 123 ml of reaction mixtures, incubated at 70 °C for 30 min. The reaction was terminated by the addition of 200 ml of 33% (v/v) acetic acid. The control was prepared by adding the enzyme solution after inactivation. Upon termination of the reaction, the concentration of glucosamine residues produced by the deacetylation reaction was estimated by oxidation using  $\text{NaNO}_2$ , followed by a spectrophotometric method using indole/HCl (Dische & Borenfreund 1950).



Chitin deacetylase activity is defined as the amount of enzyme that liberated 1 mmol of glucosamine (GlcNA) per min at 420. nm. Protein content was determined using the dye-binding method with bovine serum albumin fraction V as a standard (Bradford 1976).

### Effect of pH, temperature and additives

The pH and temperature optima of the enzyme as well as the effect of metal ion concentration on the activity were measured after incubation for 30 min at various pHs, temperatures (60-100 °C) and metal ion concentration, respectively. The pH was adjusted by using the following buffers: glycine-HCl (pH 3.0-5.0),  $\text{NaH}_2\text{PO}_4 - \text{Na}_2\text{HPO}_4$  (pH 6.0-8.0), borate (pH 8.0-9.0), and glycine-NaOH (pH 10.0-12.0). Metal ions analysed included LiCl,  $\text{MnCl}_2$ ,  $\text{ZnCl}_2$ ,  $\text{CoCl}_2$ ,  $\text{MgCl}_2$ , and  $\text{NiCl}_2$ . The effect of EDTA upon the enzyme activity was also

examined at a concentration of 1 mM. The thermostability of enzyme was measured by the residual CDA activities after incubation of enzyme solution at optimum temperature with the corresponding buffers. Samples were withdrawn every 15 minutes and the activities were measured.

## RESULTS

### Isolation and Identification of Microorganism

Microorganisms isolated from spring water collected at different locations in Langoan, North Sulawesi were screened on agar plates containing 1% of colloidal chitin at 70 °C for 4 days. Colonies that grew well under such conditions were isolated and retained for subsequent screening. Eighty strain of bacteria were obtained from the first screening and then were subcultured in liquid media containing 0.5% colloidal chitin. After incubation, the supernatants





Figure 1. Time course of CDA production by *B. thermoleovorans* LW-4-11

**Table 1.** Partial Purification Steps of CDA Produced by *B. thermoleovorans* LW-4-11

Steps	Total Activity (mU)	Protein (mg)	Specific Activity (mU/mg)	Purification (Fold)	Recovery (%)
Free-Cell Medium	21.4	2.10	10.2	1.00	100.0
Ammonium Sulfate ppt (70%)	18.5	1.21	15.3	1.50	86.4
Heat Treatment (70 °C, 1 h)	7.9	0.18	43.7	4.28	36.7



**Figure 2.** The optimal pH of chitin deacetylase from *B. thermoleovorans* LW-4-11; ●, 0.2 M citrate buffer; ▲, 0.2 M phosphate buffer; and ■, 0.2 M glycine-NaOH buffer

were collected for measurement of CDA activity. One strain showed the highest activity and used throughout the study. It produced CDA extracellularly at the maximum level after 4<sup>th</sup> days cultivation (Figure 1). According to 16S rRNA sequence, the strain was identified as *B. thermoleovorans* (Suhartono 2002).

**Enzyme Preparation**

The result of partial purification procedure are summarized in Table 1. The

enzyme from 4 days cell-free medium of LW-4-11 was precipitated by the addition of ammonium sulphate up to 70% saturation. This step resulted in 1.50-fold increase in the purity of enzyme. The heat treatment of enzyme increased the purity of enzyme to be 4.28 fold.

**Characterization of Enzyme**

The effect of pH, temperature and addition of additives on CDA activity






Figure 3. The optimal temperature of chitin deacetylase from *B. thermoleovorans* LW-4-11




Figure 4. Enzyme stability at; ◆, 70 °C; ■, 80 °C; ▲, 90 °C; and ●, 100 °C

was studied by using glycol chitin as a substrate under the standard assay conditions. The dialyzed CDA exhibited maximum activity at pH 6.0 (Figure 2). The optimum temperature of enzyme was found to be 80 °C (Figure 3). To examine the thermostability of the enzyme, CDA solution in 50 mM sodium tetraborate buffer (pH 7.0) was allowed to stand during 5 hour at various tempe-

ratures, and the residual activity was measured. The enzyme was stable up to 80 °C with a loss of activity less than 80% for 30 min (data not shown). However about 50% of the original CDA activity was lost after incubation at 90 °C for 20 minutes and at 100 °C after 5 minutes, respectively (Figure 4). The CDA was activated by EDTA. It was inhibited by  $\text{Li}^+$ ,  $\text{Mn}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Co}^{2+}$  and  $\text{Ni}^+$




Figure 5. Effect of various cations and EDTA (1 mM) on CDA activity

tested as chlorides at 1 mM concentration (Figure 5).

## DISCUSSION

Chitosan is usually produced from chitin by thermochemical deacetylation. This process results in a polymer product having a broad distribution of molecular mass and a heterogeneous extent of deacetylation. However, for many potentially important applications, uniform material with specific physical and chemical properties is required (Martinou *et al.* 1993). An alternative procedure that would exploit the enzymatic deacetylation of chitin was using chitin deacetylase. The method did not cause the degradation of sugar chain and produces no alkaline wastes (Tokuyasu *et al.* 1996). Chitin deacetylases (EC 3.5.1.41) which represent a class of hydrolytic enzymes were mainly found in fungi. It was first demonstrated in extracts of the fungus *Mucor rouxii*

(Davis & Garcia 1984). Chitin deacetylase activity also had been found in other *Zygomycetes* (Alfonso *et al.* 1995), *Colletotrichum lindemuthianum* (Tsigos *et al.* 2000), *Uromyces viciaefabae* (Deisinga & Siegrist 1995), *Saccharomyces cerevisiae* (Mishra *et al.* 1997; Christodoulidou *et al.* 1999) and in cucumber leaves infected with *Colletotrichum lagenarium* (Siegrist & Kauss 1990).

CDA from other microorganisms mainly bacteria was rarely reported. We have isolated and characterized CDA from acidophilic *Bacillus* sp. isolated from Kamojang Crater West Java (Natsir 2000; Rahayu 2000) and *B. stearothermophilus* from Langoan, North Sulawesi (Toharisman *et al.* 2001). In this study, we tried to isolate another thermophile from hot spring water and sediment from North Sulawesi having different characteristics. *Bacillus thermoleovorans* LW-4-11 was able to grow at 70°C and had the high CDA activity. It produced



CDA extracellularly having optimum temperature of 80 °C. It might be one of the highest optimum temperatures for CDA as all optimum temperature of CDAs reported so far was 50 °C (Alfonso *et al.* 1995; Tokuyasu *et al.* 1996; Christodoulidou *et al.* 1999; Natsir 2000; Rahayu 2000).

The activity of the enzyme was retained by over 50% when it was kept for 45 minutes at 70 °C. The half-lives of the activity ( $T_{1/2}$ ) at 80 and 90 °C were about 30 min and 20 min, respectively. When the enzyme was incubated at 70°C for a hour, heat-labile proteins were denatured. This strategy was applied in partial purification processes described in this paper and could purify enzyme of 4.28 fold with overall yield of 36.7%. Similar procedures were conducted for CDA purification from *M. rouxii* at 50 °C for 10 minutes (Martinou *et al.*, 1993) and from *Bacillus sp.* at 50 °C for 24 h (Rahayu 2000).

Other characteristics of the enzyme were also performed. The optimum pH of the enzyme was 6.0, similar to CDA from *M. rouxii* (Kafetzopoulos *et al.* 1993). Most of the optimum pH of CDAs reported so far was between 4.5 and 8.5 (Tsigos *et al.* 2000). The effect of cations were those similar to those of CDA of *C. lindemuthianum* (Tokuyasu *et al.* 1996) and *Aspergillus nidulans* (Alfonso *et al.* 1995). The enzyme was activated by EDTA, whereas and inhibited by metal ions such as  $Zn^{2+}$ ,  $Mn^{2+}$  and  $Cu^{2+}$  (1 mM).

## ACKNOWLEDGMENTS

This study was supported by Research Center for Biotechnology Bogor Agricultural University and Research Grant from Bioproducts Research Center Yonsei University, Korea.

## REFERENCES

- Alfonso, C., OM. Nuero, F. Santamaria, & F. Reyes. 1995. Purification of a heat-stable chitin deacetylase from *aspergillus nidulans* and its role in cell wall degradation. *Curr. Microbiol.* 30 (1): 49-54.
- Chang, KLB, G. Tsai, J. Lee, & WR Fu. 1997. Heterogeneous N-deacetylation of chitin in alkaline solution. *Carbohydr. Res.* 303: 327-332.
- Christodoulidou, A., P. Briza, A. Ellinger, & V. Bouriotis. 1999. Yeast ascospore wall assembly requires two chitin deacetylase isozymes. *FEBS Lett.* 29 (2):275-279.
- Davis, LL., & SB. Garcia. 1984. Chitosan synthesis by the tandem action of chitin synthetase and chitin deacetylase from *Mucor rouxii*. *Biochemistry.* 23: 1065-1073.
- Deising & J. Siegrist. 1995. Chitin Deacetylase Activity of the rust *Uromyces viciae-fabae* is controlled by fungal morphogenesis: *FEMS Microbiol. Lett.* 127: 207-212.
- Dische, Z. & E. Borenfreund. 1950. A Spectrophotometric methods for the



- microdetermination of hexosamines. *J. Biol. Chem.* 184: 517-522.
- Gao, XD, T. Katsumoto, & K. Onodera. 1995. Purification and characterization of chitin deacetylase from *Absidia coerulea*. *J. Biochem.* 117: 257-263.
- Jaworska, MM & E. Konieczna. 2006. Stability of chitin deacetylase. *Polish Chitin Society, Monograph XI*: 15-19.
- Kafetzopoulos, D., A. Martinou, V. Bouriotis. 1993. Bioconversion of chitin to chitosan: purification and characterization of chitin deacetylase from *Mucor rouxii*. *Proc. Natl. Acad. Sci. USA.* 90: 2564-2568.
- Kolodziejaska, I., A. Wojtasz-Pajak, G. Ogonowska, & ZE Sikorski. 2000. Deacetylation of chitin in a two-stage chemical and enzymatic process. *Bul. Sea Fish. Inst.* 2 (150): 15-24.
- Martinou, A., D. Kafetzopoulos, & V. Bouriotis. 1993. Isolation of chitin deacetylase from *Mucor rouxii* by immunoaffinity chromatography. *J. Chromatogr.* 644: 35-41.
- Mishra, C., SE. Semino, KJ. McCreath, H. de la Vega, BJ. Jones, CA. Specht, & PW. Robbins. 1997. Cloning and expression of two chitin deacetylase genes of *Saccharomyces cerevisiae*. *Yeast.* 13(4): 327-336.
- Muzzarelli, RAA. 1996. Chitosan-based dietary foods. *Carbohydr. Polym.* 29: 309-316.
- Patil, RS, V. Ghormade, & M.V. Deshpande. 2000. Chitinolytic enzymes: an exploration. *Enzyme and Microbial Technology.* 26; 473-483.
- Natsir, HD. 2000. Biochemical Characteristics of chitinase enzyme from *Bacillus* Sp. of Kamojang Crater, Indonesia. [Master Thesis]. Bogor Agricultural University.
- Rahayu, S. 2000. Biochemical characteristics of thermostable chitinase and chitin deacetylase enzymes from the Indonesian *Bacillus* K29-14. Master Thesis, Bogor Agricultural University.
- Sandford, PA. 1989. Chitosan: Commercial Uses and Potential Applications. In. Skjak-Braek *et al* (Editors). Chitin and Chitosan. Elsevier Applied Science, London.
- Shahidi, FJ, K.V. Arachchi, & J. You-Jin. 1999. Food applications of chitin and chitosans. *Trends Food Sci. & Technol.* 10: 37-51.
- Siegrist, J., & H. Kauss. 1990. Chitin deacetylase in cucumber leaves infected by *Colletotrichum lagenarium*. *Physiol. Mol. Plant Pathol.* 36:267-275.
- Somashekar, D., & R. Joseph. 1996. Chitosanases-properties and applications: a review. *Biores. Technol.* 55: 35-45.
- Srinivasan, VR. 2000. Biotransformation of chitin to chitosan. *US Patent* 5,739,015.
- Suhartono, MT. 2002. Personal Communication.
- Takayanagi, T., K. Ajisaka, Y. Takiguchi, & K. Shimahara. 1991. Isolation and characterization of thermostable chitinases from *Bacillus*



- licheniformis* X-7u. *Biochim Biophys Acta*. 1078: 404-410.
- Toharisman, A., E. Chasanah, EY. Purwani, JFL Jayanti, V. Welan, MT. Suhartono, JK. Hwang & YR Pyun. 2001. Screening of thermophilic microorganisms producing thermostable chitin deacetylase. *Indonesian Biotechnology Conference; An International Seminar and Symposium*. Yogyakarta, 2000.
- Tokuyasu, K., M. Ohnishi-Kameyama, & K. Hayashi. 1996. Purification and characterization of extracellular chitin deacetylase from *Colletotrichum lindemuthianum*. \chitin deacetylase from *Colletotrichm lindemuthianum*. *J. Biol. Chem.* 270: 26280-26291.
- Tsigos, I., A. Martinou, D. Kafetzopoulos & V. Bouriotis. 2000. Chitin deacetylases: new, versatile tools in biotechnology. *Trends Biotechnol.* 18 : 305-312.