

## CYPERACEAE OF THAILAND (EXCL. CAREX)

by

J. H. KERN\*

## SUMMARY

This is an enumeration of 174 examined species and of 11 not examined ones, in 16 genera, collected in or recorded from Thailand. The treatment is largely in the same way as in the 'Florae Siamensis Enumeratio'. Synonyms and references are given. All examined specimens are listed for each of the 7 provinces of Thailand indicated on a map, and their field notes are recorded. Distribution outside Thailand is given in some detail, and occasionally nomenclatural or other notes are added.

In *Lepironia* 1 species is recognized, in *Mapania* 5, in *Hypolytrum* 1, in *Fuirena* 2, in *Lipocarpha* 2, in *Hemicarpha* 1, in *Scirpus* 10, in *Eleocharis* 8, in *Fimbristylis* 51, in *Bulbostylis* 3, in *Cyperus* 53, in *Rhynchospora* 7, in *Triocostularia* 1, in *Schoenus* 2, in *Gahnia* 1, in *Scleria* 23.

Novelties are *Fimbristylis gracilenta* var. *psilopoda*, *Scleria biflora* ssp. *ferruginea* (Ohwi), and *Scleria reticulata* (Holt.).

An index to collectors' numbers is given at the end.

During my studies on the Cyperaceae of Malaysia for the Flora Malesiana I have also identified all materials from Thailand which came into my hands, with the exception of the species of the genus *Carex*, the elaboration of which it was agreed should be performed by Dr. M. Raymond. As most members of the sedge family are widely distributed these identifications were necessary for plant geographical reasons.

It appeared during this work that numerous species had never properly been recorded for the Thailand flora. This induced me to borrow all the cyperaceous sheets which are contained in the Herbaria of the Department of Agriculture (Curator Mr. Kasin Suvatabandhu) and of the Royal Forest Department (Curator Mr. Tem Smitinand) at Bangkok. The former herbarium consists mostly of a very large set of Kerr's collections, the latter incorporates many collections newly made by Mr. Smitinand and other members of his department. In addition sheets of Thailand Cyperaceae preserved in the Leyden, Kew, and Copenhagen Herbaria have been investigated.

\* Rijksherbarium, Leyden.

Though I have not ventured on tracing at Paris and in other herbaria all other specimens collected in Thailand, I assume that the present enumeration represents a substantial precursor towards a future comprehensive account of the *Cyperaceae* in the "Florae Siamensis Enumeratio".

In it I have evaluated former publications on Thailand *Cyperaceae* in a similar way as has been done in the "Enumeratio".

The specimens cited have been enumerated under the 7 floristic regions or provinces of Thailand as adopted by Mr. Smitinand in his study on the genus *Dipterocarpus* in Thailand (cf. Thai Forest Bull., Bot., no. 4, 1958): — Northern (N.), North-Eastern (N.E.), Eastern (E.), Central (C.), South-Eastern (S.E.), South-Western (S.W.), and Peninsular (PEN.).

I express my sincere thanks to Messrs Smitinand and Suvatabandhu for their interest and cordial co-operation in entrusting me their material and verifying the exact spelling of names of persons and localities which were sometimes difficult to copy from the herbarium labels.

Eleven names of *Cyperaceae* have been cited in literature of which I have not seen the specimens on which the records were based. These names are listed at the end of the enumeration. Some of them are probably based on erroneous identifications or are in my opinion doubtful. It falls beyond the scope of my work to try to verify them.

Finally, for reasons of facility, I have added an index to collectors' numbers.

The following abbreviations have been used for literature references:

F.B.I. — Hooker, J. D., Flora of British India 6. 1893-1894. *Cyperaceae* by C. B. Clarke.

F.I.C. — Lecomte, H., Flore Générale de l'Indo-Chine 7. 1912-1922. *Cyperacées* par E. G. Camus.

F.K.C. — Schmidt, J., Flora of Koh Chang. Contributions to the knowledge of the vegetation in the Gulf of Siam. — *Cyperaceae* by C. B. Clarke. Bot. Tidsskr. 24: 79-94. 1901.

F.M.P. — Ridley, H. N., Flora of the Malay Peninsula 5. 1925.


Hoss. 1910 — Hosseus, C. C., Beiträge zur Flora Siams. Bot. Centralbl. Beih. 27, 2: 455-507. 1910.

Hoss. 1911a — Hosseus, C. C., Beiträge zur Flora von Wang Djao am M. Ping in Mittel-Siam. Bot. Jahrb. 45: 366-374. 1911.

Hoss. 1911b — Hosseus, C. C., Die botanischen Ergebnisse meiner Expedition nach Siam. Bot. Centralbl., Beih. 28, 2: 357-457. 1911.

Kern — Kern, J. H., Florae Malesianae Precursores XXV. — Notes on Malayan and some S. E. Asian *Cyperaceae* VIII. Blumea 10: 635-651. 1960.

- Kill.—Killip, E. P., List of sedges (Cyperaceae) collected in Siam by Dr. Hugh M. Smith. Journ. Siam Soc. Nat. Hist. Suppl. 7: 55-57. 1927.
- Koy. 1955 — Koyama, T., An Enumeration of Hayata's Indo-Chinese collection of Cyperaceae: Carex and Cyperus. Naturaliste canad. 82: 194-216. 1955.
- Koy. 1957 — Koyama, T., An Enumeration of Hayata's Indo-Chinese collection of Cyperaceae. Contrib. Inst. Bot. Univ. Montréal 70: 5-64. 1957.
- Mat. F. M. P.—Ridley, H. N., Materials for a Flora of the Malayan Peninsula 3. 1907.
- Ost.—Ostenfeld, C. H., A list of plants collected in the Raheng district, Upper Siam, by Mr. E. Lindhard. Bull. Herb. Boiss., sér. 2, 5: 709-724. 1905.
- Raym.—Raymond, M., Additional notes on some S.E. Asiatic Scirpus. Naturaliste canad. 86: 225-242. 1959.
- Ridl. 1911 — Ridley, H. N., An account of a botanical expedition to Lower Siam. Journ. Str. Br. Roy. As. Soc. no 59: 27-234. 1911.
- Ridl. 1912 — Ridley, H. N., A botanical excursion to Pulau Adang. Journ. Str. Br. Roy. As. Soc. no 61: 45-65. 1912.
- Ridl. 1915 — Ridley, H. N., The plants of Koh Samui and Koh Pennan. Journ. F. M. S. Mus. 5: 158-168. 1915.
- Ridl. 1920 — Ridley, H. N., On a collection of plants from Peninsular Siam. Journ. F. M. S. Mus. 10: 65-126. 1920.
- Turr.—Turrill, W. B., in Crabb, Contributions to the Flora of Siam. Kew Bull. (1912): 421-427. 1912.
- Will.—Williams, F. N., Liste des plantes connues du Siam. Bull. Herb. Boiss., sér. 2, 4: 217-232. 1905.



### I. LEPIRONIA L. C. Rich.

1. *L. ARTICULATA* (Retz.) Domin, Bibl. Bot., Heft 85: 486. 1915. — *Ressia articulata* Retz., Obs. 4: 14. 1786. — *L. mucronata* L. C. Rich. in Pers., Syn. 1: 70. 1805; F.B.I.: 684; Mat. F.M.P.: 107; F.I.C.: 179, f. 25; F.M.P.: 175.

C. — Bangkok, cultivated: *Sarabandhu* n.n. (BK).

S.W. — Chumphawn, Bang Son: *Put* 1587 (BK); Ban Tung Maha, in marsh, near sealevel: *Kerr* 11897 (BK).

Puket, Takuapa, in large clumps in stagnant water, 10 m: *Kerr* 17108 (BK). Trang, Huay Yawt, Thale Sawng Hawn, common on sandy bank of freshwater lake, under 100 m, local name *chut*: *Smitinand* 3024 (RFD 16194) (L).

DISTR. Madagascar, Ceylon, Indo-China, S. China, Malaysia, N. and E. Australia, Caroline Islands, New Caledonia, Fiji.

The stems are used for matting.

### II. MAPANIA Aubl.

2. *M. CUSPIDATA* (Miq.) Uitt., Journ. Arn. Arb. 20: 213. 1939. — *Lepironia cuspidata* Miq., Fl. Ind. Bat., Suppl.: 603. 1861. — *M. humilis* (non F.-Vill.) F.B.I.: 683; Mat. F.M.P.: 105; F.M.P.: 174, non *Pandanophyllum humile* Steud. — *M. triquetra* Ridl., Journ. Str. Br. Roy. As. Soc. no 41: 51. 1904; Mat. F.M.P.: 106; F.M.P.: 174.

#### VARIETY CUSPIDATA

PEN. — Trang, Chum Het, in evergreen forest, 100 m: *Kerr* 15215 (BK).

Satul, Klawng Ton, very common in evergreen forest, 100—300 m: *Kerr* 14432 (BK).

var. PETIOLATA (Clarke) Uitt., Journ. Arn. Arb. 20: 213. 1939. — *M. petiolata* Clarke, Kew Bull. add. ser. 8: 54. 1908. — *M. humilis* var. *petiolata* (Clarke) Ridl., Fl. Mal. Pen. 5: 174. 1925.

PEN. — Langsuan, Ban Kragi, in evergreen forest, 200 m: *Kerr* 12005 (BK).

Pang Ngu, Kao Kathakwam, in evergreen forest, 100 m: *Kerr* 18399 (BK).

Nakawn Sritthamarat, Khao Luang, common on slope and near stream in evergreen jungle, 600 m: *Smitinand* 1028 (BKF).

Trang, Kao Soi Dao, in evergreen forest, 500 m: *Kerr* 19151 (BK).

Pattani, Betong, in evergreen forest, 200 m: *Kerr* 7505 (BK).

Satul, Adang, marshy ground in evergreen forest, 20 m: *Kerr* 15029 (BK).

DISTR. Malaysia, Solomon Islands. \*

The varieties are based on the shape of the leaf-blades and connected by many intermediates. Probably they are merely ecological forms.

3. *M. PALUSTRIS* (Hassk. ex Steud.) F.-Vill., Nov. App.: 309. 1882; F.B.I.: 681; Mat. F.M.P.: 103; F.M.P.: 172. — *Pandanophyllum palustre*

Hassk. ex Steud., Syn. 2: 134. 1855. — *M. andamanica* Clarke, Fl. Br. Ind. 6: 681. 1894; Ridl. 1920: 124. — *M. albescens* Clarke, Kew Bull. add. ser. 8: 54. 1908; F.M.P.: 172.

PEN.— Takuapa, Bangwan, common in light evergreen forest, 10 m: Kerr 17089 (BK).  
Pattani, Kao Kalakiri, common in evergreen forest, 800 m: Kerr 14993 (BK).

DISTR. From Assam to S. China and Malaysia.

*Mapania andamanica* is more slender and has smaller spikelets and flowers than typical *M. palustris*. *Mapania albescens* differs from *M. andamanica* only by the pale colour of the dried specimens. In my opinion they are not specifically distinct from *M. palustris*.

4. *M. KURZII* Clarke, Fl. Br. Ind. 6: 681. 1894; Mat. F.M.P.: 104; F.M.P.: 172. — *M. longispica* Ridl., Journ. Str. Br. Roy. As. Soc. no 44: 205. 1905; Mat. F.M.P.: 104; F.M.P.: 172. — *M. valida* Ridl., Journ. Str. Br. Roy. As. Soc. no 44: 205. 1905; Mat. F.M.P.: 104; F.M.P.: 172.

PEN.— Phang-Nga, Kao Katakwam, in evergreen forest, 200 m: Kerr 18404 (BK). Krabi, Lanta, in evergreen forest, 400 m: Kerr 19407 (BK).

DISTR. Malay Peninsula, Central Sumatra.

5. *M. ENODIS* (Miq.) Clarke, Kew Bull. add. ser. 8: 53. 1908. — *Lepironia enodis* Miq., Fl. Ind. Bat., Suppl.: 603. 1861. — *M. longa* Ridl. & Clarke, Fl. Br. Ind. 6: 683. 1894, p.p.; Mat. F.M.P.: 105; F.M.P.: 173. — *M. tumida* Uitt., Gard. Bull. S.S. 10: 182. 1939.

S.E.— Trat, Dan Chumphon, on gravel bank in stream, under 50 m: Kerr 17693 (BK). PEN.— Ranawng, Klang Kampuan, on bank of stream in evergreen forest, 100 m: Kerr 17004 (BK); same locality, 50 m: Smitinand 4161 (L).

DISTR. Malay Peninsula, Sumatra, W. Borneo.

6. *M. TENUISCAPA* Clarke, Fl. Br. Ind. 6: 683. 1894; Mat. F.M.P.: 105; p.p.; Ridl. 1911: 224; 1912: 63; F.M.P.: 173, p.p.

S.E.— Chanthabun, Kao Sabap, in evergreen forest, 200 m: Kerr 18035 (BK).

Trat, Kaw Chang, Khiawng Nonsi, by streams in evergreen jungle, 5 m: Smitinand 5682 (L).

PEN.— Ranawng, Kaw Chang, in evergreen forest, 5 m: Kerr 16595 (BK).

Takuapa, Bangwan, common in tight evergreen forest, 10 m: Kerr 17088 (BK). K.

Phang-Nga, Kao Katakwam, in dry evergreen forest, 100 m: Kerr 18500 (BK). Trang, Chawng, very common on slope in evergreen jungle, 150 m: Smitinand 4116 (RFD 16889) (L).

Satul, Klawng Ton, common in evergreen forest, 100 m: Kerr 14626 (BK, BO K); Rawei Island: Ridley 15715 (BM, K, SING).

DISTR. Indo-China, Malaysia.

### III. HYPOLYTRUM L. C. Rich.

7. *H. NEMORUM* (Vahl) Spreng., Syst. 1: 233. 1825. — *Schoenus nemorum* Vahl, Symb. Bot. 3: 8. 1794. — *H. latifolium* L. C. Rich. in Pers., Syn. 1: 70. 1805; F.B.I.: 678; F. KC.: 92; Will.: 223; Mat. F.M.P.: 100; Hoss. 1910: 457; F.I.C.: 171, f. 22, 2—8; Ridl. 1920: 124; F.M.P.: 170.

E.—Korat, Katok, by stream in evergreen forest, 400 m: Kerr #158 (BK); 500 m: Kerr 6177 (BK).

S.E.—Nawng Plu: Mrs Collins 845 (BK).

Trat, Kao Kuap, in light evergreen forest, 700—800 m: Kerr 17787 (BK); Kao Saming: Put 516 (BK); Kaw Chang, Khlawng Mayom, common by stream on sandy bank in evergreen forest, 5 m: Smitinand 2172 (RFD 11554) (BKF); Kaw Chang, Khlawng Mayom, common in evergreen forest, 100 m: Kerr 6800 (BK); Kaw Chang, Khlawng Nonsi, scattered along stream bank in evergreen jungle, 50 m: Bunnak 338 (RFD 12317) (BKF).

S.W.—Chumphawn, Kuring, in evergreen forest, 50 m: Kerr 11606 (BK); Kao Tong, in evergreen forest, 500 m: Kerr 12894 (BK).

PEN.—Ranawng, La-un, in evergreen forest, 20 m: Kerr 18494 (BK).

Langsuan, Tako: Put 1683 (BK); same locality, in evergreen forest, under 50 m, local name *tao sawn*: Kerr 11895 (BK); Kao Nom Sao, in evergreen forest, 900 m: Kerr 12038 (BK). Satul, Adang, in evergreen forest, 50 m: Kerr 14025 (BK).

DISTR. From India to Formosa, N. E. Queensland, and Polynesia.

### IV. FUIRENA Rottb.

8. *F. UMBELLATA* Rottb., Progr.: 27. 1772; F.B.I.: 666; Mat. F.M.P.: 80; F.I.C.: 141, f. 18, 6; Turr.: 425; F.M.P.: 162. — *F. uncinata* (non Kunth) F.I.C.: 141.

C.—Chalbadan, Dong Pya Yen, in marshy savannah, 100 m: Kerr 8008 (BK).

S.E.—Chanburi, Makham, Ban Saenghwang, common in wet localities, 140 m: Bunnak 165 (L.).

Prachinburi, Aran Prathet: Put 3093 (BK); Watana: Put 1952 (BK).

S.W.—Chumphawn, Bang Son: Put 1593 (BK).

PEN.—Ranawng, Kaw Chang, at edge of ditch in *Melaleuca* forest, 5 m: Kerr 16554 (BK).

Satul, Kuan Po, on edge of stream, 20 m: Kerr 13788 (BK).

DISTR. Tropical and subtropical regions of the whole world.

9. *F. CILIARIS* (L.) Roxb., Fl. Ind. 1: 184. 1820. — *Scirpus ciliaris* L., Mant. 2: 182. 1771. — *F. glomerata* Lamk, Tabl. Enc. Méth. Bot. 1: 150. 1791; F.B.I.: 666; F. KC.: 90; Will.: 224; Mat. F.M.P.: 81; Hoss. 1910: 460; Ridl. 1911: 224; F.I.C.: 140; F.M.P.: 163. — *F. wallichiana* (non Kunth) F.I.C.: 141.

S.E. — Prachinburi, Nah Kali, 30 ft: *Mrs. Collins* 1902 (BK).

C. — Bangkok, along railway trail, under 5 m: *Kerr* 6720 (BK); Bangkok, Bangkhen, RFD Compound, near water, 5 m: *Smitinand* 5419 (RFD 19728) (L); Bangkok, Klong side: *Ergl Smith* 225 (BK).

Dhonburee: *Thasnee Boonkong* 2 (BK).

PEN. — Koh Samui, Ban Bua Put, in scrub, under 5 m: *Kerr* 15711 (BK).

DISTR. Tropical Africa, S.E. and E. Asia, Malaysia, tropical Australia.

### V. LIPOCARPHA R. Br.

10. *L. CHINENSIS* (Osb.) Kern, Blumea, Suppl. 4: 167. 1958. — *Scirpus chinensis* Osb., Dagb. Ostind. Resa: 220. 1757. — *Scirpus senegalensis* Lamk. Tabl. Enc. Méth. Bot. 1: 140. 1791. — *Hypaelyptum argenteum* Vahl, En. 2: 283. 1806. — *L. argentea* (Vahl) R. Br. ex Nees, Linnaea 9: 287. 1835; F.B.I.: 667; Mat. F.M.P.: 81; Hoss. 1910: 457; 1911b: 370; F.I.C.: 143, t. 1, f. A, 1—2; Turr.: 425; F.M.P.: 163. — *L. senegalensis* (Lamk) Th. & Hél. Durand, Syll. Fl. Congol.: 619. 1909; Koy. 1957: 53. — *L. triceps* (non Nees) F.I.C.: 144.

N. — Chiangmai, Doi Suthep, common by stream in swampy ground, 1100 m: *Smitinand* 181 (RFD 6179) (L); 700 m, *Hosseus* 481 (L).

N.E. — Loei, Phu Kradang, in open pine forest, 1300 m: *Dee* 37 (RFD 4906) (L).

PEN. — Trang, Sikao, on low-lying open ground, near sealevel: *Kerr* 19063 (BK).

DISTR. Tropical and South Africa, Madagascar, Mascarene Is., S. and S.E. Asia, Malaysia, tropical Australia.

11. *L. SPHAELATA* (Vahl) Kunth, En. 2: 267. 1837; F.B.I.: 667. — *Hypaelyptum sphacelatum* Vahl, En. 2: 283. 1806.

E. — Korat, Bua Yai: *Put* 4294 (BK).

S.W. — Kanburi, scattered in wet locality, 50 m: *Smitinand* 3859 (RFD 18091) (BKF).

DISTR. From Nepal to Ceylon and Thailand, also recorded from tropical Africa and America.

12. *L. MICROCEPHALA* (R. Br.) Kunth, En. 2: 268. 1837; F.B.I.: 668; Mat. F.M.P.: 82, p.p.; F.I.C.: 144; F.M.P.: 163, p.p. — *Hypaelyptum microcephalum* R. Br., Prodr.: 220. 1810.

S.E. — Prachinburi, Aran Prathet, in open low-lying grassy ground, 50 m: *Kerr* 19568 (BK).

DISTR. S.E. and E. Asia, Malaysia, Australia.

### VI. HEMICARPHA Nees

13. *H. ISOLEPIS* Nees, Edinb. New Phil. Journ. 17: 263. 1834; in Wight, Contr. Bot. Ind.: 92. 1834. — *Scirpus isolepis* (Nees) Boeck., Linnaea 36: 498. 1870; F.B.I.: 663.

E. — Korat, Chan Tuk, among low herbage on open ground, 300 m: *Kerr* 8064 (BK).

DISTR. Tropical Africa, India.

VII. *SCIRPUS* L.

14. *S. grossus* L. f., Suppl.: 104. 1781; F.B.I.: 659; Ridl. 1911: 222; F.I.C.: 136; Turr.: 425; F.M.P.: 162; Koy. 1957: 60; Raym.: 227.  
 C.—Dhonburi, Bang Plat (Bangkok noi), common in wet and inundated area, under 10 m, local name *kok*: *Smitinand 2797* (BKF).  
 Bangkok, along ditch, local name *kok tu krap*: *Kerr 10714* (BK).  
 S.E.—Lemhabang-swamp near Sriracha, c. 20 ft: *Mrs Collins 1896* (BK).  
 DISTR. From India to S. China and tropical Australia.

"Chantabun mats are said to be made from the stems of this species" (Collins).

15. *S. affinis* Roth in R. & S., Syst. Veg. 2: 140. 1817; F.I.C.: 137. — *S. maritimus* L. var. *affinis* (Roth) Clarke, Fl. Br. Ind. 6: 659. 1893.  
 N.E.—Nakawn Panom, Tat Panom, in damp ground at edge of river, 200 m: *Kerr 21409* (BK).

DISTR. Widely distributed in Central and S.E. Asia.

Very near to *S. maritimus* L. and possibly a subspecies of this widespread, polymorphic species.

16. *S. WICHURAI* Boeck., Linnaea 36: 729. 1870; Raym.: 226. — *S. eriophorum* (non Michx) F.B.I.: 661.  
 N.E.—Loei, Phu Krading, not common by stream in evergreen jungle, 1300 m: *Smitinand 1974* (RFD 9459) (BKF).

DISTR. From India to S. China, Korea, Japan, and Central Sumatra.

17. *S. LITORALIS* Schrad., Fl. Germ. 1: 142, t. 5, f. 7. 1806; F.B.I.: 659.  
 PEN.—Patalung, Kaw Si Kaw Ha, in water about 1 m deep, in inland sea (water nearly fresh), very common along shore of limestone islets: *Kerr 15142* (BK).  
 DISTR. From the Mediterranean region through S. Asia (here very local) to Australia.

18. *S. MUCRONATUS* L., Sp. Pl. 1: 50. 1753; F.B.I.: 657; Mat. F.M.P.: 79; F.I.C.: 134; Turr.: 425; F.M.P.: 161; Koy. 1957: 59; Raym.: 232.  
 N.—Chiengrai, in swamp, 450 m: *Winit 777* (BKF).  
 S.W.—Chumphawn, Ban Tung Maha, in marshy ground, near sealevel: *Kerr 11503* (BK).  
 Pattani, Yala, in open marshy ground, under 50 m: *Kerr 7261* (BK).  
 PEN.—Nakawn Srithamarat, Tung Song, in rice-field, local name *chut mu*: *Rabil 149* (BK).  
 DISTR. From S. Europe through S. Asia to Japan and Australia.

19. *S. JUNCOIDES* Roxb., Fl. Ind. 1: 218. 1820; Raym.: 236. — *S. erectus* (non Poir.) F.B.I.: 656; Mat. F.M.P.: 79; Ridl. 1911: 222; F.I.C.: 136; F.M.P.: 161.

S.E. — Chanburi, Makhram, Thung Mala, common in wet localities, 130 m: *Bunnak* 19 (RFD 9609) (L).

Trat, Huay Raeng, Dong Madune, common in rice-field, under 50 m, local name *phrong klom yai*: Smitinand 1402 (RFD 8777) (BKF); Kaw Chang, Khlaem Nonsi, in flooded ricefields, under 10 m: Kerr 9198 (BK).

S.W. — Near Neekey, near Wangka, along pools, 150 m: den Hoed 229 (Krene No R. basis exp. 1948) (BK, L).

PEN. — Ranawng, Kraburi, on edge of rice-field, 10 m: Kerr 16340 (BK); Lam Lien in stagnant water, 10 m: Kerr 16382 (BK); Nam Chut, in marshy ground, near sealevel, local name *chuk nn*: Kerr 11700 (BK).

DISTR. Madagascar; from India, China, and Japan to tropical Australia.

20. *S. ARTICULATUS* L., Sp. Pl. 1: 47. 1753; F.B.I.: 656; F.I.C.: 133 Raym.: 233.

C. — Bangkok, in rice-fields, under 5 m: Kerr 6740, 11100 (BK); Bangkok, Bangkhen RFD Compound, near water, under 10 m: Smitinand 5420 (RFD 19729) (L).

S.W. — Rajaburi, Potaram, in paddy fields: Kerr 3877 (BK).

DISTR. From the Mediterranean region through Africa and S. Asia to tropical Australia.

21. *S. LATERIFLORUS* Gmel., Syst. Veg. 1: 127. 1791; Raym.: 234. — *S. lateralis* Retz., Obs. 4: 12. 1786, non Forsk. 1775. — *S. supinus* L. var. *uninodis* Clarke, Fl. Br. Ind. 6: 656. 1893, quoad specim. cit., non *Isolepis un nodis* Delile. — *S. supinus* (non L.) Mat. F.M.P.: 79; F.I.C.: 135; F.M.P.: 161 N. — Chiengmai, Mae Taeng, in rice-field, 300 m: Smitinand 3928 (L).

C. — Bangkok, in flooded rice-fields: Kerr 3860 (BK); along railway line: Kerr 672 (BK); Bangkhen, Bangkok, RFD Compound, near water, under 10 m: Smitinand 5418 (RFD 19737) (L).

DISTR. From India to S. China, Formosa, and Australia.

22. *S. ANNAMICUS* Raymond, Naturaliste canad. 84: 137. 1957; Raym. 239.

PEN. — Kukan, Hui Nua, open ground by roadside, 100 m: Kerr 8323 (BK, K).

DISTR. Annam (type locality).

23. *S. SQUARROSUS* L., Mant.: 181. 1771; F.B.I.: 663; F.I.C.: 134; Turr. 425; Raym.: 228.

S.W. — Kanburi, common in wet ground, 50 m: Smitinand 3861 (RFD 16092) (L).

DISTR. S. Asia, extending to Singapore Island.

24. *S. SIAMENSIS* (Clarke) Kern, Blumea 9: 219. 1958; Raym.: 230. — *S. squarrosus* L. var. *siamensis* Clarke in Hosseus, Beitr. Fl. Siams, Beih Bot. Centralbl. 27: 460. 1910; Hoss. 1911a: 372; 1911b: 371; F.I.C. 134. — *Lipocarpha tenera* (non Boeck.) F.I.C.: 148. — *Ascolepis* sp. n., Turr.: 425

- *Scirpus chinensis* Osb. var. *siamensis* (Clarke) Raym., Naturaliste canad. 84: 124. 1957.  
 N.—Nakawn Sawan, Wang Djao, savannah forest, common, 100 m: *Hosseus* 161 (P, type).  
 DISTR. Laos, Tonkin, Cochinchina.

### VIII. ELEOCHARIS R. Br.

25. *E. congesta* D. Don, Prodr. Fl. Nepal.: 41. 1825; F.B.I.: 630; F.I.C.: 88; Koy. 1957: 37. — *E. pellucida* Presl, Rel. Haenk. 1: 196. 1828; Koy. 1957: 37. — *E. afflata* Steud., Syn. 2: 76. 1855; F.B.I.: 629; F.I.C.: 87, f. 13, 5 — *E. subvivipara* Boeck., Linnaea 36: 424. 1870; F.B.I.: 629.  
 N.E.—Loei, Phu Kradang, Tham Saw, common in swampy ground, under water, 1300 m: Smitinand 1123 (RFD 7865).  
 E.—Chaiyaphum, Po Kio, edge of pool, 1000 m: Kerr 20219 (BK).  
 DISTR. S.E. Asia, from India to China, Japan, and Malaysia.

*Eleocharis congesta* and *E. pellucida* are treated as specifically distinct by Svenson, the monographer of the genus. He distinguishes between them only by the thicker rigid stems of the former. I think it is impossible to trace a dividing line between the two.

26. *E. RETROFLEXA* (Poir.) Urb., Symb. Ant. 2: 165. 1900. — *Scirpus retroflexus* Poir. in Lamk, Enc. 6: 753. 1804. — *E. chaetaria* R. & S., Syst. Veg. 2: 154. 1817; F.B.I.: 629; F. KC.: 84; Willd.: 224; Mat. F.M.P.: 77; Hoss 1910: 460; F.I.C.: 87, f. 12, 7—8; F.M.P.: 151.  
 S.E.—Chantabun: Venterdal 463 (SING).  
 PEN.—Ranawng, Nam Chut, on mud at edge of marshy ground, near sealevel: Kerr 11707 (BK).  
 Pattani, Kok Po, on damp sandy soil by stream, near sealevel: Kerr 7086 (BK).  
 DISTR. Pantropic.

To me the differences between *E. retroflexa* (the New World 'species') and *E. chaetaria* (the Old World one) are too trifling to justify specific separation.

27. *E. ACUTANGULA* (Roxb.) Schult. in R. & S., Mant. 2: 91. 1824. — *Scirpus fistulosus* Poir., Enc. 6: 749. 1804, non Forsk. 1775. — *Scirpus acutangulus* Roxb., Fl. Ind. 1: 216. 1820. — *E. fistulosa* Schult. in R. & S., Mant. 2: 89. 1824; F.B.I.: 626; Mat. F.M.P.: 76; Ridl. 1911: 222; F.I.C.: 84, f. 13, 3—4; F.M.P.: 151.  
 S.E.—Trat, Huay Raeng, Dong Maduae, common in rice-field, under 50 m: Smitinand 1191 (RFD 6776) (L).  
 PEN.—Songkhla, Chana, in open fields, under 20 m: Kerr 15798 (BK).  
 Pattani, Saiburi, paddy fields, 25 m: Kerr 7231 (BK).

DISTR. Widely distributed but uncommon species in the tropics of the whole world.

28. E. SPIRALIS (Rottb.) R. & S., Syst. Veg. 2: 155. 1817; F.B.I.: 627; F.I.C.: 85, f. 11, 6—7. — *Scirpus spiralis* Rottb., Descr. & Ic.: 45, t. 15, f. 1. 1773.

PEN.—Satul, in marshy ground: *Kerr 13740* (BK).

DISTR. Tropical Africa, Madagascar, Mauritius, Ceylon, S. India, S. China, Malaysia, tropical Australia, New Caledonia.

29. E. PHILIPPINENSIS Svens., Rhodora 31: 155, f. 9. 1929; Kern, Reinw. 4: 94. 1956.

C.—Bangkok, open, low lying, marshy ground: *Kerr 11105* (BK, BO, K, L).

DISTR. Tropical Australia, New Caledonia, Malaysia, Hainan.

30. E. OCHROSTACHYS Steud., Syn. 2: 80. 1855; F.B.I.: 626; Mat. F.M.P.: 76; F.M.P.: 151. — *Scirpus laxiflorus* Thwaites, En. Pl. Zeyl.: 435. 1864. — *E. variegata* (Poir.) Presl var. *laxiflora* (Thwaites) Clarke, Fl. Br. Ind. 6: 626. 1893; Turr.: 423.

S.E.—Chanthaburi, Makhram, common in wet localities in scrub forest, 160 m: *Smitinand 4056* (L).

Trat, Kao Kuap, in damp ground by stream, 600 m: *Kerr 17753* (BK).

PEN.—Phang-Nga, Kaw Yao Yai, in open marshy ground, 5 m: *Kerr 17355* (BK); Nakawn Srithamarat, Tha Samet, under 50 m: *Kerr 14309* (BK).

Trang, Kantang, common in swamp, under 50 m: *Smitinand 2952* (RFD 12612) (L); Huay Yawt, Thale Sawng Hawng, sandy bank of fresh water lake, 100 m: *Smitinand 3026* (RFD 17248) (L).

DISTR. From India and Ceylon to Formosa, the Riu Kiu Islands, Malaysia, Micronesia, and Melanesia.

31. E. DULCIS (Burm. f.) Henschel, Vita Rumph.: 186. 1833. — *Andropogon dulce* Burm. f., Fl. Ind.: 219. 1768. — *Scirpus plantagineus* Retz, Obs. 5: 14. 1789. — *E. plantaginea* (Retz.) R. & S., Syst. 2: 150. 1817; F.B.I.: 625; Mat. F.M.P.: 75; F.I.C.: 82; F.M.P.: 150. — *E. equisetina* Presl, Rel. Haenk. 1: 195. 1828; F.B.I.: 626; F. KC.: 85; Will.: 224; Mat. F.M.P.: 75; Hoss. 1910: 460; Ridl. 1911: 222; F.I.C.: 83; Turr.: 423; F.M.P.: 150.

C.—Bangkok, in marsh, local name *ya-chong-katiam*: *Boonbrahm* 8 (BK); in shallow ditch: *Kerr 7083* (BK); edge of flooded rice-fields: *Kerr 12526* (BK); rice-field, 40 m: *Ergyl Smith 931* (BK).

S.E.—Near Sriracha, 30 ft: *Mrs Colling 1893* (BK).

PEN.—Surat, in shallow pool, under 10 m: *Kerr 11289* (BK).

Satul, B. Rayak: *Ridley 14809* (BM, SING).

DISTR. Tropical West Africa, Madagascar; from India to S. China, Formosa, Riu Kiu Islands, and Japan, Malaysia, Micronesia, Melanesia, tropical Australia.

32. E. GENICULATA (L.) R. & S., Syst. Veg. 2: 150. 1817. — *Scirpus geniculatus* L., Sp. Pl. 1: 48. 1753. — *Scirpus caribaeus* Rottb., Progr.: 24. 1772. — *E. capitata* R. Br., Prodr.: 225. 1810; F.B.I.: 627; F. KC.: 84; Will.: 224; Mat. F.M.P.: 77; Hoss. 1910: 460; F.I.C.: 85; F.M.P.: 151. — *E. caribaea* (Rottb.) S. F. Blake, Rhodora 20: 24. 1918; Koy. 1957: 37.

S.E.—Trat, Kaw Chang, Lem Dan Kao, in marshy ground by sea, 5 m; Kerr 8236 (BK); Kaw Chang, Khlawng Mayom, common in damp localities in waste ground, 5 m; Smitinand 2227 (RFD 11540) (L); Kaw Kut, Khlawng Nonsi, by paths in sandy soil, 5 m; Smitinand 5648 (L).

S.W.—Chumphawn, Bang Son; Put 1544 (BK).

DISTR. Warmer parts of the whole world.

In a letter to Svenson, published in Rhodora 41: 50. 1939, Dandy pointed out that the correct name for this species under *Eleocharis* is *E. geniculata* (L.) R. & S. In a later publication (N. Am. Fl. 18, 9: 520. 1957) Svenson reverts to the binomial *E. caribaea*, as (l.c. p. 533) "the name *Scirpus geniculatus* included *E. caribaea* and the plant known as *Juncus aquaticus geniculatus, capitulis equiseti, major Sloane*" and "in the second edition of Species plantarum (1762) Linnaeus restricted the name *Scirpus geniculatus* to the latter." I cannot agree with him. The references in Sp. Pl. ed. 2 are literally the same as in the first edition. The fact that in his diagnosis Linné changed "spica subglobosa" into "spica oblonga" does not affect the choice of the specimen in the Hortus Siccus Cliffortianus as the lectotype of *Scirpus geniculatus*.

#### IX. FIMBRISTYLIS Vahl

33. F. THOMSONII Boeck., Linnaea 37: 37. 1871; F.B.I.: 646; F.I.C.: 117; Kern, Blumea 8: 110. 1955. — *F. asperrima* (non Boeck.) Ridl. 1911: 222; 1912: 63; F.M.P.: 158, p.p. (quoad specim. P. Rawei).

N.—Chiengmai, Doi Suthep, in the shade in oak forest, 1100 m; Smitinand 123 (RFD 6700) (BKF).

Lampang, Khun Tan, scattered in evergreen forest, 1500 m, local name *yak haew mu*; Put 153 (BKF).

Petchabun, Lomkao, Phu Hee, common in oak forest, 860 m; Smitinand 2591 (L). N.E.—Loei, Phu Krading, Sam Khae, common in open ground, 1100 m; Smitinand 1187 (RFD 7664) (BKF); Sithan, common in open deciduous forest, 300 m; Phloenachit 1293 (RFD 17006) (L).

E.—Chaiyaphum, Pu Kio, in open grassy ground, 700 m; Kerr 29200 (BK).

S.W.—Kanburi, Pu Keng, in deciduous forest, 100 m; Kerr 20858 (BK, K).

Bangtaphan; Keith 388 (SING).

Chumphawn, common in the plain; Khanthachai 1637 (L).

PEN.—Nakawn Srithamarat, in savannah, 100 m; Kerr 13624 (BK, K).

Rawei Island; Ridley 15729 (BM, K, SING).

DISTR. S. and S.E. Asia, extending to Central Sumatra and the Philippines.

Rhizome locally used against kidney's trouble (*Put*).

34. *F. COMPLANATA* (Retz.) Link, Hort. Berol. 1: 292. 1827; F.B.I.: 646; Mat. F.M.P.: 96; F.I.C.: 116; F.M.P.: 158. — *Scirpus complanatus* Retz Obs. 5: 14. 1789.

E. — Korat, Bua Yai; *Put* 4297 (BK).

C. — Chaibadan, Dong Pya, in marshy savannah, 100 m: *Kerr* 8004 (BK).

S.W. — 10 km S.E. of Kin Sayok, grassy plain, gravel, lateritic soil, 100—150 m: *Kostermans* 1132 (*Kiewae Noi R. basin exp. 1946*) (L).

Chumphawn, Kao Tong, in open scrub, 700 m: *Kerr* 11539 (BK, K).

DISTR. Pantropic.

35. *F. DURA* (Zoll. & Mor.) Merr., Philip. Journ. Sc. 11: Bot. 53. 1918 — *Isolepis dura* Zoll. & Mor. in Mor., Syst. Verz.: 97. 1846. — *F. aspera* Boeck., Linnaea 37: 40. 1871; F.B.I.: 643; Mat. F.M.P.: 96; F.I.C.: 116; F.M.P.: 158, excl. specim. P. Rawei.

E. — Korat, Katok, by stream in evergreen forest, 400 m: *Kerr* 8180 (BK).

S.E. — Krabin, Sakeo, in evergreen forest, 50 m: *Kerr* 9767 (BK).

Chanthaburi, Makhram, Thung Khui, common in wet localities, 130 m: *Bunnak* 20 (RFD 9623) (L).

Trat, Kaw Chang, Khlawng Mayom, in rock crevices in stream bed, 50 m: *Kerr* 6815 (BK); Kaw Kut, by the sea in coconut plantations, 5 m: *Smitinand* 5705 (L). S.W. — Bangtaphan: *Keith* 482 (SING).

PEN. — Surat, Kanchanadit, in evergreen forest, 5 m: *Kerr* 13061 (BK, K); Bangpakong, common in evergreen jungle, under 50 m: *Phloenheit* 852 (RFD 12833).

Trang, Chawng, in rocky ground by stream, 100 m: *Kerr* 15158 (BK), *Smitinand* 4083 (RFD 15866) (L).

Pattani, Banang Sta, in open scrub, 50 m: *Kerr* 7310 (BK).

Pulau Rawei: *Ridley* 15719 (BM, K, SING).

DISTR. From India to Indo-China and W. Malaysia.

36. *F. INSIGNIS* Thwaites, En. Pl. Zeyl.: 349. 1864; F.B.I.: 645; F.I.C.: 118; Kerr, Blumea 8: 112. 1955. — *F. longispica* (non Steud.) F.M.P.: 156, p.p.

N.E. — Loei, Phu Kradeng, Rest House, in savannah, 1300 m: *Smitinand* 5889 (L).

E. — Surin, in open deciduous forest, 100 m: *Kerr* 8231 (BK).

C. — Nakhawn Nayok, Khao Lon, on rock's debris: *Sucatabantu* s.n. (BK).

S.E. — Krabin, Aranya, in savannah, under 50 m: *Kerr* 19336 (BK).

PEN. — Trang, Bangsak, in open sandy ground near sea: *Kerr* 19950 (BK).

Songkhla, Kampangpet, in savannah, under 50 m: *Kerr* 15977 (BK).

Pattani, on open sandy ground, near sealevel: *Kerr* 7741 (BK, SING).

Satul, heaths: *Ridley* 14820 (BM, K, SING); in open grassy ground, 5 m: *Kerr* 13710 (BK); Adang, in rock crevices, open rocky ground, 300 m: *Kerr* 14107 (BK).

DISTR. Ceylon, Tonkin, Cochinchina, Laos, S. China, Malaysia, tropical Australia (Queensland).

37. *F. FALCATA* (Vahl) Kunth, En. 2: 239. 1837; Kern: 645. — *Scirpus falcatus* Vahl, En. 2: 275. 1806. — *F. junciformis* Kunth, En. 2: 239. 1837; F.B.I.: 647; F.I.C.: 119.

S.W. — Kanbur, Ta Salao, in open scrub, 50 m: Kerr 19524 (BK, K).

DISTR. Madagascar, S. Asia, Malaysia.

38. *F. CYMOSA* R. Br., Prodr.: 228. 1810; F. KC.: 89; Will.: 224; Hoss. 1910: 460; F.I.C.: 119. — *F. spathacea* Roth, Nov. Pl. Sp.: 24. 1821; F.B.I.: 640; F. KC.: 87; Will.: 225; Mat. F.M.P.: 93; Hoss. 1910: 462; Ridl. 1911: 222; F.I.C.: 109; F.M.P.: 156; Kill.: 56.

S.E. — Chanthaburi, Lem Sing, Nai Noe 137 (BK, K).

Trat, Kaw Chang, Khlawng Nonsi, common in sandy soil, 5 m: Smitinand 5652 (L); Koh Kahdat, on sandy sea-shores: Joh. Schmidt 572 (C).

S.W. — Prachuap, Sam Roi Yawt, open grassy ground near sea, under 50 m: Kerr 10973 (BK).

PEN. — Surat, Kaw Tao, on rocky ground near sea: Kerr 13010 (BK); Kaw Samui, on sandy ground in coconut plantation, near sealevel: Kerr 12622 (BK, K); Kaw Samui, Ban Bua Put, in sandy ground near sea: Kerr 15699 (BK, K); same locality, in open scrub, under 5 m: Kerr 15700 (BK, K).

Krabi, Koh Lanta, on sandy ground in open *Casuarina* grove, by sea: Kerr 18872 (BK).

DISTR. Pantropic.

39. *F. SERICEA* R. Br., Prodr.: 228. 1810; F.B.I.: 641; F. KC.: 86; Mat. F.M.P.: 94; Hoss. 1910: 462; F.I.C.: 111; Turr.: 425; F.M.P.: 157; Kill.: 56. — *F. junciformis* var. *latifolia* (*non* Clarke) F.I.C.: 120.

C. — Bangkok, strand: Marean 2433 (SING).

S.W. — Prachuap, on sandy sea shore: Kerr 10729 (BK).

Songkhla, in open sandy ground near sea: Kerr 15081 (BK); Songkhla, Tepa, in sand of beach: Kerr 14881 (BK).

DISTR. S. and S. E. Asia, through Malaysia to Australia.

40. *F. OBTUSATA* (Clarke) Ridl., Fl. Mal. Pen. 5: 157. 1925; Kern, Blumea 8: 116. 1955. — *F. tenera* R. & S. var. *?obtusata* Clarke, Fl. Br. Ind. 6: 642. 1893; Mat. F.M.P.: 94. — *F. nigrobrunnea* Thwaites var. *thorelli* Camus, Not. Syst. 1: 248. 1910; F.I.C.: 121, p.p.

PEN. — Surat, Kantuli: Put 4297 (BK).

DISTR. Lower Bengal, Lower Birma, N. W. Malaysia.

41. *F. GLOBULOSA* (Retz.) Kunth, En. 2: 231. 1837; F.B.I.: 644; Mat. F.M.P.: 95; F.I.C.: 424; Turr.: 424; F.M.P.: 158; Kill.: 56; Koy. 1957: 45. — *Scirpus globulosus* Retz., Obs. 6: 19. 1791.

- N.—Phetchabun, Lomkao, Phu Hee, common in moist localities in deciduous forest 860 m: *Smitinand 2609* (L).  
 S.W.—Chumphawn, Bang Son: *Put 1509* (BK).  
 PEN.—Surat, Kantuli: *Put 4128* (BK); Praseng, abundant in open marshy ground 50 m: *Kerr 12490* (BK).  
 Trang, Kantang, common in swamp, under 50 m: *Smitinand 2951* (RFD 1200) (L).  
 Songkhla, Tepa, in wet ground, under 50 m: *Kerr 15713* (BK).  
 Pattani, Bukit, in open marshy ground, under 50 m: *Kerr 7184* (BK); Bata Sta, on dried mud in open ground, 50 m: *Kerr 7308* (BK).  
 Narathiwat, Sungai Padi, Taloh Bukeh, common in swampy ground, 120 m: *Williams 17237* (RFD 11912) (BKF), *17227* (RFD 11913) (BKF).  
 DISTR. S. and S. E. Asia, Micronesia, Polynesia.

42. F. MILIACEA (L.) Vahl, En. 2: 287. 1806; F.B.I.: 644; F. KC.: 8 Will.: 225; Mat. F.M.P.: 95; Hoss. 1910: 462; Ridl. 1911: 222; F.I.C.: 11 Turr.: 424; F.M.P.: 158. — *Scirpus miliaceus* L., Syst. Veg., ed. 10: 88. 1759.

- N.—Chainat, Manorom, common in rice-fields, under 50 m: *Kerr 19074* (BK).  
 E.—Korat, Ban Chumsaeng, paddy field: *Noi Noe 196* (BK).  
 C.—Bangkok, along railway line, under 5 m: *Kerr 6721* (BK); at edge of the local name *ya nuat pladuk*: *Kerr 10712* (BK).  
 S.E.—Chanburi, Makham, Ban Ang, common in wet localities, 120 m: *Bannak 212* (L).  
 S.W.—Near Neekey, near Wangka, very common near pools, 150 m: *den Hoed 120* (*Kwae Noi R. basin exp. 1946*) (BK, L); marshy plain near Ku Jar, c. 150 m NW of Kanburi, 100—150 m: *Kostermans 1321* (*Kwae Noi R. basin exp. 1946*).  
 PEN.—Surat, Bangbao, common by path, 50 m: *Phloenchtit 538* (RFD 14435) (L); *Smitinand 2830* (RFD 12864) (BKF).  
 Trang, Chawng, common by stream, 100 m: *Smitinand 4082* (RFD 16867).  
 Pattani, Saihuri, in paddy fields, under 25 m: *Kerr 7230* (BK).  
 Satul, heaths: *Ridley 14921* (BM, SING).  
 DISTR. Pantropic.

43. F. SALBUNDIA (Nees) Kunth, En. 2: 230. 1837; F.B.I.: 646.  
 N.E.—Loei, Phu Kradang, source of Nam Phawng, partly submerged: *Smitinand 5854* (RFD 21490) (L).  
 DISTR. India, Burma.

44. F. ANISOCLADA Ohwi, Blumea 8: 97, f. 2. 1955; Kern, Blumea 120, 1955; Koy. 1957: 47.  
 S.E.—Trat, Dan Chumpon, in open, muddy ground, under 50 m: *Kerr 17652* (BKF).  
 DISTR. Indo-China, Java, Celebes.

45. F. PAUCIFLORA R. Br., Prodr.: 225. 1810; F.B.I.: 633; Mat. F.M.P.: 90; F.I.C.: 98; F.M.P.: 154.

S.E. — Chanburi, Makhram, Ban Ang, common in wet localities in savannah, 120 m; local name *yak praeng mu*: *Bunnak* 185 (L).

Trat, Kaw Chang, Khlawng Nonsi, common by paths in sandy soil, 5 m: *Smitinand* 5617 (L).

PEN. — Pattani, open sandy ground, near sealevel: *Kerr* 7739 (BK).

DISTR. S. Asia, Malaysia, tropical Australia.

46. *F. LEPTOCLADA* Benth., Fl. Hongk.: 393. 1861; F.B.I.: 647; Mat. F.M.P.: 96; Ridl. 1911: 222; F.I.C.: 118; F.M.P.: 159. — *F. disticha* (non Boeck.) F.I.C.: 125.

PEN. — Nakawn Srithamarat, Padang Besar, in light evergreen forest, 50 m: *Kerr* 3594 (BK).

Satul, heaths: *Ridley* 14819 (BM, K, SING).

DISTR. Ceylon, Indo-China, S. China, Malaysia.

47. *F. MONOSTACHYOS* (L.) Hassk., Pl. Jav. Rar.: 61. 1848; F.B.I.: 649; Ridl. 1911: 224; F.I.C.: 122; Turr.: 424; F.M.P.: 159. — *Cyperus monostachyos* L., Mant. 2: 180. 1771.

N. — Chiangmai, Muang Payao: *Put* 3987 (BK).

N.E. — Loei, Sithan, common in deciduous forest, 300 m: *Smitinand* 4792 (L).

Nakawn Panom, Dawn Tan, in open deciduous forest, 100 m: *Kerr* 21528 (BK).

E. — Korat, Bua Yai: *Put* 4270 (BK).

C. — Saraburi, Menam Sak, mixed deciduous forest, 40 m: *Kerr* 7022 (BK).

S.W. — Kanburi, Ta Salac, in bamboo forest, 50 m: *Kerr* 19473 (BK).

Prachuap: *Put* 289 (BK); Prachuap, Hui Yang: *Put* 3170 (BK); Sam Roi Yawt, common in open scrub on sandy ground, under 50 m: *Kerr* 10959 (BK).

PEN. — Surat, Kanchanadit, in wet ground by stream, evergreen forest, 5 m: *Kerr* 12225 (BK); Kaw Samui, on sandy ground in coconut plantation, near sealevel: *Kerr* 12584 (BK).

DISTR. Pantropic.

48. *F. ERAGROSTIS* (Nees) Hance, Journ. Linn. Soc., Bot. 13: 132. 1873; Turr.: 424; Kov. 1957: 38. — *Abildgaardia eragrostis* Nees & Mey. ex Nees in Wight, Contr. Bot. Ind.: 95. 1834. — *F. tortispica* Turr., Kew Bull. (1911): 348. 1911; Turr.: 425; F.I.C.: 125.

S.E. — Loei, Phu Kradung, common in savannah on rocks, 1300 m: *Smitinand* 1948 (L); Wang Saphung, Sithan, common along edge of paddy field, 300 m: *Dee 3* (RFD 5653) (BKF); Kao Kradung, open grassy ground, 1200 m: *Kerr* 8684 (BK).

Sukon, Wa Nawn, in open grassy ground, 200 m: *Kerr* 8477 (BK).

S.E. — Trat, Kao Kuap, in light evergreen forest, 600 m: *Kerr* 17750 (BK).

S.W. — Kanburi, Kao Tawng, in open, grassy, deciduous forest, 100 m: *Kerr* 19776 (BK, K).

Rathuri, common in open deciduous forest, under 50 m: *Kerr* 9037 (BK, K).

Langsuan, Tako: *Put* 1668 (BK).

Gorat, open spaces in scrub, under 10 m: *Kerr* 11251 (BK, K); Kantuli: *Put* 4170 (BK).

Songkhla, Ban Prakawp, on sandy soil in light evergreen forest, 50 m: *I*  
15828 (BK, K).

DISTR. S. Asia, extending northwards to S. China and Formosa, southwards to Queensland.

49. *F. CINNAMOMETORUM* (Vahl) Kunth, En. 2: 229. 1837; Kern: 6.  
— *Scirpus cinnamometorum* Vahl, En. 2: 278. 1806. — *F. cyperoides* R. Br. Prod.: 228. 1810; F.B.I.: 650. — *F. cyperoides* R. Br. var. *cinnamometorum* (Vahl) Clarke, Fl. Br. Ind. 6: 650. 1893; Turr.: 423.  
N. — Phetchabun, Lamkao, Phu Hee, common in oak forest, 860 m: *Smitinand* 2595.  
N.E. — Loei, Phu Krating, Rest House, in savannahs, 1300 m: *Smitinand* 5948.  
Nakawn Panom, Muk Dahan, in open deciduous forest, 200 m: *Kerr* 21451 (B).  
E. — Srisaket, Kanthraram, in open scrub jungle near village, 200 m: *Smitinand* (RFD 4938) (L).  
S.E. — Chanburi, Makham, along edge of thickets, 20 m: *Smitinand* 5757 (L).  
Trat, Huay Raeng, Dong Maduea, common among grasses in rice-field, 50 m: *Smitinand* 1381 (RFD 6772) (BKF).  
S.W. — Petchaburi, Tung Luang, in open sandy ground, under 50 m: *Kerr* 20642 (B).  
PEN. — Surat, Kanchanadit, common in savannah, 5 m: *Kerr* 13067 (BK, K).  
Satul, in savannah, 5 m: *Kerr* 13299 (BK).  
DISTR. From S. E. Asia to tropical Australia.

50. *F. ADENOLEPIS* Kern, Blumea 8: 123, f. 3. 1955; Kern: 645.  
S.E. — Prachinburi, Aran Prathet, in open low-lying, grassy ground, 50 m: *I*  
19566 (BK, K).  
PEN. — Surat, Kun Tale Lake: *Seidenfaden* 2326 (SING).  
DISTR. Also known from Indo-China and Malaysia (Kangean Island).

51. *F. SAVANNICOLA* Kern, Blumea 10: 635. 1960.  
N.E. — Loei, Pu Tong, in savannah, 1100 m, local name *ya nuat meo*: *Kerr* 8862  
type: BK.  
DISTR. Only known from the type collection.

52. *F. FUSCOIDES* Clarke in Ostenf., Bull. Herb. Boiss., sér. 2, 5: 7.  
1905; Hoss. 1910: 462; F.I.C.: 124 (specim. Siam); Turr.: 424; F.M.P.: 1.  
Kern, Blumea 8: 125. 1955; Kern: 647. — *F. angustifolia* Ridl., Journ.  
Br. Roy. As. Soc. no 59: 223. 1911. — *F. erythradenia* Camus, Not. S.  
1: 247. 1910; F.I.C.: 113.  
N.E. — Loei, Kao Krating, on open sandy ground, 1200 m: *Kerr* 5672 (BK);  
Krading, common on rocks in savannah, 1300 m: *Smitinand* 1937 (L).  
C. — Nakawn Nayok, Khao Lou, wet place: *Sovatabantu* s.n. (BK).  
S.E. — Chanburi, Makham, common in the plain, 20 m: *Smitinand* 5741 (RFD 2111)  
(L).  
PEN. — Pattani, on open sandy ground, near sealevel: *Kerr* 7738 (BK, SING).

Satul, sandy places on heaths: *Ridley 14822* (type collection of *Fimbristylis angustifolia* Ridl.) (BM, K, SING).

DISTR. Also in Cochinchina and W. Malaysia (Malay Peninsula, Billiton, Labuan Island).

53. *F. FUSCA* (Nees) Clarke, Fl. Br. Ind. 6: 649. 1893; Will.: 225; Mat. F.M.P.: 97, p.p.; Hoss. 1910: 461; 1911a: 372; 1911b: 370; F.I.C.: 123; F.M.P.: 160, p.p.; Kern: 647. — *Abildgaardia fusca* Nees in Wight, Contr. Bot. Ind.: 95. 1834. — *F. subfusca* Camus, Not. Syst. 1: 248. 1910; F.I.C.: 123. — *F. rigidifolia* Ridl., Journ. Str. Br. Roy. As. Soc. no 59: 223. 1911; F.M.P.: 159.

N. — Lampang, Ngao, Mae Huat, common in deciduous forest, 360 m: *Aminat 37* (RFD 11275) (L).

N.E. — Loei, Phu Kradung, common in open pine forest, local name *yak bai bit*: *Dee 127* (RFD 4808) (L); same locality, in savannahs, 1300 m: *Smitinand 5851* (RFD 21481) (L), 5959 (L).

S.E. — Krabün, Aranya, in deciduous forest, under 50 m: *Kerr 19304* (BK).

Chanthuri, Makhamb, common in thickets, 20 m: *Smitinand 5755* (L).

S.W. — Kanburi, Pa-Chan-dai, in mixed forest: *Bunnak 844* (L).

PEN. — Trang, Sikao, on sandy soil in light evergreen forest, under 50 m: *Kerr 19028* (BK).

Patulung, Sak, in savannah, under 50 m: *Kerr 19259* (BK).

Songkhla, Tepa, in savannah, under 50 m: *Kerr 14710* (BK).

Satul, in savannah: *Kerr 13689* (BK); Satul, heaths: *Ridley 14877* (type collection of *Fimbristylis rigidifolia* Ridl.) (K, SING).

DISTR. From India to S. China, Japan, and Malaysia.

54. *F. FIMBRISTYLOIDES* (F. v. M.) Druce, Rep. Bot. Exch. Cl. Br. Isl. 1916: 623. 1917. — *Abildgaardia fimbristyloides* F. v. M., Fragm. Phyt. Austr. 8: 273. 1874. — *F. lacei* Turr., Kew Bull. (1911): 348. 1911.

E. — Korat, Ban Chumsaeng: *Nai Noe 195* (BK, K).

DISTR. Burma, Riu Kiu Islands, S. Korea, Malaysia, tropical Australia.

55. *F. STRAMINEA* Turr., Kew Bull. (1911): 192. 1911; Turr.: 425; F.I.C.: 124. — *F. nigrobrunnea* Thwaites var. *thorelli* Camus, Not. Syst. 1: 248. 1910, p.p.; F.I.C.: 121, p.p.

N. — Chiengmai, Doi Suthep, 550—600 m: *Kerr 832* (type) (K); same locality, common in dry gravelly soil in dry deciduous forest, 700 m: *Smitinand 3999* (RFD 16098) (BK, L).

DISTR. Tonkin, Cochinchina.

56. *F. DISTICHA* Boeck., Linnaea 38: 393. 1874; F.B.I.: 651; Kern: 645.

— *F. fuscoides* (non Clarke) F.I.C.: 124 (specim. Cochinch.).

C. — Nakhawn Nayok, on the way to Nang Nawng, in rocky ground, 100 m: *Smitinand 6083* (L).

PEN.—Satul, in open sandy ground, 5 m: *Kerr 13734* (BK).

DISTR. Burma, Cochinchina, Tenasserim and Andamans, S. China, Sumatra.

57. *F. THOUARSHII* (Kunth) Merr., En. Philip. Fl. Pl. I: 127. 1923.  
*Arthrostylis thouarsii* Kunth, En. 2: 284. 1837. — *Arthrostylis filiformis* Thwaites, En. Pl. Zeyl.: 352. 1864. — *Actinoschoenus filiformis* (Thwaites) Benth. in Hook., Ic. Pl. 14: 33, t. 1346. 1881; Mat. F.M.P.: 84; F.M.P.: 165. *F. actinoschoenus* Clarke, Fl. Br. Ind. 6: 650. 1893.

N.—Chiengmai, Doi Nang Ku: *Pat 3340* (BK).

N.E.—Loei, Phu Kradung, Tham Saw, on loose sandy soil in dry evergreen forest, 1300 m: *Smitinand 393* (RFD 8866) (BKF); Kao Kradung, in open ground, e. of evergreen forest, 1200 m: *Kerr 8704* (BK).

S.E.—Trat, Kao Kuap, in open grassy ground, 700 m: *Kerr 17788* (BK).

PEN.—Ranawng, Kao Pawta Luang Keo, in open ground on ridge, 1300 m: *Kerr 16948* (BK, L).

Phang-nga, Kao Katakwam, on open ground, 1000 m: *Kerr 18484* (BK).

Songkhla, Kao Kee, in rock crevices on open rocky slopes, 300 m: *Kerr 1515* (BK, L).

Satul, Adang, in open rocky ground, 500 m: *Kerr 14130* (BK).

DISTR. Madagascar, Mauritius, Ceylon, Indo-China, S. China, Malaya.

58. *F. YUNNANENSIS* Clarke, Journ. Linn. Soc., Bot. 36: 247. 1903; Turi 425; Kern: 648.

N.—Mae Hawng Sawn, common in rocky ground in dry *Dipterocarpus* forest, 500 m: *Smitinand 4612* (L).

Chiengmai, Doi Suthep, common in gravelly soil in dry deciduous forest, 600-700 m: *Smitinand 3987* (RFD 16097) (BKF, L); Me Tun, open grassy pine forest, 1100 m: *Kerr 6248* (BK).

DISTR. Yunnan.

59. *F. HOOKERIANA* Boeck., Linnaea 37: 22. 1871; F.B.I.: 641; F. K. 87; Will.: 225; Hoss. 1910: 462; F.I.C.: 111, f. 16, 10—11; Kern, Blum. Suppl. 4: 165. 1958.

N.E.—Loei, Phu Kradung, Rest House, common by paths, 1300 m: *Smitinand 512* (L); in savannah on rocks, 1300 m: *Smitinand 1949* (BKF, L); in open pine forest, local name *yah huu baw*: *Dee 148* (RFD 4903) (BKF, L); Phu Kradung, Pha Mak Duk, common in open pine forest, 1300 m: *Dee 10* (RFD 3552) (BKF).

E.—Korat, Katok, on rocky ground in open forest, 300 m: *Kerr 8147* (BK).

S.E.—Trat, Kao Kuap, on rocks, 300 m: *Kerr 17686* (BK); grew at Bangkok in brought from Kao Kuap: *Kerr 19723* (BK, K).

Kaw Chang, Khlawng Kloi, on open rocky ground, 50 m: *Kerr 8245* (BK).

PEN.—Songkhla, Kao Keo, in rock crevices in open rocky ground, 100 m: *Kerr 1515* (BK, K).

DISTR. Khasia Hills, Chota Nagpore, Indo-China.

60. *F. FERRUGINEA* (L.) Vahl, En. 2: 291. 1806; F.B.I.: 638; F. KC.: 88; Will.: 224; Mat. F.M.P.: 92; Hoss. 1910: 461; F.I.C.: 108; F.M.P.: 576. — *Scirpus ferrugineus* L., Sp. Pl. 1: 50. 1753. — *F. polytrichoides* (non R. Br.) Mat. F.M.P.: 90, p.p.; F.M.P.: 154, p.p. — *F. longispica* (non Steud.) F.I.C.: 107.

VAR. *FERRUGINEA*.

C. — Bangkok, open low-lying ground by canal: *Kerr* 6979 (BK); Bangkok, Patanan, open ground flooded at high tide: *Kerr* 6787 (BK); Bangkok, Wat-Sing, ground inundated at high tide, under 5 m: *Kerr* 6707 (BK); Bangkok, Klong edge, sea-level: *Ergl Smith* 397 (BK).

S.E. — Kaw Chang, Khlawng Mayom, sandy ground close to sea: *Kerr* 6864 (BK, K). S.W. — Prachinap, Kao Tao, on open low-lying ground, under 5 m: *Kerr* 16182 (BK).

Surat, Kaw Prap, among stones on beach: *Kerr* 13080 (BK, K); Kaw Samui, Ban Bua Put, in *Pluchea* scrub, under 5 m: *Kerr* 15705 (BK, K).

Satthal, open ground at edge of mangrove forest: *Kerr* 14243 (BK).

DISTR. Pan-tropic.

VAR. *SIEBOLDII* (Miq. ex Franch. & Savat.) Ohwi, Journ. Jap. Bot. 14: 576. 1938. — *F. sieboldii* [Miq., Cat. Mus. Bot. Lugd. Bat.: 118. 1870, nom. nud.] ex Franch. & Savat., En. Pl. Jap. 2: 118. 1879.

S.E. — Trat, Kaw Chang, Khlawng Nonsi, by paths in sandy soil, 5 m: *Smitinand* 1650 (L).

DISTR. Japan, Korea, Riu Kiu Islands.

61. *F. TRISTACHYA* R. Br., Prodr.: 226. 1810; Kern, Blumea 8: 131. 1955. — *F. annamica* Camus, Not. Syst. 1: 245. 1910; F.I.C.: 95. — *F. allei-zethei* Camus, Not. Syst. 1: 291. 1910; F.I.C.: 112. — *F. podocarpa* (non Nees) Ridl. 1911: 222; F.M.P.: 155.

S.E. — Chanthaburi, Makhram, common in wet localities in the plain: *Bennak* 174 (RFD 5518) (L), *Smitinand* 5756 (L).

Trat, Huay Raeng, Dong Madusae, in rice-fields, under 50 m, local name *yeh hang* ia: *Smitinand* 1334 (RFD 6774) (BKF). Chumphawn, Ban Tung Maha, in open low-lying ground, near sealevel: *Kerr* 11409 (BK, K); Bang Son; *Pot* 1542 (BK).

PEN. — Surat, Kanchanadit, common in savannah, 5 m: *Kerr* 18068 (BK, K).

Patalung, Sak, in savannah, under 50 m: *Kerr* 19257 (BK).

Songkhla, Tepa, in savannah, under 50 m: *Kerr* 14713 (BK).

Satthal, heaths: *Ridley* 14878 (BM, K, SING).

DISTR. From India to Micronesia and Australia.

62. *F. SCHOENOIDES* (Retz.) Vahl, En. 2: 286. 1806; F.B.I.: 634; Mat. F.M.P.: 90; F.I.C.: 99; Turr.: 424; F.M.P.: 154; Koy. 1957: 44. — *Scirpus schoenoides* Retz., Obs. 5: 14. 1789.

S.E. — Makhram, common in the plain, 20 m: *Smitinand* 5771 (L).

Trat, common in lawn: *Smitinand* 5729 (RFD 21506) (L).

PEN. — Surat, edge of pool in open grassy ground, under 10 m: *Kerr 11245* (BK).  
 DISTR. From S. E. Asia to tropical Australia.

63. *F. SUBALATA* Kern, Blumea 8: 133, f. 2. 1955. — *F. alata* Cam. Not. Syst. 1: 244. 1910, p.p.; F.I.C.: 94, p.p.

PEN. — Surat, Kanchanadit, in savannah, 5 m: *Kerr 13142* (BK, K, L).

DISTR. Also known from Cochinchina, Cambodia, and New Guinea.

64. *F. DICHOTOMA* (L.) Vahl, En. 2: 287. 1806. — *Scirpus dichotoma* L., Sp. Pl. 1: 50. 1753. — *Scirpus diphylloides* Retz., Obs. 5: 15. 1789. *F. diphylloides* (Retz.) Vahl, En. 2: 289. 1806; F.B.I.: 636; F. KC.: 85; W. 224; Mat. F.M.P.: 91; Hoss. 1910: 461; 1911b: 370; F.I.C.: 103; Turr. 4 F.M.P.: 155. — *F. annua* R. & S., Syst. Veg. 2: 95. 1817; Kill.: 56. — *F. longistylis* (non Steud.) Mat. F.M.P.: 93, p.p.; F.M.P.: 156, p.p. — *F. germanica* Camus, Not. Syst. 1: 246. 1910; F.I.C.: 105.

N. — Chiengmai, Doi Suthep, common in dry deciduous forest, 750 m: *Smitinand 1933* (L); same locality, in moist ground, 1100 m, local name *yah niu nn*: *Soredia (RFD 4313)* (BKF); Me Awn, along edge of open stream, 450 m: *Kerr 4702* (B).

N.E. — Loei, Phu Kradang, in open pine forest, 1300 m: *Dee 36* (RFD 3628); same locality, in savannah, 1300 m: *Smitinand 1933*, 5950 (L).

Nakawn Panom, Dawn Tan, in open deciduous forest, 100 m: *Kerr 21524* (B).

C. — Nakawn Nayok, on the way to Nang Nawng, in rocky ground and by water, 100 m: *Smitinand 6082* (L).

Bangkok, edge of open canal: *Kerr 6982* (BK).

S.E. — Kaw Chang, Khlawng Nonsi, in open, rather marshy ground, under 1 m: *Kerr 9200* (BK).

S.W. — Near Neekey, near Wangka, in meadow, in half shade, 150 m: *den Hoed (Kwae Noi R. basin exp. 1946)* (BK, L); near Rintin, near Kin Sayok, c. 140 m, NW of Kanburi, mixed forest in marshy places on limestone rock, 100—150 m: *Kuetermans 1238* (*Kwae Noi R. basin exp. 1946*) (L); Ku-Jae, c. 150 km of Kanburi, hilly, dry country, 100—150 m: *Kuetermans 1279, 1320* (*Kiae R. basin exp. 1946*) (L).

Chumphawn, Bang Son, in savannah, 100 m: *Kerr 11341* (BK, K).

PEN. — Surat, Bangbao, common in evergreen forest in sandy soil, 50 m: *Phoebe 867* (RFD 14419) (L); Kaw Samui, on open rocks by stream, 50 m: *Kerr 11342* (BK); Kaw Samui, Ban Bua Put, in open grassy scrub: *Kerr 15726* (BK, K). Phang-nga, Nop Pring, in sandy scrub, 100 m: *Kerr 18517* (BK).

Trang, Chawng, on open rocks by stream, 100 m: *Kerr 15160* (BK); Huay Yai, Khao Kaew, common in old clearing, 100 m: *Smitinand 4127* (L); Kantang, common in swamp and muddy bank, under 50 m: *Smitinand 2953* (RFD 12640); Songkhla, Khao Noi, in gravelly ground on ridge, 40 m: *Williams RFD 12640* (BKF).

Satui, Kuan Po, by road-side, 20 m: *Kerr 13765* (BK).

DISTR. All warmer parts of the whole world.

65. *F. BISUMBELLATA* (Forsk.) Bub., Dodec.: 30. 1850. — *Scirpus bisumbellatus* Forsk., Fl. Aeg.-Arab. 1: 15. 1775. — *F. dichotoma* (non Vahl) F.B.I.: 635; Hess. 1910: 461; 1911b: 370; F.I.C.: 102, p.p.; Turr.: 424.

N. — Mae Hawng Sawn, Pang Mu, Nam Pai, in stream bed, 400 m: *Smitinand* 4695 (L).  
N.E. — Nawngkai, Chaiyaburi, on sand-bank, 200 m: *Kerr* 26697 (BK).

Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr* 21365 (BK).

C. — Saraburi, Menam Sak, on muddy river bank, 30 m: *Kerr* 7054 (BK).

S.W. — Kanburi, Wang Kanai, on gravel bar at edge of river, under 50 m: *Kerr* 12872 (BK).

Near Neeckey, near Wangka, near rivers and pools, sandy soil, 150 m: *den Hoed* 150 (*Kwae Noi R. basin exp. 1946*) (BK, L).

DISTR. From the Mediterranean region through the Old World Tropics to N. S. Wales.

66. *F. MERRILLII* Kern, Blumea 8: 135. 1955. f. 6; Kern: 648.

S.E. — Prachinburi, Aran Prathet, in open low-lying grassy ground, 50 m: *Kerr* 19572 (BK, K).

DISTR. Also known from S. China and Malaysia.

67. *F. TRICHOHYLLA* Ridl., Fl. Mal. Pen. 5: 155. 1925; Kern, Blumea 8: 140. 1955; Kern: 648.

In Thailand:

VARI. *ERECTA* Holtt. ex Kern, Blumea 8: 141. 1955.

PEN. — Langsuan, Tako, in clefts of limestone rocks, 100 m: *Kerr* 11876 (BK).

Surat, Ban Kawp Kap, in crevices of limestone rocks, 100 m: *Kerr* 18177 (BK).

DISTR. The var. *trichophylla* is only known from Pulu Langkawi, var. *erecta* occurs also on limestone rocks in the Malay Peninsula.

68. *F. RIGIDULA* Nees in Wight, Contr. Bot. Ind.: 99. 1834; F.B.I.: 640.

N. — Kampeng Pet, Me Lamung, open grassy ground, 800 m: *Kerr* 6104 (BK).

S.W. — Birmese border, Tripagodas, c. 40 km N of Wangka, scattered in grassy vegetation on sandy clay, 280 m: *Wickham* 332 (*Kwae Noi R. basin exp. 1946*) (BK, L).

DISTR. From India to China and the Philippines.

69. *F. SPICIGERA* Kern, Blumea 8: 144, f. 8. 1955.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr* 21366 (K, type; BK, L).

DISTR. Only known from the type collection.

70. *F. PSAMMOPHILA* Kern, Reinw. 4: 96. 1956. — *F. arenicola* Kern, Blumea 8: 146. 1955, non Wiggins 1950.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr* 21367, type collection (BK, K, L).

DISTR. Only known from the type collection.

71. *F. NELMESII* Kern, Act. Bot. Neerl. 7: 800. 1958. — *Tyloca cylindrostachya* Nelmes, Kew Bull. (1949): 139. 1949, non *F. cylindrostachya* Steud. 1855.

S.W. — Brangkasi, on sand-bank in river, rather rare, 150 m: *den Hoed & Koster 368 (Kwae Noi R. basin exp. 1956, type collection)* (L).

DISTR. The species is also in Laos: Hat Kam, Wiengchan, on gravel by river, 200 m: *Kerr 21294* (L), and in Tonkin: prov. de Sontay, baie de la Rivière-Noire entre Van Yen et Van Bu, sur le sable, commun: E. Coe 4055 (Raymond in litt.).

72. *F. JUCUNDA* (Clarke) Kern, Blumea 10: 637. 1960. — *Cyperus jucundus* Clarke, Kew Bull. add. ser. 8: 3. 1908. — *F. brunnea* Clarke E. G. Camus, Not. Syst. 1: 248. 1910; F.I.C.: 106.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr 21295 (BK)*.

DISTR. Tonkin, Laos.

73. *F. BRUNNEOIDES* Kern, Blumea 8: 146, f. 10. 1955.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr 21296 (K, type; BK, L)*.

DISTR. Only known from the type collection.

74. *F. GRIFFITHII* Boeck., Flora 43: 241. 1860; Kern, Blumea 8: 1955. — *F. aestivalis* (non Vahl) Mat. F.M.P.: 92; F.M.P.: 155. — *F. diatomica* (non Vahl) F.I.C.: 102, p.p.

E. — Korat, Katek, on sandy bank by river, 200 m: *Kerr 8190 (BK, K)*.

S.W. — Chumphawn: *Hauiff & Nur SF 4363 (K, SING)*.

DISTR. Bengal, Burma, Tenasserim and Andamans, Indo-China, Mala-

sia.

75. *F. AESTIVALIS* (Retz.) Vahl, En. 2: 288. 1806; F.B.I.: 637; Hoss. 1946; 1911b: 370; F.I.C.: 106; Turr.: 423; Kill.: 56. — *Scirpus aestivalis* Retz. Obs. 4: 12. 1786.

E. — Korat, Ban Chumsaeng: *Pot 2787 (BK)*.

PEN. — Surat, Ta Rong Chang, on sand-bank in river, under 50 m: *Kerr 12310 (B)*.

DISTR. S. and E. Asia, Malaysia, tropical Australia.

76. *F. SQUARROSA* Vahl, En. 2: 289. 1806; F.B.I.: 635; F.I.C.: 101, f. 3—6.

In Thailand:

var. *ESQUARROSA* Makino, Bot. Mag. Tokyo 17: 47. 1903. — *F. verna* R. Br., Prodr.: 227. 1810.

E. — Srisaket, Kantharom, common along river bank, 200: *Smitinand 602 (R 5137 (BKF))*.

S.E.—Chanthaburi, Makham, near water, 20 m: Smitinand 5746 (L).

DISTR. *F. squarrosa* is pantropic; var. *esquarrosa* is known from tropical Africa, Indo-China, Japan, Malaysia, Polynesia, and Australia.

*Fimbristylis squarrosa* is very near to *F. aestivalis* (Retz.) Vahl. Typical *F. squarrosa* is characterized by the long recurved awns of the glumes, in var. *esquarrosa* the glumes are mucronate and the mucros as short as those of *F. aestivalis*. So the long pendent hairs of the style-base (absent or very short in *F. aestivalis*) are the only reliable character to distinguish *F. squarrosa* var. *esquarrosa* from *F. aestivalis*. In Smitinand 5746 the hairs are shorter than usual.

Koyama (1957: 44) thinks that there is a clear line of demarcation between the areas of *F. squarrosa* and *F. velata*, the former being an Eurasian continental species, the latter occurring in Malaysia and Australia. I cannot agree with him.

77. *F. GRACILENTA* Hance, Journ. Bot., London 6: 89. 1868; F.I.C.: 103; Kern: 647. — *F. thorelli* Camus, Not. Syst. 1: 246. 1910; F.I.C.: 105.

N.E.—Loei, Phu Krating, common on rocks in savannah, 1300 m: Smitinand 1951, 5888 (L).

C.—Nakhawn Nayok, on the way to Nang-Nawng, in rocky ground: Smitinand 6034 (L).

R.W.—Chumphawn, Bang Son, in savannah, 50 m: Kerr 11344 (BK).

DISTR. S. China, Indo-China, Sumatra.

var. *psilopoda* Kern, var. nov.

A type rhachi, pedunculus pedicellisque glabris differt.

N.E.—Nakhawn Panom, Dawn Tan, in open deciduous forest, 100 m: Kerr 21522 (BK; type).

The numerous specimens of *Fimbristylis gracilenta* I have seen were all characterized by the puberulous rays and raylets of the inflorescence, a feature of rare occurrence in *Fimbristylis*. In Kerr 21522, otherwise perfectly agreeing with typical *F. gracilenta*, the inflorescence is quite glabrous. This collection may therefore be the type of a new variety.

78. *F. ARGENTEA* (Rottb.) Vahl, En. 2: 294. 1806; F.B.I.: 640; Mat. F.M.P.: 93; F.I.C.: 110; F.M.P.: 157. — *Scirpus argenteus* Rottb., Progr.: 27. 1772.

PHN.—Surat, Ta Rong Chang, on sand-bank in river: Kerr 12308 (BK).

DISTR. Mauritius?; S. E. Asia; also in the Malay Peninsula.

79. *F. POLYTRICHOIDES* (Retz.) R. Br., Prodr.: 226. 1810; F.B.I.: 632; F.K.C.: 88; Willd.: 225; Mat. F.M.P.: 90, p.p.; Hoss. 1910: 462; F.I.C.: 96; F.M.P.: 154, p.p.; Kill.: 56. — *Scirpus polytrichoides* Retz., Obs. 4: 11. 1786.

C.—Tachin, on rather sandy ground by sea: *Kerr 8976* (BK, K); in swampy ground with *Suaeda* sp., near sealevel: *Kerr 16100* (BK).  
DISTR. Tropical Africa, Asia, and Australia.

80. *F. ACUMINATA* Vahl, En. 2: 285. 1806; F.B.I.: 631. Mat. F.M.P.: 89; Ridl. 1911: 222; F.I.C.: 95; F.M.P.: 153. — *F. setacea* (non Benth.) F.B.I. 632, p.p.; Mat. F.M.P.: 89; F.M.P.: 153. — *F. nutans* Vahl var. *minor* Camu Fl. Gén. I.C. 7: 96. 1912.

S.E.—Near Sriracha, rice-fields, 20—30 ft: *Mrs Collins 1889, 1908* (BK).

Trat, Huay Raeng, Dong Maduae, common in rice-field, under 50 m, local name *phrong klom noi*: *Smitinand 1403* (RFD 6775) (BKF); Khlawng Yai, common along edge of thicket, under 10 m: *Smitinand 246* (RFD 6252) (L).

S.W.—Chumphawn, Ban Thung Maha, in open, low-lying ground, near sealevel: *Kerr 11502* (BK); Bang Son: *Put 1548* (BK).

PEN.—Langsuan, Na Sak, in open grassy ground, under 50 m: *Kerr 11863* (BK); Phuket, Khaw Pok-Kasum: *Haniff & Nur SF 3881* (SING).

DISTR. S. E. Asia, extending to the Lesser Sunda Islands, Borneo, and the Philippines.

Clarke and Ridley wrongly referred dwarfish specimens of *F. acuminata* to *F. setacea*, a species restricted to Australia and Eastern Malaysia.

81. *F. NUTANS* (Retz.) Vahl, En. 2: 285. 1806; F.B.I.: 632; Mat. F.M.P.: 89; F.I.C.: 96; Turr.: 424; F.M.P.: 154. — *Scirpus nutans* Retz., Obs. 4: 12. 1786.

E.—Surin, Sangka, in open marshy ground, 300 m: *Kerr 8308* (BK).

PEN.—Ranawng, Kaw Chang, on sandy ground in *Melaleuca* forest, 5 m: *Kerr 16100* (BK).

Pattani, on sandy ground, under 25 m: *Kerr 7237* (BK).

Satul, in open marshy ground, forming an extensive growth, 5 m: *Kerr 13739* (BK).

DISTR. India, Indo-China, S. China, Formosa, Riu Kiu Islands, Malaysia, tropical Australia.

82. *F. TETRAGONA* R. Br., Prodr.: 226. 1810; F.B.I.: 631; Mat. F.M.P.: 88; F.I.C.: 93; Turr.: 425; F.M.P.: 153.

E.—Surin, Sangka, 100 m: *Kerr 8341* (BK).

DISTR. India, Indo-China, S. China, Formosa, Malaysia, tropical Australia.

83. *F. DIPSACEA* (Rottb.) Clarke, Fl. Br. Ind. 6: 635. 1893. — *Scirpus dipsaceus* Rottb., Descr. & Ic.: 56, t. 12, f. 1. 1773. — *Echinolytrum dipsaceum* (Rottb.) Desv., Journ. Bot. 1: 21, t. 1. 1808; F.I.C.: 130.

N.—Chiengmai, Me Ping Rapids: *Umpai 73* (BK); Chainat: *Put 2668* (BK).

N.E.—Nakawn Panom, Tat Panom, on sand-bank in river, 200 m: *Kerr 21358* (BK).

C.—Dhonburi, Bang Plat, common in alluvial deposit around cultivated areas, under 5 m: Smitinand 82 (RFD 5273) (L).

PEN.—Surat, Ta Rong Chang, on sand-bank in river: Kerr 12309 (BK).

DISTR. Tropical Africa, S. and E. Asia, Malaysia.

### X. BULBOSTYLIIS Kunth

84. B. DENSA (Wall.) Hand.-Mazz. in Karsten & Schenk, Vegetationsb. 20, 7: 16. 1930. — *Scirpus densus* Wall. in Roxb., Fl. Ind. 1: 231. 1820. — *Isolepis trifida* Nees in Wight, Contr. Bot. Ind.: 108. 1834. — *B. capillaris* var. *trifida* (Nees) Clarke, Fl. Br. Ind. 6: 652. 1893. — *B. capillaris* (non Clarke) F.I.C.: 127.

N.E.—Loei, Phu Krading, Rest House, in wet localities, 1300 m: Smitinand 3044 (L).

DISTR. Tropical Africa, S. and E. Asia to China and Japan, Malaysia, tropical Australia.

85. B. BARBATA (Rottb.) Clarke, Fl. Br. Ind. 6: 651. 1893; Mat. F.M.P.: 77; Hoss. 1910: 462; 1911b: 369; Ridl. 1911: 222; F.I.C.: 126; Turr.: 425; Koy. 1957: 7. — *Scirpus barbatus* Rottb., Progr.: 27. 1772; Ridl. 1915: 168. — *Stenophyllus barbatus* (Rottb.) Cooke, Fl. Bombay 2: 887. 1908; Kill.: 56. N.—Mae Hawngsawn, lawn, 400 m: Smitinand 4576 (L).

Tak, 30 km N of Tak, common in dry deciduous forest, 100 m: Smitinand 5897 (L). S.W.—Kanburi, in open deciduous forest: Kerr 20492 (BK); Kanburi, Ta Taknaw, local name *ya nuat sia*: Put 10 (BK).

Rathuri, in deciduous forest, 50 m, local name *ya nuat meo*: Kerr 9040 (BK).

Prachuap, Hua Hin, in sand by sea beach: Kerr 13495 (BK); Hui Yang: Put 2179 (BK); Sam Roi Yawt, open sandy ground near sea, local name *yd nusat meo*: Kerr 10881 (BK); Kan Kradai: Put 2304 (BK).

PEN.—Surat, Kaw Tao, in rock cleft near sea: Kerr 13014 (BK).

Satthal, open sandy ground, 5 m: Kerr 13742 (BK).

Pattani, open sandy ground, near sealevel: Kerr 7248 (BK); Yala, open sand-bank in river: Kerr 7726 (BK).

DISTR. Warm parts of the Old World; also in the southern U.S.A.

86. B. PUBERULA (Poir.) Clarke, Fl. Br. Ind. 6: 652. 1893; Mat. F.M.P.: 78; F.I.C.: 128, f. 17, 1—2; F.M.P.: 160. — *Scirpus puberulus* Poir. in Lamk., Enc. 6: 767. 1804. — *Scirpus setaceus* (non L.) F.I.C.: 132. — *Stenophyllus puberulus* (Poir.) Killip, Journ. Siam Soc. Nat. Hist. Suppl. 7: 57. 1927.

PKN.—Pattani, Ban Sai Kao, open sandy ground, 50 m: Kerr 15024 (BK).

DISTR. Tropical Africa and Asia.

### XI. CYPERUS L.

87. C. DIGITATUS Roxb., Fl. Ind. 1: 209. 1820; F.B.I. 618; F.I.C.: 74, t. 3, 5—6. — *Mariscus sieberianus* var. *evolutior* (non Clarke) Mat. F.M.P.:

73, p.p.; Ridl. 1911: 221; F.M.P.: 149, p.p. — *C. elatus* (non L.) F.I.C.: 7 (incl. var. *laxus* Camus).

E. — Korat, Ban Chumsaeng: *Nai Noe* 203 (BK).

C. — Bangkok, edge of open ditch in fruit garden: *Kerr* 8975; Bangkok, Bangkok RFD Compound, under 10 m: *Smitinand* 5412 (RFD 19742) (L).

S.W. — Kanburi, Hin Dat: *Put* 105 (BK).

Brangkasi, along river-bank, 150 m: *den Hoed & Kostermans* 695 (Kwae Noi basin exp. 1946) (L).

PEN. — Surat, Tassateng, Bandon River, along the river, 5 m: *Seidenfaden* 2111 (SING).

Nakawn Srithamarat, Kao Chum Tawng, in shallow ponds, under 50 m: *Kerr* 14339 (BK).

DISTR. Pantropic.

88. *C. ELATUS* L., Cent. Pl. 2: 301. 1756; F.B.I.: 618; Mat. F.M.P.: 7 F.M.P.: 148; Kill.: 55. — *C. digitatus* (non Roxb.) Mat. F.M.P.: 71; F.M.P. 147. — *C. radiatus* (non Vahl) F.I.C.: 72, f. 9, 1—4.

C. — Bangkok, Klong, 5 m: *Ergil Smith* 109 (BK).

DISTR. S. and S. E. Asia, Malaysia.

89. *C. IMBRICATUS* Retz., Obs. 5: 12. 1789. — *C. radiatus* Vahl, En.: 369. 1806; F.B.I.: 617; Mat. F.M.P.: 70; F.M.P.: 147; Turr.: 422.

N. — Mae Hawngsawn, Pang Mu, Nam Poi, in streambed, 400 m: *Smitinand* 4609 (L).

N.E. — Nakawn Panom, on sand-bank in river, 200 m: *Kerr* 21387 (BK), 21391 (BK).

C. — Angthawng, Ayuthia: *Suvatabandhu* s.n. (BK).

S.W. — Kanburi, Brangkasi, river-banks, 150 m: *den Hoed & Kostermans* 697 (Kwae Noi R. basin exp. 1946) (L).

DISTR. Pantropic.

90. *C. EXALTATUS* Retz., Obs. 5: 11. 1789; F.B.I.: 617. — *C. elatus* var. *macronyx* (non Clarke) F.I.C.: 72.

S.W. — Petchaburi, the common species in stagnant water of open ditches, 10 m: *Kerr* 11085 (BK).

DISTR. Tropical Africa, S. and E. Asia, Australia.

91. *C. STOLONIFERUS* Retz., Obs. 4: 10. 1786; F.B.I.: 615; Mat. F.M.P. 69; Ridl. 1911: 221; F.M.P.: 145. — *C. bulbosus* (non Vahl) F.I.C.: 64, ind. var. *elatus* Camus.

S.E. — Chantabun, Lem Sing, in rock crevices close to sea, near sealevel: *Kerr* s.n. (BK).

Trat, Kaw Chang, Khlawng Nonsi, in sand by sea, 5 m: *Smitinand* 5651 (L).

S.W. — Prachuap, Hua Hin, in sand by sea: *Kerr* 12418 (BK).

Kaw Tao, in sandy ground by sea: *Kerr* 16187 (BK), 11157 (BK).

PEN. — Kaw Samui, Ban Bua Put, in sandy ground near sea: *Kerr* 13760 (BK).

DISTR. From S. E. Asia to Melanesia and tropical Australia.

92. *C. ROTUNDUS* L., Sp. Pl. 1: 45. 1753; F.B.I.: 614; Mat. F.M.P.: 68; Hass. 1910: 459; 1911a: 372; 1911b: 370; Ridl. 1911: 221; F.I.C.: 65; Turr.: 422; F.M.P.: 145; Kill.: 55.

N.—Petchabun, Lom Kao, Phu Hi, common in damp localities, in village, 860 m: Smitinand 2590 (L).

Nakhon Sawan, Wang Diao, savannah forest: *Hossenk 133a* (BM, L).

C.—Bangkok, rice-fields: *Feilberg s.n.* (C); Krungthep, Bangkhen, common in damp localities, under 10 m: Smitinand 4032 (RFD 18095) (L).

S.W.—Kanchanaburi, Wang Pra: *Pnt 24* (BK).

Brangkasi, c. 100 km S of Wangka, along paths in half shade, 150 m: *den Hoed & Kostermans 985* (*Kwae Noi R. basin exp. 1946*) (BK, L).

FEN.—Pattani, Gwynne-Vaughan 586 (L).

DISTR. Warmer parts of the whole world.

ssp. *TUBEROSUS* (Rottb.) Kük., Pflanzenr., Heft 101: 113. 1935.

C.—Bangkhen, along ditches, local name *ya takrap*: *Kerr 10713* (BK); Bangkhen, RFD Compound, common in wet ground: *Smitinand 4556* (RFD 17839) (L).

S.W.—Petchaburi, edge of open ditch, 10 m: *Kerr 11684* (BK).

DISTR. of the subspecies. Widely spread in Africa and S. Asia.

93. *C. TENUICULMIS* Boeck., Linnaea 36: 286. 1870; Kern, Reinw. 3: 30. 1954. — *C. zollingeri* (non Steud.) F.B.I.: 613; Mat. F.M.P.: 68; F.I.C.: 67, f. 7, 3—5; Turr.: 422; F.M.P.: 144.

N.—Chiangmai, Doi Suthep, common in oak forest, 1100 m, local name *yah dawk daeng*: *Soradet 113* (RFD 3857) (BKF).

Petchabunn, Lomkao, Na Phaw Sawng, common in pine and oak forest, 900 m: *Smitinand 2632* (RFD 15186) (L); Lomkao, Phu Hee, scattered in oak forest, 800 m: *Smitinand 2593* (RFD 15137).

S.E.—Loei, Phu Kradung, Rest House, in open pine forest, 1300 m: *Smitinand 6073* (L); Kao Kradung, *Suvatavabandhu 211* (PNH).

Nakorn Panom, Muk Dahan, in deciduous forest, 200 m: *Kerr 21414* (BK).

S.E.—Chanthaburi, Khlung, Map Phae, common in wet localities in the plain, 180 m: *Bunak 218* (RFD 9523) (L).

FEN.—Surat, in scrub, under 10 m: *Kerr 11262* (BK); Surat, Kantuli: *Pnt 4131* (BK).

DISTR. Tropical Africa, S. E. Asia to Micronesia and Australia.

94. *C. DISTANS* L.f., Suppl.: 103. 1781; F.B.I.: 607; Mat. F.M.P.: 69; F.I.C.: 60; Turr.: 422; F.M.P.: 145; Kill.: 55.

N.—Chiangmai, Huay Kaew, common in moist localities, 350 m: *Smitinand 4758* (RFD 17849) (L).

C.—Bangkok, open waste ground recently cleared: *Kerr 6990* (BK).

Suraburi, Menam Sak, on sand-bank with coarse grass (*Saccharum* sp.), 20 m: *Kerr 7012* (BK, SING).

S.E.—Trat, Kaw Kot, in cocoanut plantation, 5 m: *Smitinand 3709* (RFD 21567) (L).

S.W.—Kanchanaburi, Ta Salao, in scrub on river bank, 50 m: *Kerr 19532* (BK); Brangkasi, scattered, open spaces, 150 m: *den Hoed & Kostermans 696* (*Kwae Noi R. basin exp. 1946*) (L).

Near Neeckey, near Wangka, sand and gravel banks along Ran Ti River, 150 m: *den Hoed* 122 (*Kwae Noi R. basin exp. 1946*) (BK, BO, L, PNH, SING); *ibid.* *Bloembergen* 604 (L).

Prachnap, Pak Tawan, on sand-bank by river, 20 m: *Kerr* 20527 (BK).  
DISTR. Warmer parts of the whole world.

95. C. NUTANS Vahl, En. 2: 363. 1806; F.B.I.: 607.

- C.— Saraburi, Muak Lek, sand-bank in stream, 200 m: *Marcan* 1855 (BK).  
Bangkok, in ditch in fruit garden: *Kerr* s.n. (BK).  
Chaibadan, Dong Pya Yen, in open marshy savannah, 100 m: *Kerr* 8009 (BK).  
S.E.— Chanburi, Pong Nam Rawn, Wang Kaphae, common by stream, 160 m: *Smitinand* 3348 (BKF).  
S.W.— Near Neeckey, near Wangka, Ran Ti River, sandy river banks, 150 m: *den Hoed* 211 (*Kwae Noi R. basin exp. 1946*) (L); near Kin Sayok, c. 120 km W. of Kanburi, mostly dry limestone rocks in rivulet, 100 m: *Kostermans* 11000 (*Kwae Noi R. basin exp. 1946*) (L).  
Bangtaphan: *Keith* 406 (SING).  
DISTR. S. and S. E. Asia, Malaysia, ? W. Africa.

96. C. PROCERUS Rottb., Descr. & Ic.: 29. 1773; F.B.I.: 610; Mat. F.M.P.

- 70; F.I.C.: 62; F.M.P.: 140. — *Pycreus puncticulatus* (non Nees) Mat. F.M.P. 61; Ridl. 1911: 220; F.M.P.: 140. — *Pycreus baccha* (non Nees) F.I.C.: 31.  
C.— Bangkok, on bank of open canal: *Kerr* 6984 (BK); in flooded rice-fields: *Kerr* 11101 (BK); local name *kok takluk*: *Susatavandhu* s.n. (BK).  
DISTR. Madagascar; from India, Cochinchina, E. China, and Formosa southward to Queensland.

97. C. PILOSUS Vahl, En. 2: 354. 1806; F.B.I.: 609; Mat. F.M.P.: 61.

- F.I.C.: 62, f. 6, 12; Turr.: 422; F.M.P.: 144; Koy. 1955: 207; 1957: 25.  
S.W.— Near Neeckey, near Wangka, on sandy soil along Ran Ti River, 150 m: *den Hoed* 139 (*Kwae Noi R. basin exp. 1946*) (BK, L).  
PEN.— Trang, Chawng, common by stream, 100 m: *Smitinand* 4085 (L).  
DISTR. From Central Asia and Japan through tropical Asia to N. Australia; tropical W. Africa.

98. C. BABAKAN Steud., Syn. 2: 6. 1855; Kern, Reinw. 2: 105, f. 1. 1952. — *C. babakensis* Steud. ex Miq., Fl. Ind. Bat. 3: 257. 1856; F.B.I.

- 610; F.I.C.: 61; Turr.: 421; F.M.P.: 144.  
S.E.— Kaw Chang, Lem Dan Kao, marshy ground by sea, 5 m: *Kerr* 9287 (BK).  
PEN.— Surat, Kanchanadit, pond in savannah, 5 m: *Kerr* 13099 (BK).  
Nakawn Srithamarat, Padang Besar, in light evergreen forest, 50 m: *Kerr* 13098 (BK, BO).  
Songkhla, Kampangpet, in edge of marshy ground, under 50 m: *Kerr* 15982 (BK).  
Pattani, Bansang Sta, in rather marshy paddy fields, 50 m: *Kerr* 7294 (BK).  
DISTR. From Hindustan and Assam to Cochinchina and Malaysia.

99. *C. CORYMBOSUS* Rottb., Descr. & Ic.: 42, t. 7, f. 4. 1773; F.B.I.: 612; F.I.C.: 66 — *C. longus* (non L.) F.I.C.: 68.

S.—Lampang, Mae Paen, in marshy place, 160 m: *Winit* 1722 (BK, BKF).

C.—Pradumdhani, common in ditches by roadsides, under 50 m: *Smitinand* 4530 (RFD 18590).

Bangkok, along open tidal canals: *Kerr* 6985 (BK); Bangkok, local name *kok kluu*, *kok chantaburee*: *Suvatibandhu* s.n. (BK).

SE.—Chanthaburi, cultivated, used for mat-making, under 50 m, local name *kok*: *Kerr* 18051 (BK).

DISTR. Tropics and subtropics of the whole world.

Sometimes cultivated for matting purposes.

100. *C. MALACCENSIS* Lamk, Ill. 1: 146. 1791; F.B.I.: 608; Mat. F.M.P.: 70; Ridl. 1911: 221; F.I.C.: 63; Ridl. 1920: 124; F.M.P.: 147; Kill.: 55.

C.—Bangkok, along edge of open canal: *Kerr* 6980 (BK, SING); along ditch: *Kerr* 10716 (BK).

SE.—Chanthaburi: *Vesterdal* 459 (C).

SEN.—Surat, Bandon: *Seidenfaden* 2039 (SING).

Satui, *Ridley* 14829 (SING).

DISTR. Widely spread in the Eastern Hemisphere.

101. *C. TONKINENSIS* Clarke, Kew Bull. add. ser. 8: 11. 1908; Camus, Not. Syst. 1: 241. 1910; F.I.C.: 70.

NE.—Nakawn Panom, Tat Panom, on sand-bank in river, 200 m: *Kerr* 21397 (BK).

SW.—Kanburi, Wang Kanai, on gravel bar at edge of river, under 50 m: *Kerr* 12371 (BK).

DISTR. Also known from Tonkin (type locality).

*Cyperus brevicaulis* Clarke, Kew Bull. add. ser. 8: 11. 1908, was based on two *Balanusa* collections (2832 and 2833). I am pretty certain that the duplicates of *Balanusa* 2831, type collection of *C. tonkinensis*, and that of *Balanusa* 2832, both in the Leyden Herbarium, represent but one species.

102. *C. IRIA* L., Sp. Pl. 1: 45. 1753; F.B.I.: 606; Mat. F.M.P.: 66; Ridl. 1911: 221; F.I.C.: 59, f. 6, 4—6; Turr.: 422; F.M.P.: 143; Kill.: 55.

S.—Nakhon Sawan, Chainat, Manorom, rice-field weed, under 50 m: *Kerr* 19673 (BK); Takli: *Put* 2145 (BK); Muang Sing: *Put* 2630 (BK).

NE.—Loei, Phu Kradeng, swamp, 1300 m, local name *hangka khaow*: *Dee* 158 (RFD 2679) (L); ibid.: *Smitinand* 1944 (L); *Smitinand* 5558 (RFD 21492) (L).

Nakhon Panom, on sand-bank in river, 200 m: *Kerr* 21379 (BK), 21385 (BK).

C.—Bangkok, road side: *Kerr* 11633 (BK), 19727 (BK); Bangkok, open marshy ground: *Kerr* s.n. (BK); Bangkok, Klong side: *Eryl Smith* 1037 (BK).

Angthaweng, Ayuthia: *Put* 2599 (BK).

SE.—Kao Chang, Lern Dan Kao, open marshy ground by sea, 5 m: *Kerr* 9299 (BK).

S.W.—Ran Ti River, near Neeckey, near Wangka, along river banks, 150 m: *den Hoed* 206 (*Kwae Noi R. basin exp. 1946*) (BK, BO, G, L, PNH, SING).

DISTR. Tropical Asia to China and Japan, Malaysia, tropical E. Africa, Australia.

103. *C. COMPRESSUS* L., Sp. Pl. 1: 46, 1753; F.B.I.: 605; Mat. F.M.P.: 67; F.I.C.: 57, f. 5, 3—5; F.M.P.: 144.

C.—Paknam, in crevices of cement path, near sealevel: *Kerr* 12946 (BK).

S.W.—Prachuap, Hua Hin, in open grassy, sandy ground, near sealevel: *Kerr* 1312 (BK).

DISTR. Pantropic.

104. *C. FLABELLIFORMIS* Rottb., Progr.: 22, 1772.

C.—Bangkok, in stagnant water, local name *ya kok*: *Marcou* 653 (SING), *Marcou* 1924 (SING); in a fruit garden ditch, probably planted, under 5 m: *Kerr* 1907 (BK).

DISTR. Tropical Africa, elsewhere sometimes escaping from cultivation.

105. *C. DIFFUSUS* Vahl, En. 2: 321, 1806; F.B.I.: 603; Mat. F.M.P.: 45; *Hoss.* 1910: 458; 1911a: 374; *Ridl.* 1911: 221; F.I.C.: 54; *Turr.*: 421; *Ridl.* 1920: 124; F.M.P.: 142; *Koy.* 1955: 209; 1957: 27 (incl. var. *microstachys*).

N.—Lampang, Huay Luang, along edge of stream in evergreen forest, 140 m: *Wittmer* 1887 (BK, BKF).

Nakawn Sawan, Takli: *Put* 2108 (BK).

E.—Korut, Lat Buu Kao: *Put* 4315 (BK).

S.E.—Prachinburi, near Sriracha, 30 ft: *Mrs Collins* 1891 (BK); Nawng Plu: *J. Collins* 846 (BK).

Trat, Koh Chang: *Einar Nielsen* 724 (C).

S.W.—Kanburi, in deciduous forest, under 50 m: *Kerr* 20557 (BK); Ta Salai, bamboo (*Bambusa arundinacea*) forest, 50 m: *Kerr* 19535 (BK).

Near Kin Sayok, c. 120 km NW of Kanburi, bamboo thicket, mostly dry limestone rocks, 100 m: *Kostermans* 1098, 1100 (*Kwae Noi R. basin exp. 1946*) (L); near Neeckey, near Wangka, open dry place, 150 m, local name *neh gay song* (Kam language): *den Hoed* 131 (*Kwae Noi R. basin exp. 1946*) (BK, BO, L, PNH, SING); *ibidem* 233 (BK, L); near Ka Tha Lai, in Pan Paung River valley, very common, bamboo grove, 300 m: *den Hoed* 923 (*Kwae Noi R. basin exp. 1946*) (L); Brangkasi, along paths, 150 m: *den Hoed & Kostermans* 457 (*Kwae Noi R. basin exp. 1946*) (L).

Prachuap, Kao Luang, common in evergreen forest, 300 m: *Kerr* 10873 (BK). Pak Tawan, in bamboo forest: *Kerr* 20521 (BK).

PEN.—Surat, Kaw Tao, glade in evergreen forest, 30 m: *Kerr* 11176 (BK); *12783* (BK, L).

Songkhla, Kaw Maew, shady places, 30 m: *Williams* 17281 (RFD 8833) (BK). Satul, Adang, in evergreen glade, under 5 m: *Kerr* 14085 (BK).

DISTR. From India to S. China and Malaysia.

VIT. MACROSTACHYUS Boeck., Linnaea 35: 534. 1868. — *C. pubisquama* Steud., Syn. 2: 20. 1855; F.B.I.: 604; F.I.C.: 56, f. 57. — *C. diffusus* var. *pubisquama* (Steud.) Hook. f. in Trimen, Handb. Fl. Ceyl. 5: 28. 1900; Mat. F.M.P.: 65; Ridl. 1920: 124; F.M.P.: 143.

NE — Nongkai, Pon Pisai, in evergreen forest, 200 m: Kerr 8554 (BK).

SE — Bayawng, Ban Phae: Put 2779 (BK).

DISTR. As the species.

106. *C. TRIALATUS* (Boeck.) Kern, Reinv. 3: 32. 1954, f. 1, 2. — *Scirpus trialatus* Boeck., Flora 42: 445. 1859. — *C. bancanus* Miq., Fl. Ind. Bat., Suppl.: 599. 1861; Mat. F.M.P.: 65; Ridl. 1911: 221; F.I.C.: 56, f. 5, 10; F.M.P.: 143; Kill.: 55. — *C. turgidulus* Clarke, Journ. Linn. Soc. 21: 130. 1884; F.B.I.: 604. — *C. helferi* (non Boeck.) F.I.C.: 55.

NE — Loei, Phu Krating, Dawn Phra Kaew, scattered in evergreen forest, 1300 m: Smitinand 1864 (RFD 6446) (BKF); Loei, Pu Tong, among rocks in open grassy forest, 1000 m: Kerr 8832 (BK).

Nakorn Panum, Dong Bang-i, in evergreen forest, 200 m: Kerr 21490 (BK).

C — Saraburi, Menam Sak, in rocky limestone hill, 100 m: Kerr 7041 (BK); Saraburi, Phu Khao Arboretum, common in dry evergreen forest, 30 m: Smitinand 3884 (L).

EE — Chantaboon: Vesterdal 8M (SING); Chanburi, Khlung, Map Phae, common in wet localities in the plain, 130 m: Bunnak 217 (RFD 9524) (L).

Trat, Kaw Chang, Khlawng Nonsi, in sandy soil, 5 m: Smitinand 3834 (RFD 12470) (L); Kaw Chang, Pak Khlawlk, common in paddy fields and scrub forest, under 10 m: Bunnak 465 (RFD 12503) (L).

SW — Bangtaphan, jungle: Keith 452 (SING); same locality, beach: Keith 494 (SING).

PEN.— Surat, in evergreen scrub, under 10 m: Kerr 11264 (BK).

Songkhla, Kaw Mae, gregarious in dry shady places, 30 m: Williams 17275 (RFD 12021) (BKF).

Satui: Ridley 14833 (BM, SING).

Without exact locality: Mrs. Collins 849 (BK); Kerr 4196 (P).

DISTR. Indo-China, S. China, W. Malaysia.

107. *C. HELFERI* Boeck., Linnaea 38: 360. 1874; F.B.I.: 604.

PEN.— Ranawng, Lam Lieng, on shingle-bank by stream, 10 m: Kerr 16384 (BK).

DISTR. Burma, Pegu, Mergui, Tenasserim or Andamans.

108. *C. RADIANA* Nees & Mey. ex Kunth, En. 2: 95. 1837; F.B.I.: 605; Mat. F.M.P.: 66; F.M.P.: 143; Kill.: 55. — *C. sinensis* Debeaux, Act. Soc. Linn. Bord. 31: 14. 1877, t. 12; F.I.C.: 52, f. 6, 1—3.

EW — Prachuap, Hua Hin, in sand by sea: Kerr 18417 (BK), s.n. (BK).

PEN.— Phuket, Lem Ta, in sand by sea: Kerr 17390 (BK).

Songkhla, open sandy ground near sea: Kerr 15083 (BK).

DISTR. S. E. Asia to S. and E. China, Formosa, and W. Malaysia.

109. *C. DIFFORMIS* L., Cent. Pl. 2: 6. 1756; F.B.I.: 599; F.I.C.: 49, 1-2; F.M.P.: 142.

N.E.—Nawngkai, Chaiyaburi, on sand bank in Me Kong, 200 m: *Kerr* 20700 (B).  
*C.*—Bangkok, rice-fields: *Feilberg* s.n. (C); Bangkok, Bangkhen, RFD Compound near water, under 10 m: *Smitinand* 5413 (RFD 19741) (L); *Smitinand* 5533 (RFD 17833) (L).

Tachin, in swampy ground with *Suaeda* sp., near sealevel: *Kerr* 16108 (B).  
 Ayuthia, on damp mud at edge of river, 10 m: *Kerr* 7007 (BK).

PEN.—Kaw Samui, Ban Bua Put, in low-lying scrub, under 5 m: *Kerr* 15724 (B).

DISTR. Tropical and warm temperate regions of the Old World + Australia; in C. America presumably introduced.

110. *C. PULCHERRIMUS* Willd. ex Kunth, En. 2: 35. 1837; F.B.I.: 48.

Mat. F.M.P.: 64; Hoss. 1910: 459; 1911b: 370; Ridl. 1911: 221; F.I.C.: 422; F.M.P.: 142. — *C. silletensis* (non Nees) F.I.C.: 47.

N.—Phrae: *C. W. Franck* 489 (C).

E.—Korat, Ban Chumsaeng, in paddy-field: *Noi Noe* 188 (BK).

C.—Chalbadan, Dong Pya Yen, damp mud in mixed forest, 50 m: *Kerr* 7925 (BK).  
 Bangkok, edge of open ditch in fruit garden: *Kerr* 6978 (BK); Bangkok, Ban Kok: *Umpai* 112 (BK).

S.E.—Chantaburi: *Vesterdal* 489 (C, SING).

S.W.—Kanburi, Ta Salao, marshy ground in bamboo (*Bambusa arundinacea*) 50 m: *Kerr* 19522 (BK).

Near Ku-Jae, c. 150 km NW of Kanburi, marshy plains, 100—150 m: *Kostermans* 1319 (*Kwae Noi R. basin exp. 1946*) (L).

PEN.—Nakawn Srithamarat, Tung Song, local name *koh ki ma*: *Rabil* 110 (B).  
 DISTR. S. and S. E. Asia, Malaysia.

111. *C. HASPAN* L., Sp. Pl. 1: 45. 1753; F.B.I.: 600; F. KC.: 82; WI: 224; Mat. F.M.P.: 64; Hoss. 1910: 458; 1911b: 370; Ridl. 1911: 221; F.I.C.: 50; Turr.: 422; Ridl. 1920: 124; F.M.P.: 142; Koy. 1955: 210; 1947: 29. — *C. flavidus* var. *africanus* Camus, Fl. Gén. I.-C. 7: 51. 1912.

N.—Chiengmai, Doi Suthep, swamp, 1100 m: *Smitinand* 114 (RFD 8157) (B).  
 Muang Fang, rocky soil, gray mud near hot spring: *Kostermans* 64 (L).

N.E.—Loei, Phu Krading, common by path near Rest House, 1300 m: *Smitinand* 115 (L).

E.—Chaiyaphun, Pu Kio, in open ground, 1000 m: *Kerr* 20211 (BK, L).

C.—Bangkok, along open ditches: *Kerr* 6977 (BK); on road side: *Kerr* 11057 (BK).  
 Bangkok, Bangkhen, RFD Compound, common near water, under 10 m: *Smitinand* 5414 (RFD 19740) (L).

S.E.—Prachinburi, Aran Prathet: *Pat* 1978 (BK); Lemchabang, swamp, 20 ft: *Collins* 1895 (BK, L).

Trat, Kaw Chang, Khlawng Mayom, on rock crevices by stream, 100 m: *Kerr* 11058 (BK); Kaw Kut, sandy soil in plantations, 5 m: *Smitinand* 5730 (L).

S.W.—Ku-Jae, about 150 km NW of Kanburi, 100—150 m: *Kostermans* 1310 (*Kwae Noi R. basin exp. 1946*) (BO).

- PEN.—Ranswng, Nam Chuet, edge of marshy ground, near sealevel: *Kerr 11711* (BK).  
 Langsuan, Kao Noem Sao, open spaces in evergreen forest, 900 m: *Kerr 12061* (BK).  
 Surat, in open scrub, under 10 m: *Kerr 12205* (BK, L); Kaw Samui, in rock clefts by stream, 100 m: *Kerr 22546* (BK).  
 Trung, Ambo Kao Kao, near stream: *Katul 211* (BK); Kanchang, common in swamp, under 50 m: *Smithson 2949* (RFD 12646) (L).  
 Songkhla, in sandy ground by sea: *Kerr 35113* (BK).  
 Satul: *Ridley 14821*, *14824* (BM, SING).

DISTR. Tropical and subtropical regions of the whole world.

112. *C. TENUISPICA* Steud., Syn. 2: 11. 1855; Kern, Reinw. 2: 116. 1952; 3: 38. 1954. — *C. flavidus* (non Retz) F.B.L: 600; F.I.C.: 51, f. 4, 12, excl. var. *affricanus* Camus.

- N.—Nakawn Sawan, Takd: *Pst 2144* (BK).  
 S.E.—Chanthaburi, Makham, Ban Ang, common in wet localities, 120 m: *Baumak 211* (L).  
 S.W.—Kambari, common in wet ground, 50 m: *Smithson 1855* (RFD 12094) (L).  
 Also in Hainan: Ang Mao, dry paddy fields, 1790 ft: *Eryl Smith 82* (BK).

DISTR. Tropical Africa, Asia (extending to China and Japan), and Australia.

113. *C. CUSPIDATUS* Kunth in H.B.K., Nov. Gen. & Sp. 1: 264. 1815; F.B.L: 598. — *C. squarrosum* (non L.) F.I.C.: 58. — *C. uncinatus* (non Poir.) Mat. F.M.P.: 64; F.I.C.: 47, f. 4, 5—6; Turf.: 422; F.M.P.: 141.

- N.—Mae Hawngsawn, lawn, -400 m: *Smithson 4378* (L); Lampang, Hang Chat, common in open dry *Dipterocarpus* forest, 300 m, local name *kok rangka pak*: *Winit 1750* (BK, BK.F.).

- S.E.—Nakawn Panom, Muk Tahan, on gravel in open deciduous forest, 100 m: *Kerr 8333* (BK).

- E.—Nakawn Rachasima, Korat, on sandy soil in open deciduous forest, 200 m: *Kerr 8085* (BK).

- S.E.—Chanthaboon: Westerval 51 (SING); Lem Sing: Nni Noc 1ff (BK).  
 Trat, Kaw Chang, Pak Khawlk, common in paddy fields and scrub jungle, under 10 m: *Baumak 241* (RFD 12515) (L).

- S.W.—Prachuap, Huai Hin, in open, grassy, sandy ground, near sealevel: *Kerr 12155* (BK).

- PEN.—Surat, Kantuli: *Pst 4-183* (BK); Surat, Banghao, common in sandy soil in evergreen jungle, under 500 m: *Piloschitz 347* (RFD 12826) (L).

DISTR. Pantropic.

114. *C. CASTANEUS* Willd., Sp. Pl. 1: 278. 1797; F.B.L: 598; Mat. F.M.P.: 63; F.I.C.: 46, f. 4, 8—11; F.M.P.: 141. — *C. amabilis* (non Vahl) F.I.C.: 45.  
*C.*—Nakawn Namyok, on the way to Nang Nawng, rocky ground, 100 m: *Smithson 6087* (L).

DISTR. Tropical Asia and Australia.

115. *C. squarrosus* L., Cent. Pl. 2: 6. 1756; Kern: 642. — *C. aristata* Rottb., Progr.: 22. 1772; F.B.I.: 606.

S.W. — Prachuap, Hua Hin, on sandy ground in scrub, 5 m: *Kerr 13467* (BK); S. Roi Yawt: *Put 2501* (BK).

DISTR. Tropical Africa, extending to S. Africa; tropical Asia to S. China and Malaysia; Australia; from S. Canada through the U.S.A. and tropic America to Argentina and Chile.

116. *C. LEUCOCEPHALUS* Retz., Obs. 5: 11. 1789; F.B.I.: 602, p.p. Turr.: 422; Kern, Reinw. 3: 39. 1954; Kern: 644. — *Kyllinga pierreana* Camus, Not. Syst. 1: 290. 1910; F.I.C.: 27.

N. — Mae Hawngsawn, in dry *Dipterocarpus* forest, 400 m: *Smitinand 5568* (L).

N.E. — Loei, Wang Saphung, Sithan, 300 m, local name *yak faek mai*: *Dee 2* (RF 4907).

Nakawn Panom, Muk Dahan, in open deciduous forest, 200 m: *Kerr 21447* (BK). Khawnkaen, Chumpae, common by roadsides in sandy soil, 250 m: *Smitinand 5836* (L).

E. — Srisaket, Kantharom, in scrub jungle near village, 200 m: *Smitinand 588* (RF 5136) (L).

S.E. — Chanburi, Makham, common in the plain, 130 m: *Bunnak 231* (RFD 810) (L); along edge of thicket, 20 m: *Smitinand 5743* (RFD 21509) (L).

Krabi, Aranya, in deciduous forest, under 50 m: *Kerr 19365* (BK).

S.W. — Kanburi, Pa-pu-thong, on level area of mixed forest, 165 m: *Bunnak 451* (L).

DISTR. S.E. Asia.

117. *C. NIVEUS* Retz., Obs. 5: 12. 1791; F.B.I.: 601; Kern: 644. — *C. leucocephalus* (non Retz.) F.I.C.: 51.

N. — Mae Hawngsawn, in dry *Dipterocarpus* forest, 400 m: *Smitinand 4971* (L). Nakawn Sawan, Hua Wai: *Put 4067* (BK).

S.W. — Kanburi, Kao Tawng, in open deciduous forest on limestone hill, 100 m: *Kerr 19715* (BK); Wang Kanai, abundant in open deciduous forest, 50—100 m: *Kerr 12846* (BK).

DISTR. S. Asia.

118. *C. CEPHALOTES* Vehl, En. 2: 311. 1806; F.B.I.: 597; F.I.C.: 46 f. 4, 1—4; Turr.: 421.

E. — Nakawn Rachasima, Korat, open marshy ground, 200 m: *Kerr 8987* (BK).

C. — Bangkok, in open marsh with *Salvinia* sp.: *Kerr 8978* (BK).

PEN. — Surat, forming, with *Salvinia* sp., floating islands in the river, under 10 m: *Kerr 11284* (BK), *Kerr 11288* (BK); Bandon: *Seidenfaden 2008* (SING); Kha Tale Lake: *Seidenfaden 2301* (SING), 2360 (SING).

DISTR. S. and S.E. Asia, Malaysia, tropical Australia.

In *Kerr 11288* and *Seidenfaden 2008*, both from Surat, the leaves are much narrower than usual, but the glumes, nuts, etc. are as in typical *Cyperus cephalotes*.

119. *C. PYGMAEUS* Rottb., Descr. & Ic.: 20, t. 14, f. 4, 5. 1773. — *Juncellus pygmaeus* (Rottb.) Clarke, Fl. Br. Ind. 6: 596. 1893; F.I.C.: 38; Turr.: 421.

N. — Me Ping Rapids, Kaeng Um Luk, on sandy heath; *Sucentabandhu* s.n. (BK).

N.E. — Nakawn Panom, on sand-bank in river, 200 m: *Kerr 21382* (BK).

C. — Bangkok: *Feilberg* s.n. (C); Bangkok, in dried up mud of rice-field: *Kerr 6764* (BK).

S.E. — Chanburi, Makham, near water, 20 m: *Smitinand 5747* (L).

DISTR. N. and E. Africa, S. and E. Asia to Australia.

120. *C. UNILOIDES* R. Br., Prodr.: 216. 1810; Koy. 1955: 213; 1957: 32; Kern: 645. — *C. angulatus* Nees in Wight, Contr. Bot. Ind.: 73. 1834. — *Pycreus angulatus* (Nees) Nees, Linnaea 9: 283. 1834; F.B.I.: 593; F.I.C.: 35. — *Pycreus unioloides* (R. Br.) Urb., Symb. Ant. 2: 164. 1900; Turr.: 421.

N.E. — Loei, Phu Kradung, in savannah, 1300 m: *Smitinand 1952, 5845, 5933* (L).

C. — Chaibadan, Dong Pya Yen, in marshy savannah, 100 m: *Kerr 8007* (BK).

DISTR. Pantropic.

121. *C. SANGUINOLENTUS* Vahl, En. 2: 351. 1806. — *Pycreus sanguinolentus* (Vahl) Nees, Linnaea 9: 283. 1834; F.B.I.: 590; Mat. F.M.P.: 60; F.I.C.: 30; F.M.P.: 139.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr 21385* (BK).

DISTR. Warmer parts of the Eastern Hemisphere.

122. *C. GLOBOSUS* All., Fl. Pedem. Auct.: 49. 1789; Koy. 1955: 213; 1957: 32. — *C. capillaris* Koenig ex Roxb., Fl. Ind. 1: 198. 1820. — *Pycreus globosus* (All.) Reichenb., Fl. Germ. Exc. 2: 140. 1830; F.I.C.: 31, f. 3, 10; Turr.: 421. — *Pycreus capillaris* (Roxb.) Nees, Linnaea 9: 283. 1834; F.B.I.: 591.

N.E. — Nakawn Panom, Paknam Songkram, on sand-bank in river, 200 m: *Kerr 21363* (BK).

S.W. — Kin Sayok, about 140 km NW of Kanburi, on limestone rock in marshy place, 100—150 m: *Kostermans 1239* (*Kwae Noi R. basin exp. 1946*) (BO, L).

DISTR. Warmer parts of the Eastern Hemisphere.

123. *C. POLYSTACHYOS* Rottb., Progr.: 21. 1772; Ridl. 1915: 168. — *Pycreus polystachyos* (Rottb.) Beauv., Pl. Oware & Benin 2: 48, t. 86, f. 2. 1807; F.B.I.: 592; F. KC.: 81; Will.: 223; Mat. F.M.P.: 60; Hoss. 1910: 458; 1911b: 370; Ridl. 1911: 220; F.I.C.: 34; F.M.P.: 139. — *C. odoratus* (non L.) Kill.: 55.

C. — Bangkok, waste ground: *Kerr 3864* (BK); on low-lying open ground: *Kerr s.n.* (BK).

S.E. — Mouth of Chantabun River, on edge of road running through salt marsh, under 5 m: *Kerr 17586* (BK).

Koh Chang, Lem Dan: *Joh. Schmidt 748a* (C).

S.W. — Prachuap, Ban Nawng Kang, open ground, under 50 m: *Kerr 10022* (BK); Kaw Tao, in rock crevices near sea: *Kerr 16023* (BK).

Chumphawn, Bang Son: *Pat 1547* (BK, L).

PEN. — Surat, Ban Dawn, in tidal swamp at mouth of Ban Dawn R.: *Kerr 10023* (BK, L); Kaw Samui, on open rocks by stream, 200 m: *Kerr 12577* (BK); Kaw Samui, Ban Bua Put, in open scrub, under 5 m: *Kerr 15708* (BK).

Trang, Ampo Kao Kao, near stream, local name *kok kima*: *Rabil 240* (BK); Satul: *Ridley s.n.* (SING); Satul, Butang, in rocky ground by sea: *Kerr 10024* (BK).

DISTR. Warmer parts of the whole world.

124. *C. SULCINUX* Clarke, Journ. Linn. Soc. 21: 56. 1884. — *Pyrenula sulcinux* (Clarke) Clarke, Fl. Br. Ind. 6: 593. 1893; F. KC.: 82; Willd. 224; Hoss. 1910: 458.

S.E. — Koh Chang, Lem Dan: *Joh. Schmidt s.n.* (C).

Also in Cambodia: "Mainland opposite Koh Kong": *Joh. Schmidt 300 p.p.* (C).

DISTR. From India to S. China and tropical Australia.

125. *C. PUMILUS* L., Cent. Pl. 2: 6. 1756. — *C. nitens* Retz., Obs. 13. 1789; Hoss. 1911a: 372. — *Pyrenula nitens* (Retz.) Nees, Nov. Act. & Nat. Cur. 19, Suppl. 1: 53. 1843; F.B.I.: 591; Mat. F.M.P.: 61; Hoss. 1911b: 458; 1911b: 370; F.I.C.: 33; Turr.: 421; F.M.P.: 140.

N. — Chiengmai, 1000 ft: *Kerr 862* (L).

Phrae, Mo Ban; *C. W. Franek 1326* (C).

N.E. — Loei, Phu Krading, 1300 m: *Smitinand 1947* (L).

E. — Korat, Pak Tong Chai, open sandy ground, 200 m: *Kerr 8111* (BK).

C. — Nakawn Sawan, Wang Djao, common in savannah-forest: *Hossens 104* (BM).

S.E. — Chantabun, Lem Sing: *Noi Noe 136* (BK); Chantaburi, Makhram, Khlaeng Pong, common by stream, 50 m: *Smitinand 4538* (RFD 19723) (L).

S.W. — Kanburi, common in wet locality, 50 m: *Smitinand 3860* (RFD 16093) (L).

PEN. — Surat, Bandon, Aerodrome Road, humid sandy flats: *Seidenfaden 3428* (L 24297) (C, SING).

DISTR. From India to S. China and tropical Australia.

126. *C. DIAPHANUS* Schrad. ex R. & S., Mant. 2: 477. 1824; Kerr 644. — *C. latespicatus* Boeck., Flora 42: 441 ("433"). 1859. — *Pyrenula latespicatus* (Boeck.) Clarke, Fl. Br. Ind. 6: 590. 1893; F.I.C.: 30.

N.E. — Loei, Phu Krading, Rest House, in savannah, 1300 m: *Smitinand 3953* (L).

S.E. — Prachinburi, Aran Prathet, in open marshy ground, 50 m: *Kerr 19577* (BK, L).

DISTR. From Central Asia to India and Malaysia.

127. *C. SUBSTRAMINEUS* Kük., Pflanzenr. Heft 101: 398. 1936. — *C. stramineus* Nees in Wight, Contr. Bot. Ind.: 74. 1834, non Desf. (1820). — *Papyrus stramineus* (Nees) Clarke, Fl. Br. Ind. 6: 589. 1893; F.I.C.: 29. PEN. — Phaket, Khan Pok Kasum; *Hawiff & Nur SF 3805* (BO, K, SING).

DISTR. From Khasia and Ceylon through India to Farther India and the Malay Peninsula.

128. *C. JAVANICUS* Houtt., Nat. Hist. 2, 13: Aanw. pl. (1), t. 88, f. 1. 1782. — *C. pennatus* Lamk, Ill. 1: 144. 1791; Will.: 224; Hoss. 1910: 458; Kill.: 56. — *Mariscus albescens* Gaudich. in Freyc., Voyage Bot.: 415. 1826; F.B.I.: 623; F. KC.: 82; Mat. F.M.P.: 74; Ridl. 1911: 222; F.I.C.: 78; F.M.P.: 149.

C. — Bangkok, waste ground: *Kerr 3888* (BK); Bangkok, Wat Sing, shady ground inundated at high tide, under 5 m: *Kerr 8706* (BK); Bangkok, Bangkapi, bank of pond: *Boonbrahm 9* (BK).

SE. — Trat, Kaw Chang, Khlawng Nonsi, sandy soil, 5 m: *Smitinand 5055* (L); Koh Lem W of Koh Chang: *Joh. Schmidt 716a* (C); Lem Ngob, inner margin of mangrove: *Joh. Schmidt 40* (C).

PEN. — Surat, Kaw Tao, in tidal marsh: *Kerr 11227* (BK).

DISTR. From tropical Africa and Madagascar through S. Asia to S. China, the Riu Kiu Islands and Formosa, to the Pacific Islands and through Malaysia to tropical Australia.

129. *C. COMPACTUS* Retz., Obs. 5: 10. 1789. — *C. dilutus* Vahl, En. 2: 357. 1806; Will.: 224; Hoss. 1910: 458; Kill.: 56. — *Mariscus microcephalus* Presl, Rel. Haenk. 1: 182. 1828; F.B.I.: 624; F. KC.: 83; Mat. F.M.P.: 74; Ridl. 1911: 221; 1920: 124; Turr.: 423; F.M.P.: 149. — *Sphaeromariscus microcephalus* (Presl) Camus in Lecomte, Not. Syst. 1: 239. 1910; F.I.C.: 79.

C. — Chaibadan, Dong Pya Yen, edge of savannah, 100 m: *Kerr 8012* (BK).

Bangkok, waste ground: *Kerr 3865* (BK); by marshy pool: *Kerr s.n.* (BK). Nonthaburi, local name *kok*: *Varnichgorn 19* (BK).

SE. — Prachinburi, near Sriracha, 30 ft: *Mrs. Collins 1886* (BK); Chanburi, Pong Nam Rawn, Wang Kaphae, 120 m: *Smitinand 3347* (RFD 15677) (L).

S.W. — Kanburi, Wang Ka: *Put 22* (BK).

Near Neekey, near Wangka, rather common in open space among other Cyperaceae and Gramineae, 150 m: *den Hoed 325* (*Kwae Noi R. basin exp. 1946*) (L); marshy plain near Ku Jae, c. 150 km W of Kanburi: *Kostermans 1322* (*Kwae Noi R. basin exp. 1956*) (L).

PEN. — Satul, Khuan Pho, in savannah, 20 m: *Kerr 13761* (BK).

— Dhu Luang (Royal Landing): *Hugh M. Smith 158* (SING).

DISTR. From India to S. China, Formosa, and Malaysia.

130. *C. STENOPHYLLUS* Valck. Sur., Nova Guinea 8: 701, t. 114. 1911.  
 S.T. Blake, Journ. Arn. Arb. 28: 221. 1947.  
 S.E.—Prachinburi, Aran Pratet, marshy ground in open deciduous forest, *Kerr 19618* (BK).  
 DISTR. Solomon Islands, Melanesia, Malaysia.

131. *C. CYPEROIDES* (L.) O. Kuntze, Rev. Gen. Pl. 3, 2: 333. 1898.  
*Scirpus cyperoides* L., Mant, 2: 181. 1771. — *Mariscus sieberianus* Nees ex Clarke, Fl. Br. Ind. 6: 622. 1893; Mat. F.M.P.: 72; F.M.P.: 148. — *Mariscus umbellatus* var. *sieberianus* (Clarke) Camus, Fl. Gen. I.-C. 7: 77. 1912.  
 N.—Chiengrai, Doi Hu Mot, Mae Lao, on ridge, 1450 m: *Smitinand* 772 (BKF).  
 Lampang, Ngao, Mae Huat, common in deciduous forest, 360 m: *Bunnak* 5 (RFD 11260) (L).  
 Mae Hawngsawn, in dry *Dipterocarpus* forest, 400 m: *Smitinand* 4572 (L).  
 Phetchabun, Lomkao, Phu Hee, common in oak forest, 860 m: *Smitinand* 200 (RFD 15139) (L).  
 N.E.—Nakawn Panom, Muk Dahan, in deciduous forest, 200 m: *Kerr* 21418 (BK).  
 S.W.—Kin Sayok, c. 120 km NW of Kanburi, mixed forest along river: *Kostermans* 1212 (*Kwae Noi R. basin exp. 1946*) (L); Birmese border, Tripagodas, 40 km N of Wangka, sandy soil near railroad, 280 m: *Wichian* 337 (*Kwae Noi R. basin exp. 1946*) (BK, BO, L, PNH, SING); near Neeckeey, near Wangka, sandy soil near Ran Ti River, 150 m: *Wichian* 192 (*Kwae Noi R. basin exp. 1946*) (BK, L, PNH, SING); Brangkasi, along paths: *den Hoed & Kostermans* 886 (*Kwae Noi R. basin exp. 1946*) (L).  
 PEN.—Surat, Kaw Tao, in evergreen forest near sea, 10 m: *Kerr* 11180 (BK); same locality, on open sandy ground by sea, near sealevel: *Kerr* 12736 (BK).  
 DISTR. Tropics and subtropics of Africa, Asia, and Australia; in the West Indies presumably introduced.

132. *C. CYPERINUS* (Retz.) Valck. Sur., Gesl. Cyp. Mal. Arch.: 15. t. 6, f. 10. 1898. — *Kyllingia cyperina* Retz., Obs. 6: 21. 1791. — *Mariscus cyperinus* (Retz.) Vahl, En. 2: 377. 1806; F.B.I.: 621; Mat. F.M.P.: 72; Ridl. 1911: 221; Turr.: 423; F.M.P.: 149. — *Mariscus umbellatus* var. *cyperinus* (Retz.) Camus, Fl. Gén. I.-C. 7: 77. 1912.  
 N.—Mae Hawngsawn, in dry *Dipterocarpus* forest, 400 m: *Smitinand* 4572 A (L).  
 Nakawn Sawan, Me Wong, damp ground by stream in mixed forest, 300 m: *Kerr* 6053 (BK).  
 N.E.—Loc, Phu Krading, Sain Khae, by path in evergreen forest, 1100 m: *Smitinand* 5934 (L).  
 Nakawn Panom, Muk Dahan, Pu Mano, in deciduous forest, 200 m: *Kerr* 21418 (BK).  
 S.E.—Krabin, Aranya, in deciduous forest, under 50 m: *Kerr* 19331 (BK).  
 Trat, Kaw Chang, Khlawng Nonsi, in sandy soil, 5 m: *Smitinand* 5653 (L); Ra Kut, in coconut plantation, 5 m: *Smitinand* 5707 (L).

S.W.—Prachup, Kaw Luang, open ground, 1200 m: *Kerr 19843* (BK).

TH.—Surat, Kaw Samui, sandy ground in coconut plantation, near sealevel: *Kerr 12626* (BK).

Nakawn Sritthamarat, Sichon, in scrub, under 50 m: *Kerr 15687* (BK).

Satui: *Ridley 14811* (BM, K, SING).

DISTR. From S.E. Asia through Malaysia to Melanesia, Polynesia, and Queensland.

133. *C. PANICEUS* (Rottb.) Boeck., Linnaea 36: 381. 1870. — *Kylloge paniceus* Rottb., Deser. & Ic.: 15, t. 4, f. 1. 1773.

In Thailand:

var. *ROXBURGHIANUS* (Clarke) Kük., Pflanzenr., Heft 101: 526. 1936.

— *Mariscus paniceus* var. *roxburghianus* Clarke, Fl. Br. Ind. 6: 621. 1893; Turr. 423.

E.—Korat, Ban Chumsaeng, in paddy field: *Nai Nor 197* (BK).

DISTR. of the variety: India, Java, Madura.

134. *C. DUBIUS* Rottb., Deser. & Ic.: 20, t. 4, f. 5. 1773. — *C. kyllogeoides* Vahl, En. 2: 312. 1806; Kill.: 55. — *Mariscus dregeanus* Kunth, En. 2: 120. 1837; F.B.I.: 620; Mat. F.M.P.: 72; Ridl. 1911: 221; Turr.: 423; F.I.C.: 75; F.M.P.: 148. — *Mariscus merrillii* Clarke, Philip. Journ. Sc. Bot.: 87. 1907. — *C. merrillii* (Clarke) Kük., Pflanzenr., Heft 101: 562. 1936.

NE.—Nakawn Panom, Muk Dahan, Pu Mano, in deciduous forest, rocky ground, 200 m: *Kerr 21426* (BK).

E.—Korat, Ban Chumsaeng: *Put 2832* (BK).

C.—Saraburi, Menam Sak, mixed deciduous forest, 40 m: *Kerr 7023* (BK, SING).

S.W.—Kanchanaburi, Ta Salao, in open grassy forest, 50 m: *Kerr 19589* (BK), *Put 3050* (BK).

Prachup, in crevices of limestone rocks by sea: *Kerr 10724* (BK).

Plant brought from Bangtaphan, now (4. 3. 1928) flowering in Bangkok: *Kerr 21883* (BK).

TH.—Surat, Kaw Tao, in rock clefts near sea: *Kerr 11241* (BK); Kaw Samui,

Ban Bua Put, sandy ground near sea: *Kerr 15696* (BK).

Satui, Adang, in crevices of rock near sea, 5 m: *Kerr 14089* (BK); Satui: *Ridley 14812* (K, SING).

DISTR. In the African steppes; S. Asia to S. China and Malaysia.

In Reinwardtia 3: 59. 1954 I referred *Kerr 7023* to *Cyperus merrillii* on account of the narrow glumes and nuts. I then already came to the conclusion that the only reliable characters to distinguish between *C. dubius* and *C. merrillii* seem to be found in the shape of the glumes and the nuts. Since I have seen numerous collections from continental Asia perfectly connecting the two, and it appeared that no doubt they are conspecific. —

135. *C. PEDUNCULATUS* (R. Br.) Kern, Act. Bot. Neerl. 7: 798. 1890.  
*— Remirea maritima* Aubl., Hist. Pl. Guian. Franç. 1: 45, t. 16. 1770; F.B.I.: 677; F. KC.: 92; Will.: 225; Mat. F.M.P.: 99; Hoss. 1910: 46; Ridl. 1911: 224; 1912: 63; F.I.C.: 155; F.M.P.: 169, *non Cyperus remitimus* Poir. — *R. pedunculata* R. Br., Prodr.: 236. 1810.  
 S.E. — Kaw Chang, Khlawng Ktoi, on sandy ground close to sea: *Kerr* s.n. (BK).  
 S.W. — Prachuap, Kao Tao, sandy ground by sea: *Kerr* 18188 (BK).  
 PEN. — Surat, Kaw Samul, in sand on sea-beach: *Kerr* 12586 (BK).  
 Satul, islet off Tenga, in sand of beach: *Kerr* 14756 (BK); Pulau Aleg, *Ridley* 15675 (SING).  
 DISTR. All tropical seacoasts.

136. *C. ODORATUS* L., Sp. Pl. 1: 46. 1753; Dandy in Exell, Cat. Var. Pl. S. Thomé: 360. 1944. — *Torulinium ferax* Ham., Prodr. Pl. Ind. Om. 15. 1825. — *Mariscus fernx* (Ham.) Clarke, Fl. Br. Ind. 6: 624. 1893. — *Mariscus sieberianus* var. *evolutior* (*non* Clarke) Mat. F.M.P.: 73, p.p.; F.M.P.: 149, p.p.  
 N.E. — Nakawn Panom, Muk Dahan, on sand-bank in river, 200 m: *Kerr* 21416 (BK).  
 DISTR. All warm parts of the Old and New World.

137. *C. TRICEPS* (Rottb.) Endl., Cat. Hort. Ac. Vindob. 1: 94. 1824.  
*— Kyllinga triceps* Rottb., Descr. & Ic.: 14. 1773; F.B.I.: 587. — *Kyllinga cylindrica* (*non* Nees) Mat. F.M.P.: 59; F.M.P.: 137.  
 E. — Srisaket, Kanthraram, common in scrub jungle near village, 200 m, local name *sam pi baw haeng*: *Smitinand* 587 (RFD 5135) (L).  
 S.W. — Kanburi, Pa-pu-thong, in mixed forest, 165 m: *Bunnak* 840 (L).  
 DISTR. Tropical Africa, Asia, and Australia.

138. *C. SESQUIFLORUS* (Torr.) Mattf. & Kük., Pflanzenr., Heft 100: 591. 1936.

- In Thailand:  
 var. *CYLINDRICUS* (Nees) Kük., Ic., p. 593. — *Kyllinga cylindrica* Nees in Wight, Contr. Bot. Ind.: 91. 1834; F.B.I.: 588; F.I.C.: 23.  
 N. — Chiangmai, Me Ten, in open evergreen forest, 700 m: *Kerr* 8472 (BK); Do Suthep, common in hill evergreen forest, 1100 m, local name *yab dawk*: *Smitinand* 112 (RFD 3555) (BKF).  
 DISTR. of the variety. Tropical Africa, S.E. Asia to Yunnan and Formosa, Malaysia.

139. *C. BREVIFOLIUS* (Rottb.) Hassk., Cat. Hort. Bog.: 24. 1844. — *Kyllinga brevifolia* Rottb., Descr. & Ic.: 13, t. 4, f. 3. 1773; F.B.I.: 588; Mat. F.M.P.: 58; F.I.C.: 24, f. 3, 3—6; Turr.: 421; F.M.P.: 138; Kill.: 55.

S.—Chiengmai, Doi Suthep, common by stream in damp ground, 1100 m: Smitinand 109 (RFD 6762) (BK); Doi Chiengdao, abandoned clearings, 1400—1500 m: Kerr 6029 (BK).

SE.—Chanthaburi, Thung Maia, common in wet localities, 130 m: Hunnak 197 (L).

NW.—Near Neeckeey, near Wangka, 150 m: den Hoed 145 (Kwae Noi R. basin exp. 1946) (BK, L).

PEK.—Surat, under 10 m: Kerr 11396 (BK); Bangbao, common by river in old clearings, under 50 m: Phloenchit 858 (RFD 12832) (L); Kantuli: Put 4182 (BK). Nakawn Srithamarat, Khiriwong, 100 m: Phloenchit 358 (RFD 11576) (L).

Sangkhla, Saha Yoi, open wet ground, under 50 m: Kerr 14795 (BK).

DISTR. Pantropic.

140. *C. KYLLINGIA* Endl., Cat. Hort. Ac. Vindob. 1: 94. 1842. — *Sydiagn monocephala* Rotth., Descr. & Ic.: 13, t. 4, f. 4. 1773; F.B.I.: 588; F. KC.: 80; Mat. F.M.P.: 58; Hess. 1910: 459; Ridl. 1911: 220; F.I.C.: 25, f. 1, 1—2; Turr. 421; Ridl. 1920: 124; F.M.P.: 138. — *C. monocephalus* (Rotth.) F. v. M., Fragm. 8: 271. 1874; Will.: 224, non Roxb. (1832).

X.—Chiengmai: Kerr s.n. (BK).

C.—Bangkok, in wet place (collector?); Krungthep, Khao Din, common in waste ground, under 10 m, local name *yak tuwhu*: Dee RFD 9134 (L).

SE.—Trat, Kub Chang: Einer Nielsen 696 (C).

NW.—Kanchanaburi, Wang Pra: Put 26 (BK); Brangkasi, common along paths, 150 m: den Hoed & Koetermans 888 (Kwae Noi R. basin exp. 1946) (L); near Neeckeey, near Wangka, 150 m: den Hoed 146 (Kwae Noi R. basin exp. 1946) (BK, BO, L, SING).

Bangtaphan: Keith 541 (SING).

PEK.—Surat, Handon River: Seidenfaden 2225 (C, SING).

Nakawn Srithamarat, Khao Luang, common in evergreen jungle, 600 m, local name *yak wua fa*: Suwan 251 (RFD 13007) (L); Khiriwong, common on sandy bank of stream, 200 m: Phloenchit 853 (RFD 8958) (L).

Tung Song, Ao Wang Kram, in valley, local name *ya uni fa*: Rabil 210 (BK). Without exact locality: Mrs. Collins 850 (BK).

DISTR. Pantropic.

## XII. RHYNCHOSPORE Vahl

141. *R. CORYMBOSA* (L.) Britt., Transact. N.Y. Ac. Sc. II: 84. 1892. — *Scirpus corymbosus* L., Cent. Pl. 2: 7. 1756. — *R. aurea* Vahl, En. 2: 229. 1806; F.B.I.: 670; F. KC.: 91; Will.: 225; Mat. F.M.P.: 83; Hess. 1910: 463; Ridl. 1911: 224; F.M.P.: 164.

C.—Nonthaburi: Varmiekgorn 29 (BK).

Bangkok, on bank of open tidal canal: Kerr 6986 (BK); Bangkok, Klong side: Engl. Smith 339 (BK).

PEK.—Trang, Ampo Kao Kao, near stream, local names *ya kombang* or *ya tamkong*: Rabil 234 (BK); Chawng, common by stream, 100 m: Smitinand 4681 (L).

DISTR. Pantropic.

142. *R. HOOKERI* Boeck., Linnaea 37: 621. 1873; F.B.I.: 671; K. Blumea 9: 230. 1958.

S.E. — Prachinburi, Aran Prathet: Put 1991 (BK).

S.W. — Marshy plain near Ku-Jae, c. 150 km NW of Kanburi, 100—150 m: Kostermans 1369 (L), 1324 (Kwae Noi R. basin exp. 1946) (BO, L, SING).

PEN. — Surat, Kanchanadit, in marshy ground, 5 m: Kerr 13064 (BK, K, L). Without exact locality: Howard 27 (K).

DISTR. India, Burma, Tonkin, Malay Peninsula, W. Java, everywhere scattered.

143. *R. RUBRA* (Lour.) Makino, Bot. Mag. Tokyo 17: 180, t. 7, f. b. 1903. — *Schoenus ruber* Lour., Fl. Cochinch. 1: 41. 1790. — *R. ritchiana* Kunth, En. 2: 289. 1837; F.B.I.: 668; Mat. F.M.P.: 82; Ridl. 1024; F.I.C.: 145; Turr.: 426; F.M.P.: 164.

N. — Chiengmai, Doi Suthep, in gravelly soil in dry deciduous forest, 700 m: Smitinand 3998 (RFD 16096) (L).

N.E. — Loei, Phu Krating, Rest House, in savannah, 1300 m: Smitinand 5764 (RFD 21617) (L).

S.E. — Chanburi, Makham, common in the plain, 20 m: Smitinand 5764 (L). Trat, Huay Raeng, Dong Madue, common in rice-field, under 50 m: Smitinand 1332 (RFD 6771) (L).

S.W. — Ku-Jae, c. 150 km NW of Kanburi, grey loam soil, dry gravel, hilly, 100—150 m: Kostermans 1268 (Kwae Noi R. basin exp. 1946) (L).

Chumphawn, Ban Tung Maha, common in savannah, 20 m: Kerr 11415 (L). PEN. — Surat, Kanchanadit, very common in savannah, 5 m: Kerr 13103 (BK). Patalung, Sak, very common in savannah, under 50 m: Kerr 19258 (BK).

Pattani, Banang Sta, on dry mud in open ground, 50 m: Kerr 7309 (BK). Satul, in grassy savannah, 5 m: Kerr 13688 (BK).

DISTR. From S.E. Asia through Malaysia to tropical Australia and Oceania.

144. *R. LONGISETIS* R. Br., Prodr.: 230. 1810; F.B.I.: 669; Ost. Trop. Hoss. 1910: 463; 1911a: 372; 1911b: 371; Turr.: 426. — *R. museiformis* Camus, Not. Syst. 1: 249. 1910; F.I.C.: 147.

N. — Chiengmai: Kerr 1595 B (L).

Nakawn Sawan, Wang Djao, wet localities in savannah forest, common: Howard 132 (E, L).

S.W. — Petchaburi, Tung Luang, in open sandy ground, under 50 m: Kerr 2664 (L). Prachuap, Hua Hin, in open grassy sandy ground, 5 m: Kerr 13498 (BK).

DISTR. Central India, Burma, Cambodia, New Guinea, tropical Australia.

145. *R. SUBMARGINATA* Kük., Bot. Jahrb. 74: 498. 1949; Kern, Blumea 9: 231. 1958. — *R. marginata* Clarke, Kew Bull. add. ser. 8: 89. 1900 non Steud. (1855). — *R. wightiana* (non Steud.) F.I.C.: 146; F.M.P.: 16.

S.E.—Chanthuri, Maekham, Noen Tabok, common in the field, 120 m: *Bonnak* 195 (RFD 9512) (L); Chantahoon: *Vesterdal* s.n. (SING).

S.W.—Rathavi, open grassy ground, under 50 m: *Kerr* 8988 (BK).

PEN.—Surat, on rice-field bunds, under 10 m: *Kerr* 11248 (BK); Kanchanadit, open sandy ground, 5 m: *Kerr* 13086 (BK); Bandon, Aerodrome Road: *Seidenfaden* 2123 (SING).

Phuket, Khaw Pok Hill: *Haniff & Nur SF* 3822 (SING).

Trang Huay Yawt, Thale Sawng Hawng, common on sandy banks of freshwater lake, 100 m: *Smitinand* 3027 (RFD 17250) (L).

Satul, heaths: *Ridley* 14814, 14815 (SING).

Kastang: *Haniff & Nur SF* 3708 (SING).

DISTR. India, Tonkin, Malay Peninsula, Sumatra, N. Australia.

146. *R. RUGOSA* (Vahl) Gale, Rhodora 46: 275, t. 835, f. 1A—B. 1944.

—*Schoenus rugosus* Vahl, Eclog. Am.: 5. 1798. — *R. glauca* Vahl, En. 2: 233, 1806; F.B.I.: 671. — *R. chinensis* (vix Nees) Boeck., Linnaea 37: 356, 1873. — *R. glauca* Vahl var. *chinensis* Clarke, Fl. Br. Ind. 6: 672, 1893; Mat. F.M.P.: 84; Ridl. 1911: 224; F.M.P.: 165.

S.E.—Loei, Phu Krating, common in savannah, 1000—1300 m: *Smitinand* 1951, 2109 (L); Kao Krating, in marshy ground, 1200 m: *Kerr* 8724 (BK).

E.—Surin, Sangka, in open marshy ground, 300 m: *Kerr* 5309 (BK).

Trat, Kao Kuap, in open grassy ground, 600 m: *Kerr* 17729 (BK).

PEN.—Langsuan, by pool in open scrub, under 50 m: *Kerr* 11922 (BK).

Krabi, Ao Lok, in marshy ground, 50 m: *Kerr* 18566 (BK).

Nakawa Srithamarat, Ta Samet, open grassy ground, under 50 m: *Kerr* 14306 (BK).

Songkhla, Salsa Yoi, in sandy ground, under 50 m: *Kerr* 14779 (BK).

Satul, in open marshy ground, 5 m: *Kerr* 13743 (BK).

DISTR. Tropics and subtropics of the whole world.

147. *R. GRACILLIMA* Thwaites, En. Pl. Zeyl.: 435, 1864; F.B.I.: 671;

*Kern*: 649.

S.E.—Loei, Phu Krating, common in savannah, wet localities, 1300 m: *Smitinand* 1809 (L), 5848 (RFD 21587) (L).

DISTR. Tropical Africa, S.E. Asia, Sumatra, New Guinea, N. Queensland.

### XIII. TRICOSTULARIA Nees

148. *T. UNDULATA* (Thwaites) Kern, Act. Bot. Neerl. 8: 267, 1959.

—*Cladium undulatum* Thwaites, En. Pl. Zeyl.: 353, 1864; F.B.I.: 674; Mat. F.M.P.: 86; Ridl. 1911: 224; F.I.C.: 152, f. 19, 2—8; F.M.P.: 167. —

*Cladium pulchrum* Ridl., Journ. Fed. Mat. St. Mus. 6: 192, 1915; F.M.P.: 167.

- PEN.—Surat, Kanchanadit, in savannah, 5 m: *Kerr 18138* (BK, BO), *18148* (BK).  
 Nakawn Srithamarat, Ta Samet, under 50 m: *Kerr 14279* (BK).  
 Pattani, in sandy ground, 25 m: *Kerr 7241* (BK, SING).  
 Satul, in open grassy ground: *Kerr 18718* (BK); Satul, heath: *Ridley 150* (SING).

DISTR. From N. Australia to Ceylon and Indo-China.

#### XIV. SCHOENUS L.

149. *S. CALOSTACHYUS* (R. Br.) Poir., Enc. Suppl. 2: 251. 1811; F.B.I. 673; Mat. F.M.P.: 85; F.I.C.: 150, t. 1, f. C, 6—9; F.M.P.: 166. — *Chlorospora calostachya* R. Br., Prodr.: 233. 1810.
- PEN.—Ranawng, Kaw Chang, on sandy ground in *Melaleuca* forest, 5 m: *Kerr 18137* (BK, BO, K).  
 Surat, Kantuli, near sea: *Pat 4164* (BK); Kanchanadit, in savannah, 5 m: *Kerr 18137* (BK, BO, K).  
 Nakawn Srithamarat, Ta Samet, common in marsh, under 50 m: *Kerr 14278* (BK).  
 Pattani, on open grassy sandy ground, in loose tufts, near sealevel: *Kerr 700* (BK, K).

DISTR. From Australia and Micronesia to Thailand and the Riu Kiu Islands.

150. *S. FALCATUS* R. Br., Prodr.: 232. 1810; F.I.C.: 150. — *Lophocarpus tonquinensis* Boeck., Allg. Bot. Zeitschr. 2: 111. 1896. — *Neolophocarpus tonquinensis* (Boeck.) Camus, Fl. Gén. I.-C. 7: 149. 1912, quod syn.
- E.—Ubon, Muang Samsip, in tufts on saline sand, 100 m: *Kerr 8355* (BK).  
 S.W.—Kin Savok, c. 120 km NW of Kanbari, 100—150 m: *Kosterwax 1137* (Kor Noi R. basin exp. 1946) (BO, L).  
 PEN.—Surat, Nawng Wai, in savannah, 50 m: *Kerr 12269* (BK, BO, L).  
 Satul, in savannah, 50 m: *Kerr 14411* (BK).

DISTR. From Northern Australia and the Solomon Islands to Thailand, Formosa, and the Riu Kiu Islands.

In typical *Schoenus falcatus* the nuts are distinctly scrobiculate, i.e. marked by numerous shallow pits. In *Kerr 8355* they are markedly different by the inflated epidermal cells which give a bullate appearance to their surface. The specimen may represent a separate race of *S. falcatus*.

#### XV. GAHNIA J.R. & G. Forst.

151. *G. TRISTIS* Nees in Hook. & Arn., Bot. Beech. Voy.: 228. 1837; Mat. F.M.P.: 98; F.M.P.: 168; Raymond, Mém. Jard. Bot. Montréal n. 54: 75. 1959. — *G. boniana* Boeck., Allg. Bot. Zeitschr. 2: 141. 1896; F.I.C.: 154. — *G. stricta* Boeck., l.c. 142; F.I.C.: 155.

N.E. — Loei, Phu Krading, near Rest House, common by stream along edge of evergreen jungle, 1300 m: *Smitinand 1896 (RFD 19935)* (L).

S.E. — Trat, Kao Kuap, rocky ground in light evergreen forest, 700 m: *Kerr 17778 (BK)*.

Trang, Sikao, in open evergreen forest on sandy soil, under 50 m: *Kerr 19006 (BK)*.

Songkhla, Kampangpet, in open forest on sandy ground, 50 m: *Kerr 15915 (BK)*.

DISTR. S. China, Riu Kiu Islands, Indo-China, W. Malaysia. The record "Moluccas" (Benl. Bot. Arch. 40: 183. 1940; Raymond, l.c.: 76) refers to a specimen of *G. aspera* (R. Br.) Spreng., a closely related eastern species.

## XVI. SCLERIA Berg.

152. *S. CARICINA* (R. Br.) Benth., Fl. Austr. 7: 426. 1878; F.B.I.: 688; Mat. F.M.P.: 111; Ridl. 1911: 225; F.M.P.: 178. — *Diplacrum caricinum* R. Br., Prodr.: 241. 1810; F.I.C.: 157.

S. — Den Jaya, *Franch 882* (C).

C. — Bangkhen, evergreen forest, marsh: *Mureau 2559* (BM, K).

S.E. — Near Sriracha, rice-fields, very frequent, 10—20 ft: *Mrs Collins 1888* (BM, K), 1821 (BK, K), 1932 (BK, K), 2251 (BM).

Chanthaburi: *Vesterdal 401* (C, SING).

Trat, Hui Reng, in evergreen scrub, 5 m: *Kerr 17593* (BK, BM); Kaw Chang, Salak Kawk, in open scrub, 10 m: *Kerr 9225* (BM, K).

PEN. — Ranawng, La-un, in open grassy ground, 10 m: *Kerr 10349* (BK, BM, K).

Surut, in open scrub: *Kerr 11287* (BM, K); Khun Tale Lake: *Seidenfaden 2335* (C, SING); Kaw Samui, Ban Bua Put, in open scrub, under 5 m: *Kerr 15709* (BK, BM, K).

Phuket, Khaw Pok Hill, in lallang (*Imperata*), 1500 ft: *Haniff & Nur SF 3847* (K, SING).

Satul, heaths: *Ridley 14806* (BM, K, SING); Satul, Khuan Pho, common roadside weed, 20 m: *Kerr 12770* (BK, BM, K).

DISTR. S. and S.E. Asia, Malaysia, tropical Australia.

153. *S. reticulata* (Holtt.) Kern, comb. nov. — *Diplacrum reticulatum* Holtt., Gard. Bull. Sing. 11: 295, f. 6. 1947.

PEN. — Surat, Ban Na: *Yuang 51* (K).

DISTR. Also known from the Malay Peninsula (Pahang, Gua Tipus; type locality), East Bengal, and Lower Burma (Kyaukpyu Isl.).

A very remarkable species, closely similar in habit to *S. caricina*. Holttum distinguishes it by the turbinate, not cylindric, female spikelets with entire glumes, and by the broader reticulate nuts. An additional differentiating character is to be found in the nervation of the glumes of the female spikelets: strongly many-nerved in *S. caricina*, faintly few-nerved

with only the midrib prominent in *S. reticulata*. To me the differences in the nut are inconspicuous if any. The nut in *S. caricina* is very variable and in some specimens I found it hardly distinguishable from that of *S. reticulata*.

154. *S. LEVIS* Retz., Obs. 4: 13. 1786; S. T. Blake, Journ. Arn. Arb. 35: 226. 1954. — *S. hebecarpa* Nees in Wight, Contr. Bot. Ind.: 117. 1831; F.B.I.: 689; Mat. F.M.P.: 113; Ridl. 1911: 225; F.I.C.: 166.  
 N.—Chiengmai, Doi Suthep, in dry deciduous forest, 700 m: Smitinand 3989 (BKF); Kerr 1269 (BM).  
 Kamphaeng Phet, common in teak forest, 200 m: Smitinand 6110 (L).  
 S.E.—Chanthaburi, Makham, common in thickets, 50 m: Smitinand 5745 (L).  
 Trat, among undergrowth along edge of evergreen forest, 50 m: Smitinand 122 (BKF).  
 S.W.—Kanburi, Hin Dat: Put 62 (BK, BKF, BM, BO, L).  
 Brangkasi, along forest paths, 150 m: den Hoed & Kostermans 659 (Kwae Noi R. basin exp. 1946) (L).  
 Near Kin Sayok, c. 120 km NW of Kanburi, mostly dry limestone rocks, bamboo thicket, 100 m: Kostermans 1099 (Kwae Noi R. basin exp. 1946) (BO, L).  
 Ratburi, in deciduous forest, under 50 m: Kerr 9030 (BK, BKF, BM).  
 PEN.—Surat, in evergreen scrub, under 10 m: Kerr 11259 (BK, BKF, BM, BO, L).  
 Satul, heaths: Ridley 14807 (SING).  
 DISTR. From India through S. Asia to Japan and through Malaysia, Queensland and New Caledonia.

155. *S. TERRESTRIS* (L.) Fass., Rhodora 26: 159. 1924; S. T. Blake, Proc. Roy. Soc. Queenal. 62: 89. 1952; Journ. Arn. Arb. 35: 228. 1954. — *Zizaniopsis terrestris* L., Sp. Pl. 2: 991. 1753. — *S. elata* Thwaites, En. Pl. Zeyl.: 32. 1864; F.B.I.: 690; Mat. F.M.P.: 114; F.I.C.: 167; Turr.: 426; F.M.P. 179.  
 N.—Chiengmai, Chiengdao, common in evergreen forests, 1030 m: Khanthachai Du-chuai 215 (RFD 18179) (L).  
 Tak, on road to Ma-sawt, in old clearings, 600 m: Smitinand 5288 (L).  
 Phetchabun, Lomkao, Phuhee, in deciduous forest, 800 m: Smitinand 2810 (det. Raymond).  
 S.E.—Krabin, Bankeng, evergreen forest, 25 m: Marcan 2556 (BM).  
 Chantaburi, Kao Petehakut, 3200 ft: Nai Noe 75 (BM).  
 Kaw Chang, common by stream on sandy bank, 100 m: Smitinand 2173 (L).  
 S.W.—Near Neeckey, near Wangka, 150 m: den Hoed 183 (Kwae Noi R. basin exp. 1946) (K, L), Wichian 323 (L).  
 Prachuap, Kao Luang, in open scrub, 1200 m: Kerr 10852 (BK, BM).  
 Chumpawn, Kao Tong, open scrub, 700 m: Kerr 11536 (BK, BM, K).  
 PEN.—Ranawng, Kao Pawta Luang Keo, in open evergreen ridge, 1300 m: Eo 16965 (BM, K); Kao Pawta Chenglong, in evergreen scrub, 900 m: Kerr 1671 (BK, BM, K).

Langsuam, Kao Nom Suo, in evergreen forest, 900 m: *Kerr 12043* (BK, BM).  
 Pang-nga, Kao Katakwam, in evergreen forest, 900 m: *Kerr 18433* (BK, BM).  
 Krahl, in open spots, 1200 m: *Kerr 18709* (BK, BM).

Trang, Chawng, by stream in evergreen forest, 100 m: *Smitinand 4079, 4080* (L).  
 Pattani, Betong, Gunong Ina, in evergreen forest, 1200 m: *Kerr 7556* (BK, BM);  
 Bachaw, in evergreen forest, 500 m: *Kerr 7188* (BK, BM).

DISTR. From Ceylon and India to China, Formosa, the Riu Kiu Islands,  
 and through Malaysia to Australia.

156. *S. KERRII* Turr., Kew Bull. (1910) : 384. 1910; F.I.C.: 165; Turr.: 426.

N.—Chiangmai, Doi Suthep, in jungle, 1000—1500 ft: *Kerr 1239* (K; type).  
 Chiangrai (?), lateritic soil, open country, 1000 ft: *Howard 20* (K).

NE.—Loei, Wang Saphung, Sithan, swamp, 300 m, local name *yah poom khao*: *Dee et al. (RFD 4813)* (BKF, MTJB).

DISTR. Annam.

157. *S. CILIARIS* Nees in Wight, Contr. Bot.-Ind.: 117. 1834; S.T. Blake, Journ. Arn. Arb. 35: 227. 1954. — *S. chinensis* Kunth, En. 2: 357. 1837; F.B.I.: 690; Mat. F.M.P.: 114; F.I.C.: 167; F.M.P.: 180. — *S. bancana* Miq., Fl. Ind. Bat., Suppl.: 602. 1861; F.B.I.: 693; Mat. F.M.P.: 112; Ridl. 1911: 225; F.I.C.: 168; F.M.P.: 178. — *S. bancana* Miq. var. *nana* Ridl., Journ. Str. Br. Roy. As. Soc. no 59: 225. 1911; F.M.P.: 178.

SE.—Chanthaburi, Makham, common in thickets, 50 m: *Smitinand 5748* (L); common in deciduous forest, 100 m: *Smitinand 4050* (L, MTJB); Trat, Kao Kuap, in light evergreen forest, 600 m: *Kerr 17748* (BK, BM).

PEN.—Surat, Kantuli: *Put 4169* (BK, BM).

Trang, Huay Yawt, Thale Sawng, very common in evergreen forest, 100 m: *Smitinand 4137* (L).

Songkhla, Tepa, in open waste ground: *Kerr 14701* (BKF, BM).

Satul, in scrub, 5 m: *Kerr 15726* (BK, BM, K, P); Satul, heaths: *Ridley 14804* (type collection of *S. bancana* var. *nana* Ridl.) (K, SING).

DISTR. From Burma through Thailand and Indo-China to S. China;  
 Malaysia, Solomon Islands, tropical Australia.

158. *S. OBLATA* S. T. Blake, Blumea 11, ined. 1961. — *S. levis* (non Retz.) F.B.I.: 694; F.K.C.: 93; Will.: 225; Mat. F.M.P.: 111; Hoss. 1910: 463; F.I.C.: 169; Ridl. 1920: 124; F.M.P.: 177.

C.—Bangkok, waste ground in forest garden: *Kerr 8974* (BK, BKF, BM, K, SING); *Murace 848* (BM).

SE.—Trat, Koh Chang, edge of water: *Eryl Smith 201* (BK, BKF); Khlawng Nonst, in open rather marshy ground, under 10 m: *Kerr 9203* (BK, BKF, BM, K); Lem Dan: *Joh. Schmidt 230* (C); Kaw Kut, common in coconut plantations, 5 m: *Smitinand 5706* (L).

S.W. — Bangtaphan, in marshes: *Keith 232* (SING).

PEN. — Langsuan, by pond in open scrub, under 50 m: *Kerr 11921* (BK, BKF, BM, K). Nakawn Srithamarat, Tung Song, Ao Wang Kram, in valley: *Rabil 277* (BKF, BM).

Songkhla, Tepa, in scrub: *Kerr 14664* (BK, BKF, BM); Koh Gah: *Kloss 6605* (K). Satul, in scrub, 5 m: *Kerr 18726 A* (K).

DISTR. Ceylon, Assam, Burma, Indo-China, Malaysia.

The name *Scleria levis* Retz. has wrongly been applied to this well characterized species, for which the new name *Scleria oblata* will be published by Dr. S. T. Blake, who pointed to the confusion (Journ. Arn. Arb. 35: 227. 1954).

*With regard to the name *Scleria levis**

159. *S. POAEFORMIS* Retz., Obs. 4: 13. 1786. — *S. oryzoides* Presl, Re Haenk. 1: 201. 1828; F.B.I.: 691; Mat. F.M.P.: 110; Ridl. 1911: 225; F.I.C.: 164; F.M.P.: 177; Kill.: 57.

E. — Korat, Ban Chumsaeng, in marsh: *Noi Noe 263* (BK, BM, K); Pak Chawng, in stagnant pools, 300 m: *Kerr 20461* (BK, BM).

S.E. — Chanburi, common in wet places, 120 m: *Smitinand 4069* (L).

S.W. — Chumphawn, Ban Tung Maha, in marshy ground, near sealevel: *Kerr 1190* (BK, BM, K, L).

PEN. — Surat, Kanchanadit, marshy ground in savannah, 5 m: *Kerr 18191* (BK, BM, K).

Songkhla, Tepa, in open swampy ground, under 50 m: *Kerr 14704* (BK, BM). Pattani, Bang Nana, very common in open marshy ground, near sealevel: *Kerr 7158* (BK, BM, K).

Lake Ta Lei Nawn: *Gwynne-Vaughan 254* (K, L).

Klawng Rangsit, Tangaburi, near canals, used for building matting huts: *Ladei 11* (BK, K).

DISTR. Tropical E. Africa; from India and Ceylon to S. China and tropical Australia.

160. *S. PSILORHIZA* Clarke, Fl. Br. Ind. 6: 691. 1894; F.I.C.: 164.

N. — Nakawn Sawan, Hua Wai: *Pat 4091* (BK, BM).

C. — Saraburi, Menam Sak, among grass in savannah: *Kerr 7034* (BK, BM, K, SING).

DISTR. Very local and scattered from India through Thailand and Malaysia to N. Australia.

161. *S. SCROBICULATA* Nees & Mey. ex Nees in Wight, Contr. Bot. Ind.

117. 1834.

E. — Korat, Katok, by stream in evergreen forest, 400 m: *Kerr 8157* (BK, BM).

C. — Bangkok, in waste shady ground, tidal: *Kerr 4069* (BM, K).

DISTR. From S. E. Asia through Malaysia to Queensland and Samoa.

*Scleria acrobiculata* and *S. purpurascens* are so closely related that I feel inclined to treat them as races of a single species.

162. *S. PURPURASCENS* Steud., Syn. 2: 169. 1855. — *S. multifoliata* Boeck., Linnaea 38: 510. 1874; F.B.I.: 693; F.K.C.: 93; Willd.: 225; Mat. F.M.P.: 112; Hoss. 1910: 463; F.I.C.: 169; F.M.P.: 178.

S.E. — Koh Chang, jungle near Klong Munsé, gravel: Joh. Schmidt 273 (C).

PEN. — Ranawng, Lam Lieng, in evergreen forest, 50 m: Kerr 16393 (BK, BM, BO, K).

Surat, in evergreen scrub, under 10 m: Kerr 11256 (BK, BM); Bandon River: Seidenfaden 2159 (C, SING).

DISTR. Burma, Indo-China, Malaysia.

163. *S. SUMATRENSIS* Retz., Obs. 5: 19, t. 2. 1789; F.B.I.: 693; Mat. F.M.P.: 113; F.I.C.: 170; F.M.P.: 179.

S.E. — Koh Chang, Lem Dan, plains: Joh. Schmidt 89 (C).

PEN. — Surat, Khun Talae Lake, swamp: Seidenfaden 2319 (C); Bandon, along a Klong: Seidenfaden 2092 (C).

DISTR. From Ceylon and India to Formosa and Queensland.

164. *S. CORYMBOSA* Roxb., Fl. Ind. ed. 2, 3: 574. 1832; F.B.I.: 686; Mat. F.M.P.: 109, p.p.; F.M.P.: 176, p.p. — *S. ridleyi* Clarke, Fl. Br. Ind. 6: 686. 1894; Mat. F.M.P.: 109; F.I.C.: 160.

E. — Sangka-Samsawng, by stream in scrub jungle, 300 m: Kerr 8801 (BK, BM, P).

SE. — Trat, Talaum, glade in evergreen forest, under 50 m, local name *ya kom bang*: Kerr 17863 (BK, BM).

PEN. — Ranawng, La-un, marshy ground in evergreen forest, 10 m: Kerr 16483 (BK, BM, BO, K, L).

Hnopp: Haniff & Nur SF 4634 (K, SING).

DISTR. From India and Ceylon to S. China, W. and N.W. Malaysia.

165. *S. PARVULA* Steud., Syn. 2: 174. 1855. — *S. tessellata* (non Willd.) F.B.I.: 686, p.p.; F.I.C.: 162.

N.E. — Loei, common in savannah, 1300 m: Smitinand 5021 (L).

DISTR. Tropical Africa; from Nepal, India, and Ceylon to S. China, Japan, Korea, and Malaysia.

166. *S. BIFLORA* Roxb., Fl. Ind. ed. 2, 3: 573. 1832; F.B.I.: 687; Mat. F.M.P.: 114; F.I.C.: 163, f. 21, 7; F.M.P.: 180.

PEN. — Phang-Nga, Pulau Teban: Haniff & Nur SF 3620 (K, SING).

DISTR. S.E. Asia, from Ceylon and India to S. China, Formosa, the Riu Kiu Islands, eastwards to the Philippines and Celebes, southwards to Java.

ssp. *ferruginea* (Ohwi) Kern, stat. nov. — *S. ferruginea* Ohwi, Ad. Phytotax. Geob. 7: 37. 1938.

S.E. — Prachinburi, Aran Prathet, in open grassy low-lying ground, 50 m: Kerr 1516 (BK, BM, K).

DISTR. of the subspecies: Liu Kiu Islands, Formosa, Indo-China, Malay Peninsula.

167. *S. MIKAWANA* Makino, Bot. Mag. Tokyo 27: 57. 1913. — *S. tessellata* (non Willd.) F.B.I. 686, p.p.

N.E. — Loei, Phu Krating, common by stream and in damp place, 1300 m: Smitinand 1845 (herb. Raymond).

DISTR. Tropical Africa; from Ceylon and India to Japan and New Guinea, very scattered.

168. *S. RUGOSA* R. Br., Prodr.: 240. 1810. — *S. flaccida* Clarke, Fl. Br. Ind. 6: 688. 1894, non Steud. 1855. — *S. zeylanica* (non Poir.) F.B.I.: 687, p.p.; Mat. F.M.P.: 110; F.I.C.: 163; F.M.P.: 177.

S.E. — N. N. Kheo, Sriracha forest, 400 m: Mrs Collins 1900 A (BK, K).

DISTR. From Ceylon and India to S. China, Formosa, and Japan through Malaysia to N. and N. E. Australia and New Caledonia.

169. *S. THWAITESIANA* Boeck., Linnaea 38: 454. 1874. — *S. zeylanica* (non Poir.) F.B.I.: 687, p.p. (quoad specim. ceyl.).

N.E. — Loei, Phu Krating, 1300 m: Smitinand 1428 B (BKF).

S.E. — Trat, Kao Saming, in open grassy ground, 20 m: Kerr 9484 (BK, BM, E. PEN). — Surat, Kantuli: Put 4163 (BK, BM).

Between Trang and Krabi, common in wet locality, 100 m: Smitinand 413 (RFD 16865) (BKF, L).

DISTR. Ceylon (type locality).

Clarke and subsequent authors reduced this species to the synonymy of *S. rugosa* R. Br. (*S. zeylanica* sensu Clarke). In my opinion it is readily distinguishable by the following characters: Stems strictly erect, very slender. Leaves rigid, canaliculate, narrow, 1—2 mm wide, glabrous. Terminal panicle not pseudo-lateral, its bract not or only slightly overtopping the inflorescence. Peduncles of the lateral panicles setaceous. Nut obtusely trigonous, smooth, very shining. Disk thin, hardly lobed, not cellular-glandular.

170. *S. NEESII* Kunth, En. 2: 358. 1837; F.B.I.: 688; F.I.C.: 164 t. 21, 6. — *Hypoporum capitatum* Nees, Edinb. New Phil. Jour. 17, no 34. 267. 1834. — *S. capitata* (Nees) H. Pfeiff. in Fedde, Rep. 26: 262. 1920, in syn., non Willd. (1805).

S.E.—Chanthaburi, common in wet sandy soil in open *Dipterocarpus* forest, under 100 m: Smitinand 3569 (L); Makham, in open deciduous forest, 100 m: Smitinand 3041 (RFD 12618) (L).

PEN.—Surat, open scrub, under 10 m: Kerr 11283 (K).

Between Trang and Krabi, in wet localities, 100 m: Smitinand 4148 (RFD 16883) (L).

DISTR. Ceylon, Cochinchina, Annam, Malay Peninsula.

171. *S. TONKINENSIS* Clarke, Kew Bull. add. ser. 8: 57. 1908.

N.E.—Nakawn Panom, Ta Uten, by road-side in mixed forest, sandy soil, 1200 m: Kerr 21374 (BK, BM, K).

E.—Korat, Ban Chumsaeng: Nai Noe 262 (BK, BM, BO, K).

Srisaket, Kanthraram, common in scrub jungle, under shade, 100 m; used locally for labouring mothers: Smitinand 594 (RFD 4932) (K, L).

SE.—Prachinburi, Aran Prathet, railway embankment, 50 m: Kerr 19625 (BK, BO, K).

Chanthaburi, Makham, Okphok, in dry mixed deciduous forest, 100 m: Smitinand 3818 (RFD 22095) (BKF); Chanthaburi, Makham, common in the plain, along edge of thickets, 20 m: Smitinand 5758 (RFD 21502) (L).

Trat, Kao Kuap: Put 2912 (BK, BM).

S.W.—Chumphawn, Bang Son: Put 1590 (BK, BM, K); same locality, sandy heath near coast: Haniff & Nur SF 4221 (K, SING).

PEN.—Surat, in evergreen scrub: Kerr 11249 (BK, BM, K); Kantuli: Put 4166 (BK, BM); Tha Phet, common along edge of water-hole, 50 m: Smitinand 5617 (L).

DISTR. Also in Tonkin (type locality), Cambodia, Cochinchina, and Hainan.

172. *S. LITHOSPERMA* (L.) Sw., Prodr.: 18. 1788; F.B.I.: 685; Mat. F.M.P.: 108; Ridl. 1911: 225; 1912: 63; F.I.C.: 161, f. 21, 5; F.M.P.: 176; Key. 1957: 62. — *Scirpus lithospermus* L., Sp. Pl. 1: 51. 1753.

E.—Korat, Hui Taleng: Put 2227 (BK, BM).

C.—Chaibadan, Dong Pya Yen, in mixed forest, 50 m: Kerr 7995 (BK, BM, K).

SE.—Trat, Kaw Chang, Khlawng Son, along edge of scrub jungle, 10 m: Smitinand 5675 (L).

Chantaboon: Vesterdal 478 (C, SING).

Prachinburi, Nawng Plu, Bandan, 200 ft: Mrs Collins 848 (BK, K); Sriracha: Put 450 (BK, BM, BO, K); N. N. Kheo-Sriracha forest, 400 ft: Mrs Collins 1900 (BK, K).

S.W.—Prachinburi, Ban Nawng Kang, in dry evergreen forest, under 50 m: Kerr 16929 (BK, BM).

PEN.—Surat, Kantuli: Put 4158 (BK, BM); Kaw Tao, in dry evergreen forest, 10 m: Kerr 11162 (BK, BM, BO, K); same locality, in light evergreen forest, 300 m: Kerr 16039 (BK, BM).

Songkhla, Kampangpet, in evergreen forest, 50 m: Kerr 15975 (BK, BM, K).

Pattani, Banang Sta, in banana plantation, 50 m: Kerr 7408 (BK, BM, K).

Satul, Butang, on rocky ground, 60 m: *Kerr 14053* (BK, BM); Adang, in evergreen forest, 5 m: *Kerr 13993* (BK, BM); Bukit Raja Wang: *Ridley 15805* (SING).  
DISTR. Pantropic.

var. *LINEARIS* Benth., Fl. Austr. 7: 430. 1878. — *Hypoporum roxburghianum* Nees ex Boeck., Linnaea 38: 452. 1874. — *Hypoporum roxburghii* Nees ex Clarke, Fl. Br. Ind. 6: 686. 1894. — *S. lithosperma* (L.) Sw. var. *roxburghii* Clarke, Fl. Br. Ind. 6: 686. 1894; Mat. F.M.P.: 105. N.E. — Nakawn Panom, Muk Dahan, Pu Mano, in deciduous forest, 300 m: *Kerr 2121* (BK, BO, K).

S.E. — Trat, Kaw Chang, Salak Kawk, in loose damps, open deciduous forested slope near sea: *Kerr 6891* (BK, BM, K, NY, P).

DISTR. of the variety. From India and Ceylon to Indo-China, and through Malaysia to tropical Australia.

173. *S. PERGRACILIS* (Nees) Kunth, En. 2: 354. 1837; F.B.I.: 685; F.I.C.: 160, f. 21, 1-3; Turr.: 426. — *Hypoporum pergracile* Nees, Edinb. New Phil. Journ. 17, no. 34: 267. 1834; in Wight, Contr. Bot. Ind.: 118. 1834.

N. — Doi Suthep, 1200—1500 ft: *Kerr 1428* (BK, BM, P); in open marsh, 2200 ft: *Kerr 2647* (BM, K).

N.E. — Loei, Phu Krading, common among grasses in savannah, 1300 m: *Smitinand 1905* (herb. Raymond).

S.E. — Krabin, Watana, in open grassy forest, 50 m: *Kerr 9788* (BK, BM, K). Baw Luang, in open marshy ground: *Kerr 6266* (BK, BM, K, P).

DISTR. Tropical Africa; from India and Ceylon to S. China and Malaya.

174. *S. BENTHAMII* Clarke, Kew Bull., add. ser. 8: 58. 1908. — *S. khasiana* Clarke in Hook. f., Fl. Br. Ind. 6: 692. 1894; Journ. Linn. Soc., Bot. 34: 102. 1898, non Boeck. (1890).

N. — Chiengmai, Doi Suthep, common in open grassy jungle, 900—1000 m: *Kerr 3358* (BM).

N.E. — Loei, Phu Krading, common in savannah, 1300 m: *Smitinand 1942* (herb. Raymond).

DISTR. Khasia, Indo-China, Luzon, Queensland.

*Scleria thomsoniana* Boeck., Linnaea 38: 479. 1874, and *S. khasiana* Boeck., Cyp. Nov. 2: 29. 1890, were based on the same collection, viz *Scleria sp. 12*, Herb. Ind. Or., Hook. f. & Thoms. As the latter binomial is nomenclaturally superfluous, Clarke used the same combination again for a quite different species. This procedure is contrary to the present-day Code. As *Scleria benthamii* Clarke, thus far only known from a single Australian collection (Queensland, Brisbane R., F. v. Müller 61, K) appears

to be conspecific with the Asiatic plants, this is the earliest legitimate name of the species.

*Scleria benthamii* is most easily recognizable by the characteristic shape of the hypogynous disk, which reminds one of a tricorn hat. In habit it is extremely similar to *S. levis* Retz., but easily recognizable by the more distinctly reticulated, hardly apiculate nut, and the different shape and texture of the hypogynous disk. In *S. levis* the lobes of the disk are thin, lanceolate, brown, 1—1½ mm long, in *S. benthamii* thick, much shorter, ovate, whitish, with a minute, erect apicula at the top. Probably there is also a difference in the contraligule (i.e. the tongue-shaped prolongation at the top of the leaf-sheath on the anterior side). In *S. levis* this contraligule has (always?) a small, easily overlooked membranous appendage at the top, which is lacking in the few specimens of *S. benthamii* I have seen.

*Scleria benthamii* is also in Indo-China (Tonkin, route de Laokay à Chapa, savannes herbeuses, 1000 m: Pételot 8548, herb. Raymond).

#### SPECIES RECORDED FOR THAILAND NOT SEEN BY ME:

1. *Cyperus eleusinoides* Kunth; Turr.: 422.
2. *Cyperus exaltatus* Retz. var. *digynus* Williams: 224. — According to Kükenthal, Pflanzenr., Heft 101: 71. 1935 = *C. alopecuroides* Rottb. Kükenthal did not see the collection on which the record was based.
3. *Cyperus kurzii* Clarke; Koy. 1955: 209; 1957: 28.
4. *Cyperus ohwi* Kük.; Koy. 1955: 205; 1957: 24.
5. *Fimbristylis haspaniformis* Koyama, Contr. Inst. Bot. Univ. Montréal no. 70: 45. 1957.
6. *Fimbristylis tenera* R. & S.; Ost.: 719; Hoss. 1910: 462; F.I.C.: 112.
7. *Mariscus clarkei* Turr.: 422. Kükenthal, Pflanzenr., Heft 101: 518. 1936, reduced this to a variety of *Cyperus cyperoides* (L.) O. Kuntze.
8. *Scirpodendron ghaeri* (Gaertn.) Merr. (*S. costatum* Kurz; Ridl. 1911: 225). "Common in tidal swamps, Setul". I have not seen any specimen from Satul in the Singapore Herbarium.
9. *Scirpus fluitans* L.; Koy. 1957: 61.
10. *Scirpus triangulatus* Roxb.; Koy. 1957: 59. I doubt whether this is specifically distinct from the widely distributed and extremely polymorphic *S. mucronatus* L.
11. *Scirpus triquetus* L.; Koy. 1957: 62.

## INDEX TO COLLECTORS' NUMBERS

The first number of each pair is the collector's number, the second refers to the number of the species in the text. See also Addenda.

- Amsat* 57: 53.  
*Bloembergen*, S. 604: 94 — *Boonag*, see  
*Bunnak* — *Boonbrahm* 8: 31; 9: 128 —  
*Bunnak* 5: 131; 135: 45; 165: 8; 174:  
 61; 195: 145; 197: 139; 198: 19; 205:  
 35; 211: 112; 212: 42; 217: 106; 218:  
 93; 231: 116; 233: 7; 464: 113; 465:  
 106; 840: 137; 841: 116; 844: 53.  
*Colani*, E. 4055: 71 — *Collines*, Mrs D. J.  
 845: 7; 846: 105; 848: 172; 849: 106;  
 850: 140; 1888: 129; 1888: 152; 1889:  
 80; 1891: 105; 1893: 31; 1895: 111;  
 1896: 14; 1900: 172; 1900A: 168; 1902:  
 9; 1908: 80; 1931: 152; 1932: 152;  
 2251: 152.  
*Dee* 2: 116; 3: 48; 36: 64; 37: 10; 62:  
 156; 158: 59; 158: 102; 327: 53; 9135:  
 140 — *Din* 10: 59.  
*Franck*, C. W. 439: 110; 993: 152; 1826:  
 125.  
*Gwynne-Vaughan* 255: 159; 586: 92.  
*Haniff & Nur* (all SF numbers) 3620:  
 166; 3831: 80; 3805: 127; 3822: 145;  
 3847: 152; 4084: 164; 4221: 171; 4363:  
 74; 4708: 145 — *Hoed*, G. den 122: 94;  
 131: 105; 133: 155; 139: 97; 145: 39;  
 149: 140; 149: 42; 150: 65; 296: 102;  
 211: 95; 220: 19; 283: 105; 325: 129;  
 594: 64; 923: 105 — *Hoed*, G. den &  
*Kostermans*, A. 457: 105; 650: 154;  
 686: 131; 695: 87; 696: 94; 697: 89;  
 888: 140; 968: 71; 985: 92 — *Hosseus*  
 101: 24; 104: 125; 182: 144; 183a: 92;  
 481: 10 — *Howard*, A. J. 20: 156; 27:  
 142.  
*Keith*, A. 252: 158; 341: 140; 388: 33;  
 406: 95; 432: 106; 482: 35; 494: 106 —  
*Kerr*, A. F. G. 832: 55; 862: 125; 1239:  
 150; 1269: 154; 1428: 173; 1595B: 144;  
 2647: 173; 3358: 174; 3860: 21; 3864:  
 123; 3865: 129; 3877: 20; 3888: 128;  
 4069: 161; 4196: 106; 4708: 64; 6053:  
 132; 6104: 68; 6248: 58; 6266: 173;  
 6472: 138; 6629: 139; 6706: 128; 6707:  
 60; 6720: 9; 6721: 42; 6722: 21; 673:  
 20; 6764: 119; 6767: 60; 6800: 7; 681:  
 35; 6864: 60; 6891: 172 var.; 692:  
 111; 6942: 91; 6973: 158; 6975: 6;  
 6976: 110; 6977: 111; 6978: 60; 698:  
 100; 6983: 64; 6984: 96; 6985: 9;  
 6986: 141; 6990: 94; 7007: 109; 701:  
 94; 7022: 47; 7023: 134; 7035: 19;  
 7041: 106; 7054: 65; 7063: 31; 707:  
 149; 7086: 26; 7134: 41; 7156: 19;  
 7188: 155; 7230: 42; 7231: 27; 723:  
 81; 7241: 148; 7246: 85; 7261: 12;  
 7294: 98; 7308: 41; 7309: 143; 733:  
 35; 7408: 172; 7505: 2; 7546: 15;  
 7726: 85; 7738: 52; 7739: 45; 774:  
 36; 7994: 110; 7995: 172; 8004: 2;  
 8007: 120; 8008: 8; 8009: 95; 801:  
 129; 8064: 13; 8085: 113; 8087: 19;  
 8111: 125; 8147: 59; 8157: 161; 818:  
 7; 8160: 35; 8177: 7; 8190: 74; 822:  
 36; 8241: 82; 8301: 164; 8308: 8;  
 8309: 146; 8323: 22; 8355: 150; 839:  
 113; 8477: 48; 8554: 105 var.; 861:  
 52; 8684: 48; 8704: 57; 8724: 14;  
 8832: 106; 8862: 51; 8970: 79; 897:  
 118; 8988: 145; 9036: 154; 9037: 4;  
 9040: 85; 9198: 19; 9200: 64; 921:  
 158; 9225: 152; 9245: 59; 9266: 12;  
 9287: 98; 9389: 102; 9484: 169; 977:  
 35; 9785: 173; 10057: 104; 10712: 4;  
 10713: 92 ssp.; 10714: 14; 10715: 19;  
 10720: 39; 10724: 134; 10843: 12;  
 10852: 155; 10873: 105; 10919: 12;  
 10922: 123; 10958: 47; 10973: 8;  
 10981: 85; 11033: 102; 11057: 11;  
 11084: 92 ssp.; 11085: 90; 11100: 2;  
 11101: 96; 11105: 29; 11157: 91; 1116:  
 172; 11176: 105; 11180: 131; 1117:  
 128; 11241: 134; 11245: 62; 11248: 14;  
 11249: 171; 11251: 48; 11256: 12;  
 11259: 154; 11262: 93; 11264: 10;  
 11283: 170; 11284: 118; 11287: 12;  
 11288: 118; 11289: 31; 11305: 11;  
 11306: 129; 11341: 64; 11344: 7;

- 11597: 1; 11598: 159; 11400: 61;  
 11402: 80; 11403: 18; 11415: 143;  
 11536: 155; 11539: 34; 11606: 7;  
 11706: 19; 11707: 26; 11711: 111;  
 11803: 80; 11876: 67; 11895: 7; 11921:  
 159; 11942: 146; 12005: 2; 12038: 7;  
 12043: 185; 12063: 111; 12269: 150;  
 12305: 78; 12309: 83; 12310: 75; 12430:  
 41; 12340: 111; 12577: 123; 12584: 47;  
 12586: 135; 12609: 64; 12622: 38;  
 12626: 132; 12726: 131; 12783: 105;  
 12816: 117; 12871: 101; 12872: 65;  
 12884: 7; 12946: 103; 13010: 38; 13014:  
 95; 13030: 60; 13033: 123; 13061: 35;  
 13061: 142; 13067: 49; 13068: 61;  
 13096: 145; 13099: 98; 13101: 159;  
 13109: 143; 13125: 47; 13137: 149;  
 13139: 148; 13142: 63; 13148: 148;  
 13177: 67; 13417: 108; 13418: 91;  
 13442: 103; 13443: 113; 13467: 115;  
 13495: 85; 13498: 144; 13526: 31;  
 13551: 46; 13601: 98; 13624: 33;  
 13649: 143; 13682: 53; 13690: 49;  
 13710: 86; 13718: 148; 13726: 157;  
 13726A: 158; 13730: 81; 13734: 56;  
 13739: 81; 13740: 28; 13742: 85; 13743:  
 146; 13761: 129; 13765: 64; 13770:  
 152; 13788: 8; 13993: 172; 14025: 7;  
 14029: 2; 14039: 123; 14053: 172;  
 14065: 105; 14089: 134; 14107: 36;  
 14120: 57; 14156: 135; 14243: 60;  
 14278: 149; 14279: 148; 14306: 146;  
 14309: 30; 14329: 87; 14385: 134;  
 14411: 150; 14432: 2; 14626: 6; 14684:  
 1598; 14681: 39; 14704: 159; 14710: 53;  
 14713: 61; 14715: 41; 14779: 146;  
 14795: 139; 14993: 3; 15024: 86;  
 15081: 89; 15083: 108; 15113: 111;  
 15142: 17; 15158: 35; 15160: 64; 15215:  
 2; 15687: 132; 15698: 134; 15699: 38;  
 15700: 91; 15705: 60; 15706: 38; 15708:  
 129; 15709: 152; 15711: 9; 15724: 109;  
 15726: 64; 15798: 27; 15828: 48; 15915:  
 151; 15931: 57; 15969: 59; 15973: 172;  
 15977: 36; 15983: 98; 16023: 123;  
 16039: 172; 16100: 79; 16108: 109;  
 16180: 60; 16187: 91; 16188: 135;  
 16240: 18; 16352: 19; 16384: 107;  
 16393: 162; 16449: 152; 16494: 7;  
 16498: 164; 16590: 149; 16592: 81;  
 16593: 8; 16595: 6; 16795: 155; 16948:  
 57; 16963: 155; 17004: 5; 17088: 6;  
 17089: 3; 17108: 1; 17354: 30; 17390:  
 108; 17586: 123; 17593: 152; 17603:  
 5; 17652: 44; 17686: 59; 17729: 146;  
 17748: 157; 17763: 30; 17756: 48;  
 17778: 151; 17787: 7; 17788: 57; 17863:  
 164; 18035: 6; 18051: 99; 18399: 2;  
 18404: 4; 18438: 155; 18464: 57; 18500:  
 6; 18517: 64; 18566: 146; 18709: 155;  
 18872: 38; 19003: 10; 19006: 151;  
 19028: 53; 19050: 36; 19151: 2; 19257:  
 61; 19258: 143; 19259: 53; 19304: 53;  
 19305: 116; 19351: 132; 19536: 30;  
 19407: 4; 19473: 47; 19489: 134; 19522:  
 110; 19524: 37; 19532: 94; 19533: 105;  
 19565: 166 ssp.; 19566: 50; 19568: 12;  
 19572: 66; 19577: 126; 19618: 130;  
 19625: 171; 19673: 102; 19674: 42;  
 19715: 117; 19723: 59; 19727: 102;  
 19776: 48; 20200: 33; 20211: 111;  
 20219: 25; 20461: 159; 20482: 85;  
 20521: 105; 20527: 94; 20557: 105;  
 20651: 144; 20642: 49; 20658: 33;  
 20697: 65; 20700: 109; 21294: 71;  
 21353: 72; 21359: 73; 21361: 70; 21363:  
 122; 21364: 121; 21365: 65; 21366: 69;  
 21374: 171; 21379: 102; 21382: 119;  
 21387: 89; 21388: 102; 21391: 89;  
 21397: 101; 21398: 83; 21409: 15;  
 21416: 136; 21418: 131; 21419: 93;  
 21425: 132; 21426: 134; 21438: 172;  
 var.; 21447: 116; 21451: 49; 21490:  
 106; 21522: 77; 21524: 64; 21528: 47 —  
 Khanthachai Bunchuai 215: 155; 1037:  
 33 — Kloss 6605: 158; — Kostermans,  
 A. 65: 111; 1098: 105; 1699: 154;  
 1100: 105; 1100a: 95; 1132: 34;  
 1137: 150; 1212: 131; 1238: 64; 1239:  
 122; 1268: 143; 1279: 64; 1309: 142;  
 1310: 111; 1319: 110; 1320: 64; 1321:  
 42; 1322: 129; 1324: 142; — Krua Noi  
 River basin expedition 1916 122: 94;  
 131: 105; 133: 155; 139: 97; 145:  
 139; 146: 140; 149: 42; 150: 65;  
 192: 131; 206: 102; 311: 95; 220;

- 19; 233: 105; 323: 155; 325: 129;  
 332: 68; 337: 181; 457: 105; 594: 64;  
 604: 94; 650: 154; 686: 131; 695: 87;  
 696: 94; 697: 89; 888: 140; 923: 105;  
 968: 71; 985: 92; 1098: 105; 1099: 154;  
 1100: 105; 1100a: 95; 1132: 34; 1137:  
 150; 1212: 131; 1238: 64; 1239: 122;  
 1268: 143; 1279: 64; 1309: 142;  
 1310: 111; 1319: 110; 1320: 64; 1321:  
 42; 1322: 129; 1324: 142.  
*Marcus, A.* 658: 104; 848: 158; 1855: 95;  
 1924: 104; 2533: 39; 2556: 39; 2559:  
 152.  
*Noi Noe* 75: 155; 136: 125; 137: 38; 138:  
 113; 195: 54; 196: 42; 197: 183; 198:  
 110; 263: 87; 262: 171; 263: 159 —  
*Nielsen, Einer* 696: 140; 724: 105.  
*Phloeoecidit* 353: 140; 368: 139; 838: 42;  
 847: 113; 852: 35; 858: 139; 867: 64;  
 1293: 33 — *Put* 10: 85; 22: 129; 24:  
 92; 26: 140; 62: 154; 105: 87; 143: 38;  
 289: 47; 440: 172; 516: 7; 1509: 41;  
 1542: 61; 1544: 32; 1546: 80; 1547:  
 123; 1587: 1; 1590: 171; 1593: 8; 1668:  
 48; 1683: 7; 1952: 8; 1978: 111; 1991:  
 142; 2108: 105; 2144: 112; 2145: 102;  
 2227: 172; 2304: 85; 2501: 115; 2599:  
 102; 2630: 102; 2668: 83; 2779: 105  
*var.*: 2757: 75; 2832: 134; 2912: 171;  
 3056: 134; 3093: 8; 3170: 47; 3179:  
 85; 3340: 57; 3387: 47; 4067: 117;  
 4091: 160; 4128: 41; 4181: 93; 4188:  
 172; 4163: 169; 4164: 149; 4166: 171;  
 5169: 157; 4170: 48; 4182: 139; 4183:  
 113; 4207: 40; 4270: 47; 4291: 34;  
 4294: 11; 4315: 105.  
*Rabul* 140: 110; 149: 18; 210: 140; 311:  
 158; 338: 141; 240: 123; 251: 111 —  
*RFD numbers* (*Royal Forest Department Bangkok*) 3582: 59; 3628: 64;  
 3679: 102; 3857: 93; 3858: 138; 4313:  
 64; 4904: 59; 4906: 10; 4907: 116;  
 4908: 53; 4913: 156; 4932: 171; 4934:  
 49; 5135: 137; 5136: 116; 5137: 76;  
 5273: 83; 5653: 48; 6157: 111; 6160:  
 33; 6162: 139; 6179: 10; 6252: 80;  
 6771: 143; 6772: 49; 6774: 61; 6775:  
 80; 6776: 27; 6777: 19; 7064: 33;  
 7665: 25; 8841: 64; 8853: 105; 8860:  
 57; 8958: 140; 9134: 140; 9448: 196;  
 9459: 16; 9518: 61; 9523: 93; 952:  
 106; 9609: 19; 9612: 145; 9623: 36;  
 9647: 116; 10934: 151; 11260: 131;  
 11275: 53; 11540: 32; 11554: 7; 1177:  
 139; 11912: 41; 11913: 41; 12021: 19;  
 12317: 7; 12503: 106; 12515: 113;  
 12618: 170; 12639: 41; 12840: 860 —  
*Satinand* 5949 and 2953; 12612: 31;  
 12833: 35; 12884: 42; 12932: 139;  
 12936: 113; 14419: 64; 14455: 41;  
 14677: 129; 15136: 93; 15137: 93;  
 15139: 131; 16091: 11; 16092: 23;  
 16093: 125; 16094: 112; 16095: 92;  
 16096: 143; 16097: 58; 16098: 66;  
 16194: 1; 16590: 99; 16869: 179;  
 16885: 169; 16886: 35; 16887: 41;  
 16889: 6; 17006: 33; 17248: 30; 17251:  
 145; 17833: 109; 17849: 94; 18179: 155;  
 19723: 125; 19728: 9; 19729: 20; 1977:  
 21; 19740: 111; 19741: 109; 19742: 92;  
 21490: 43; 21491: 53; 21492: 189;  
 21502: 171; 21506: 62; 21509: 110;  
 21510: 52; 21537: 147; 21567: 34;  
 21573: 160; 21617: 143; 23095: 171 —  
*Ridley* 14804: 157; 14805: 172; 14806:  
 152; 14807: 154; 14809: 31; 14811:  
 132; 14812: 134; 14813: 145; 14814:  
 145; 14819: 46; 14820: 36; 14821: 42;  
 14822: 52; 14824: 148; 14829: 109;  
 14831: 111; 14833: 106; 14834: 111;  
 14877: 53; 14878: 61; 15675: 133;  
 15718: 6; 15719: 26; 15720: 33.  
*Schmidt, Joh.* 16: 128; 89: 163; 239: 114;  
 273: 162; 360: 124; 572: 38; 7160: 19;  
 748a: 128 — *Seidenfaden, Gustav*  
 2008: 118; 2039: 100; 2055: 145; 2061:  
 163; 2118: 87; 2159: 162; 2225: 146;  
 2301: 118; 2319: 163; 2326: 50; 2331:  
 152; 2360: 118; 2439 (or 3439?): 125 —  
*SF (Singapore Field) numbers*, see *Hui*  
*niff & Nor* — *Smith, Eryl* 81: 112;  
 109: 88; 224: 9; 339: 141; 301: 158;  
 367: 60; 931: 31; 1017: 102 — *Smith*  
*Hugh M.* 158: 129 — *Smitinand, Te*  
 32: 83; 114: 111; 118: 139; 121: 11;  
 122: 33; 246: 80; 393: 57; 585: 6

- 537; 137; 538; 116; 594; 171; 602; 76;  
 77; 131; 1020; 2; 1123; 25; 1187; 33;  
 1321; 49; 1332; 143; 1333; 154; 1334;  
 61; 1401; 27; 1402; 19; 1403; 80;  
 1428B; 169; 1890; 151; 1905; 173;  
 1917; 52; 1938; 64; 1939; 147; 1941;  
 73; 1942; 174; 1943; 48; 1944; 102;  
 1945; 107; 1946; 111; 1947; 125; 1949;  
 59; 1951; 146; 1952; 120; 1964; 106;  
 1974; 16; 2109; 146; 2172; 7; 2173;  
 155; 2227; 32; 2590; 92; 2591; 33;  
 2592; 93; 2594; 131; 2595; 49; 2600;  
 41; 2610; 155; 2633; 93; 2797; 14;  
 2820; 42; 2949; 111; 2951; 41; 2952;  
 36; 2953; 64; 3024; 1; 3026; 20; 3027;  
 145; 3041; 170; 3347; 129; 3348; 95;  
 3309; 170; 3318; 171; 3858; 112; 3859;  
 11; 3860; 125; 3861; 23; 3884; 106;  
 3897; 85; 3928; 21; 3987; 58; 3989;  
 154; 3998; 143; 3999; 55; 4011; 64;  
 4022; 92; 4050; 157; 4056; 30; 4069;  
 159; 4079; 155; 4080; 155; 4081; 141;  
 4082; 42; 4083; 25; 4085; 97; 4110;  
 6; 4127; 64; 4137; 157; 4147; 169;  
 1118; 170; 4164; 5; 4530; 99; 4538;  
 125; 4553; 109; 4556; 92 ssp.; 4568;  
 116; 4571; 117; 4572; 131; 4572A; 132;  
 1576; 85; 4678; 113; 4605; 65; 4609;  
 89; 4612; 58; 4758; 94; 4792; 47; 5015;  
 146; 3021; 105; 5412; 87; 5413; 109;  
 5414; 111; 5418; 21; 5419; 9; 5420;  
 20; 5617; 171; 5647; 45; 5648; 32;  
 5650; 60 var.; 5651; 91; 5652; 38;  
 5653; 132; 5654; 106; 5655; 128; 5662;  
 6; 5675; 172; 5705; 35; 5706; 158;  
 5707; 132; 5709; 94; 5730; 111; 5739;  
 62; 5741; 52; 5743; 116; 5745; 154;  
 5746; 76; 5747; 119; 5748; 157; 5755;  
 53; 5756; 61; 5757; 49; 5758; 171;  
 5764; 143; 5771; 62; 5836; 116; 5845;  
 120; 5851; 53; 5854; 43; 5858; 102;  
 5888; 77; 5889; 36; 5933; 120; 5934;  
 132; 5944; 84; 5946; 49; 5947; 59;  
 5948; 147; 5949; 53; 5950; 64; 5951;  
 143; 5953; 126; 5958; 155; 6073; 93;  
 6081; 114; 6082; 64; 6083; 56; 6084;  
 77; 6110; 154 — *Suan* 251; 140 — *So-*  
*radet* 81; 64; 112; 138; 113; 93 — *Su-*  
*vatabandhu* 211; 93.
- Thusnee Boonkong* 2; 9.
- Umpai* 73; 83; 112; 110.
- Vannichgorn* 19; 129; 20; 141 — *Vester-*  
*dal* 8J; 118; 8M; 106; 401; 152; 459;  
 100; 463; 26; 478; 172; 489; 110.
- Wanandorn*, see *Winit* — *Wickian* 192;  
 131; 323; 155; 332; 68; 337; 131 —  
*Williams* 8841; 64; 17227; 41; 17237;  
 41; 17275; 106; 17281; 105 — *Winit*  
 777; 18; 1722; 99; 1850; 113; 1887;  
 105.
- Fuang* 51; 153.

## ADDENDA

After this paper had gone to the press I have seen the following collections:

- Bunnak* 220; 102 — *Dee* 120; 162; 1151; 155 — *Franek* 267; 102; 442; 154;  
 1306; 154 — *Haniff & Nar* SF 7456; 152 — *Howard* 15; 154 — *Kerr* 3552; 146;  
 18312; 106; 18439; 2 var. *petiolata*; 20599; 154 — *Kostermans* 59; 139 — *Lakshnakara*  
 47; 41 — *Marcas* 1358; 172; 2699; 90 — *Put* 1511; 162; 1848; 82; 8134; 170;  
 1167; 170; 4387; 48 — *RFD* 9529; 102; 23026; 155 — *Smith, Hugh M.* 157; 102;  
 152; 102 — *Smithinand* 5600; 162 — *Suan* 55; 155 — *Ward, F. K.*, SF 37352; 172.