

## ON STAUDE'S NEW GENERIC NAMES FOR AGARICS

M. A. DONK\*

## SUMMARY

The author concludes that the generic names for agarics first introduced by Staude (1857) have been validly published.

When I drew attention to Staude's forgotten "Die Schwämme Mitteldeutschlands" (1857), I felt obliged to conclude that the new generic names for agarics appearing in that work were validly published (Donk, 1949: 319-320), and I still am fully convinced that this conclusion must be upheld. Rogers (1950: 22) submitted that Staude, although referring to *Collybia* and other genera as G[attungen], did not definitely accept them as genera, since he continued to refer species under them to *Agaricus*, thereby implying that they were only subgenera or sections; and that Staude, therefore, did not validly publish these generic names.

Rogers's conclusion is untenable for several reasons. If his argument became generally accepted, quite a lot of generic names might be murdered by it: the number of generic names established without simultaneously published new combinations with the new generic name, like those without any mention of species, is considerable. Such an attitude would not only be undesirable from a practical point of view, but would also disregard the declared object of the Code to promote stability of nomenclature.

The Code has deliberately and carefully avoided the stipulation that a new generic name requires simultaneously published new combinations; it has even refused to rule that new generic names ought to be associated with binomials, as was stipulated by the former American code! Certainly this was not done solely to accommodate such special cases as, for example, Tournefortian generic names published after 1753 as an overflow from the pre-binomial period: the decision was for general application.

Staude emphatically marked the names in question as generic ones; not only in the introductory pages, but also throughout his more detailed treatment of the agarics he preceded these generic names with "G[attung]" and added generic descriptions. He supplied the information that, "the genera . . . to follow are all according to Fries. Cp. Systema mycologicum

\* Keeper of Herbarium Bogoriense, Kebun Raya, Indonesia, Bogor.

etc.”<sup>1</sup> That he did not make any new combination is incorrect: I already pointed out that (as regards agarics) a couple of recombinations were made under *Pleuropus* and one under *Crepidotus* (Staude, 1857: 71). Under the generic name *Coprinus*, already established and generally accepted in his day, Staude is inconsistent in exactly the same manner as under *Pleuropus*: some specific denominations are combinations with the corresponding generic name, others, with *Agaricus*. Under *Gomphidius*, another previously established generic name, all three specific denominations are combinations with *Agaricus*, but I should feel guilty of deception if I stated that I really believed that Staude did not definitely accept that genus. In any case, Rogers may have to admit that *Crepidotus* was validly published by Staude, for the only species treated by that author bears a duly recombined name. Yet Rogers may still want to argue that by not having ‘definitely accepted’ the other equivalent taxa as genera, Staude thereby also implied that *Crepidotus* and the earlier published generic names *Coprinus*, *Russula*, *Gomphidius*, and other ones were names for subgenera and sections, too. One reason why Staude did not make recombinations may be that the agarics were popularly known as species of *Agaricus*, and another that his plates bearing the combinations with *Agaricus* may have already been finished when he prepared the text.

Rogers has forgotten to tell us of which *genus or genera* Staude intended his ‘subgenera or sections’ to form a part. Perhaps of that author’s “10. Fam. Agaricoideae. Blätterschwämme” or of the subdivisions thereof, “1. Abtheilung. Dermophylli,” “2. Abtheilung. Pleurotinae,” “3. Abtheilung. Cantharellinae,” and “4. Abtheilung. Agaricini”? But then must Staude not be corrected once more? By accepting the view that Staude implied that his genera were only subgenera or sections, must we not also accept the further implication that the family and its *Abtheilungen* were genera or subgenera? Or was Staude after all quite consistent in the ranks he assigned to his groups? Might it not be Rogers who was inconsistent when he magnified something quite unessential and irrelevant into a major objection to valid publication, an omission deliberately permitted by the Code? The Code keeps the two elements Rogers confused distinct. Art. 42 declares Staude’s new generic names unequivocally validly published, while Art. 43 (1) which he invoked deals with quite different matters as may also be gathered from a study of its text and examples. Moreover, Rogers failed to state why, with his interpretation of the facts, these

<sup>1</sup> “Die folgenden Gattungen . . . sind sämtlich nach Fries. Vergl. Systema mycologicum etc.”—Staude (*op. cit.* p. xxv). In his bibliography on pages 139-140 he fully cited both Fries’s “Systema mycologicum” and “Epicrisis.” For an “Extract from Staude’s summary of the classification of agarics,” see Donk (1949: 385-387).

generic names would not stand as validly published alternative names to the implied subgeneric ones.

If Rogers were correct, I fear that the next victim would be Quélet's "Les Champignons du Jura et des Vosges" (1872), and that such statements as the following would become standard practice. Quélet, like Staude, although dividing the agaric "F[amille]" into "Series" and "Genres," did not definitely accept the latter as genera, since he continued to use in connection with *Tricholoma*, *Clitocybe*, *Collybia*, and so on the masculine terminations for his specific epithets in both the text and the index, thereby implying that they were only subgenera or sections of *Agaricus*; and that he, therefore, did not validly publish these generic names. May I also direct attention to Bataille's "Les Bolets" (1908) and Gilbert's monograph "Amanitaceae" (1941)? These two publications may also prove fruitful subjects for elimination of *bona fide* published generic names!

I do not at all admire Staude's book, but I will not pretend to disbelieve in the valid publication of his names when I honestly believe the opposite to be true. Roger's solution asks mycologists to agree upon an unusual convention in connection with one particular case only. As this proposed convention is not supported by the Code, it will for that reason certainly not appeal to all!

Staude's names hardly form an inconvenience, and, apart from the dubious *Pleuropus-Pleurotus* case (cf. Donk, 1949: 302, 348), there is only one generic name that endangers nomenclatural stability, viz., *Derminus* (Fr.) Staude, hence I have already proposed it for rejection in favour of *Pholiota* (Fr.) Kumm. (Donk, 1949: 324, 394).<sup>2</sup> There are no inconveniences at all as to specific names, because the new combinations he made have no undersirable consequences. The fact that such household names in mycology as *Tricholoma*, *Clitocybe*, *Collybia*, and others appear to have been published much earlier than was commonly believed, is a gain.

Recently the Special Committee for Fungi approved by ballot the conservation of the fungus name *Tricholoma* with '(Fr.) Kummer' (1871: 25, 129) as the author's citation. If its members were genuinely convinced from a taxonomic point of view that this citation is preferable to '(Fr.) Staude' (1857: xxviii, 125), the name *Tricholoma* (Fr.) Staude must be added to *nomina rejicienda* as an earlier homonym. It may well be however, that they acted under the erroneous impression that Staude's generic names were not validly published. Since Staude's concept of the taxon is

<sup>2</sup> This proposal still stands.

the same as Fries's (1838) and Kummer's and includes nothing objectionable, to ignore *Tricholoma* (Fr.) Staude is to misrepresent the facts. Before the proposal is finally accepted, therefore, this error should be corrected and '(Fr.) Kummer' be replaced by '(Fr.) Staude.'

## BIBLIOGRAPHY

- DONK, M. A. (1949). Nomenclatural notes on generic names of agarics (Fungi: Agaricales). *In* Bull. bot. Gdns Buitenzorg III 18: 271-402.
- KUMMER, P. (1871). Der Führer in die Pilzkunde. Zerbst. 1871.
- ROGERS, D. P. (1949). Nomina conservanda proposita and nomina confusa—Fungi. Supplement. *In* Farlowia 4: 15-43. 1950.
- STAUDE, F. (1857). Die Schwämme Mitteldeutschlands, insbesondere des Herzogthums Coburg. [Festgabe für die Mitglieder der XIX. Versammlung deutscher Land- und Forstwirthe zu Coburg.] Coburg. xxviii & 155 pp. 10 pls. — Same: Die Schwämme Mitteldeutschlands. Erste Lieferung. Gotha. 1858.
-

## INDEX TO VOLUME II

### INDEX TO GENERA AND SPECIES

New names and the final members of new combinations are in bold face type. Pages where illustrations appear are marked with an asterisk.

- Acarospora confusa* 401.  
*Acratherum miliaceum* 232.  
*Acrospermum* 441, 487, 488, 491, 492; *compressum* 487, 488, 491, 492; *conicum* 487; *sclerotoides* 488.  
*Actaea aspera* 198, 222.  
*Affonsea* 69, 70; *lucyi* 86; *vaillantii* 95.  
*Afzelia* 359; *bijuga* 13.  
*Agaricum* sp. 466.  
*Agaricus* 460, 463, 495-497.  
*Agathis* 45; *alba* 54.  
*Agrostis*, *see* Index on p. 346; *infirmavar. sumatrana* 225; *steenisi* 226.  
*Akára-Patsjóti* 208.  
*Albizzia* 70; *sect. Pleiophaca* 70; *brevipes* 79; *hansemanni* 84; *hausemannii* 84; *lucyi* 86; *minahassae* 46; *mollis* 88; *sessilis* 93; *vaillantii* 95.  
*Alectoria* *quartiniana* 231.  
*Allantula* 443; *diffusa* 443.  
*Andropogon*, *see* Index on p. 346.  
*Anisoptera*, *see* Index on p. 67; 371.  
*Anthina* 473; *subgen. Pterula* 473; *flammea* 473.  
*Anthoceros* 411, 412, 422; *amboinensis* 418, 419\*, 420, 423\*; *argillaceus* 419, 420; *carolianus* 414; *falsinervius* 415, 418, 420, 421\*, 422\*, 423\*; *falsinervius var. lyratus* 420; *fuciformis* 422; *glandulosus* 418; *gracilis* 420; *indonesicus* 420; *laevis* 412, 413\*, 414\*, 417, 423\*; *monandrus* 415; *philippinensis* 415; *polyandrus* 411, 415, 418; *punctatus* 412; *silvaticus* 415; *tjibodensis* 411, 415, 416\*, 417\*, 423\*; *weistii* 419, 420.  
*Anthoxanthum* 227; *angustum* 227; *angustum var. borneense* 227; *luzoniense* 227, 228, 230; *odoratum* 227; *papuanum* 228, 229\*; *sumatranum* 227.  
*Aphelaria* 444, 468.  
*Aporosa?* *minahassae* 49.  
*Araecoryne* 444; *elegans* 444.  
*Archidendron* 69-71, 72, 76, 85, 96; *affine* 69, 73, 74, 75; *aruense* 72, 75, 76, 89, 90; *beguinii* 72, 73, 77; *bellum* 72, 73, 77, 78, 80, 94; *brachycarpum* 73, 78, 94, 95; *brevicalyx* 73, 79, 92; *brevipes* 73, 74, 79, 88, 89, 92; *calliandrum* 69, 72, 80; *calycinum* 72, 81; *chrysocarpum* 74, 81; *dies-Christi* 69, 72, 81, 84; *effeminatum* 72, 82; *fallax* 73, 83, 96; *forbesii* 72, 83; *gawadense* 79; *glabrum* 72, 81, 82, 84; *gogolense* 69, 74, 84, 85, 91; *graciliflorum* 91; *harmsii* 96; *incurvatum* 72, 79, 85; *kubaryanum* 83, 96; *laxiflorum* 75; *ledermannii* 73, 86, 91; *lucyi* 74, 86, 94; *lucyi var. schlechterii* 69, 74, 87; *megaphyllum* 72, 87, 88; *molle* 72, 73, 88, 89; *mollis* 88; *mucronatum* 73, 89; *nervosum* 69, 73, 89, 90; *oblongum* 72, 90; *pachycarpum* 72, 73, 85, 86, 91; *papuanum* 93, 94; *parviflorum* 74, 91, 92; *parviflorum var. longipes* 69, 73, 92; *peekelii* 83, 86; *racemosum* 75, 76; *schlechterii* 69, 87; *sessile* 73, 81, 93; *sogerense* 74, 93, 94; *solomonense* 74, 94; *sp. nov.* 96; *tenuiracemosum* 96; *trifoliolatum* 69, 73, 94; *vaillantii* 71, 72, 95; *warenense* 92.  
*Aristida* 230; *adscensionis* 230; *macroclada* 230; *meraukensis* 230; *personata* 230; *polyclados* 231; *setacea* 230.  
*Arthraxon* 231; *ciliaris subsp. quartinianus* 231; *hispidus f. quartinianus* 231; *quartinianus* 231; *quartinianus var. monostachyus* 231.  
*Arundinella*, *see* Index on p. 346; *fuscata var. celebica* 233; *holcooides var. ciliata* 231; *humilior* 232; *setosa var. culionensis* 235; *setosa var. lasiostoma* 235.

- Arundo semiannularis 262.  
 Aspiromitus 411, 412; argillaceus 419, 420; buitenzorgius 419, 420; falsinervius 420; gracilis 420, 422; gracilis var. lyratus 422; spinisporus 418, 419.  
 Assa 187, 205, 206; exotica 205, 206; indica 187, 205-207; tetragynia 206.  
 Ataxis horsfieldii 278.  
 Atylosporium 439.  
 Bacidia aspera 399; avellanea 400; elegantula 399; modesta 398; sorediosa 400.  
 Baeomyces 386.  
 Baumannella 444.  
 Baumannella 444; togoensis 444.  
 Blastenia olivacea 401.  
 Botriochloa 246; parviflora 246; parviflora var. mutispicula 247.  
 Brachiaria, see Index on p. 346; fusiformis var. pilicoronata 236; lanceata var. timorensis 237; subquadripara var. hirsuta 241; subquadripara var. pubescens 241.  
 Brachypodium, see Index on p. 347; sylvaticum var. involutum 242; sylvaticum var. pubifolium 242.  
 Brandtia holcoides 231.  
 Bromus, see Index on p. 347; insignis var. ceramicus 245; insignis var. kinabaluensis 224\*, 245; insignis var. scopulorum 246.  
 Bruinsmia 10.  
 Buellia dissipata 402.  
 Caesalpinia digyna 69.  
 Calliandra brevipes 393.  
 Calligonum asperum 198, 222, 223.  
 Calocera viscosa 477.  
 Calophyllum akara 208.  
 Calvaria 452.  
 Campium laciniatum 406; minutulum 406.  
 Cantharellus 441.  
 Capillipedium, see Index on p. 347; as-simile var. glaucophyllum 251; parviflorum var. mutispiculum 247.  
 Capitoclavaria 444; capitata 444.  
 Carex 373, 379; sect. Multiflorae 383; sect. Stellulatae 383; alta 382; baccans 377; blepharolepis 374; breviscapa 380; brunnea 377; brunnea var. dolichocarpa 378; buennemejeri 376; buruensis 379; capillacea 381; ceramica 376; continua 376; cruciata 374; cryptostachys 379; dietrichiae 374; doniana 381; filicina var. angustifolia 376; formosensis 373, 379, 380; gajonum 373, 382; gembolensis 376; horsfieldii 374; indica 374; kinabaluensis 377; lacerans 382; lateralis 381; ligata var. formosensis 379; loheri 381; longibracteata 377; mer-rillii 377; multifolia 381; myosurus 377; nodiflora 374; nubigena 382; nubigena var. fallax 383; oedorrhapha 381; oligostachya 373; palawanensis 374; phacota 382; philippinensis 381; pycnothyrsos 373; 374, 375; rafflesiana 376; rafflesiana var. macrothyrsa 376; ramosii 374; rhynchachaenium 379; rugata 381; semiglomerata 376; speciosa 381; subtransversa 381; teinogy-na 373, 378, 379; verticillata 377.  
 Caripia 444.  
 Cassia mimosoides 69.  
 Casuarina 390; sp. 388.  
 Ceiba pentandra 391.  
 Centotheca 252; latifolia 253; longilamina 253; schlechteri 252.  
 Ceratella 444, 445, 446; aculeata 445; aculina 445; acuminata 445, 446; hele-nae 445, 446; macrospora 445; queletii 445, 446.  
 Ceratellopsis 445, 446; aculeata 446; queletii 446.  
 Chaetotyphula 447.  
 Chiodecton sanguineum 438.  
 Chionachne 253; biaurita 253.  
 Chloris 253; clementis 253; cynodontoides 253; tenera 253.  
 Chromocyphella 437.  
 Chrysopogon 254; collinus 254; montanus 254; parviflorus 249; parviflorus var. spicigera 249; tenuiculmis 254.  
 Chrysothrix 436.  
 Cicilia 435, 436, 438; aeruginosa 437; noli-tangere 436.

- Cinnamomum camphora 394.  
 Cladaria 442, 443, 447-452, 454-457, 460, 475, 478-481, 484, 486.  
 Cladsterigma 451; fusispora 451.  
 Clavaria 442, 447, 448, 451-456, 461, 463-467, 470-472, 476, 478, 479, 481, 482, 484, 490-493; *subgen.* Ramariopsis 487, 488; *sect.* Ceratella 445, 446; *sect.* Holocoryne 463, 464; *sect.* Syncoryne 464; *sect.* Typhula 490; *trib.* Botryoideae 481, 484; *trib.* Corynoideae 481; *trib.* Holocoryne 454; *trib.* Ramaria 481-484, 486; *trib.* Ramaria *B.* Ochrospora 454; *trib.* Ramaria *sect.* Velutinae 460; *trib.* Ramariae 481, 482, 484, 486; acroporphyreia 478; aculina 445; acuta 464; alba, pistilliforme 451; amathystea 448, 454; amethystina 447-449, 454; angulisporea 488; apiculata 453, 454; aurea 454; botrytis 442, 443, 447, 448, 450-453, 455, 456, 475-478, 480, 481, 483-486; broomei 468; byssiseta 477; capitata 444; cinerea 456; coralloides 477, 485; cornea 448; corniculata 442, 443, 447-451, 455, 456, 459, 475-477, 483, 484; cristata 454, 478, 483, 485; cyanocephala 457; dendroides 444; erythropus 490; falcata 441, 463, 464, 490; fistulosa 453, 460; flava 453, 455; formosa 456; fossicola 435; fragilis 451, 452, 464; fragilis *var.* cylindrica 452; furcellata 443, 460-462; grisea 490; gyrans 490, 491; herbarum 487, 488, 491, 492; ignea 465; kunzei 486; ligula 453; militans, crocea 451; militaris 452; miniata 455; mucida 435, 445; muscicola 490, 491; muscoides 456; nigrescens 468; ophioglossoides 452; ophioglossoides, nigra 451; ornithopodioides 473; ovata 490, 491; palmata 448; paradoxa 453; penicillata 443, 456, 467, 472-474; phacorrhiza 490, 491; pistillaris 452, 453; plumosa 472-474; pomacea 447; pratensis 441, 443, 447, 449, 475, 476, 478, 483, 484; pyxidata ("pixidata") 456; rugosa 454, 464, 476, 482, 483, 485; sclerotioides 468, 487, 488, 491, 492; setipes 455; surculus 461, 462, 465; tubulosa 460; uncialis 445, 463; vermicularis 451, 452; vernalis 435; xylogena 474; zollingeri 449, 454, 477.  
 Clavariae ramulosae 447, 475, 481, 484; *subdiv.* Botryoideae 447-449, 475, 484; *subdiv.* Ramalinae 447-449, 475, 481, 482, 484.  
 Clavariachaeta 452, 453, 459.  
 Clavariachaete 453.  
 Clavariadelphus 453, 464.  
 Clavariella 443, 453, 464, 478-480, 484; botrytis 478.  
 Clavicorona 454.  
 Clavulina 449, 454, 455, 464, 465, 468, 476, 480, 485, 490; cinerea 490; cristata 477, 490; rugosa 443, 454, 456, 490; sp. 477.  
 Clavulinopsis 441-443, 450, 451, 455, 465, 476, 479, 480; corniculata 456, 476; miniata 455; sulcata 455.  
 Cleyera 139; mertensiana 139, 149.  
 Clitocybe 497.  
 Cnazonaria 455, 471, 488.  
 Coelorrhachis, *see* Index on p. 347; foveolata 256; rottboellioides *var.* hirsuta 255; rottboellioides *var.* intermedia 255.  
 Coenogonium 437.  
 Collybia 495, 497.  
 Colocasia esculenta 99.  
 Colysis 406, 407; ellipticum 406.  
 Coprinus 496.  
 Cora 436, 437, 438, 440; *subgen.* Cora 436; *subgen.* Eucora 436; gyrolophia 438; ligulata 439; montana 437, 438, 440; pavonia 436, 438, 440; sericea 436; textilis 435, 436.  
 Corallium 450, 455, 486.  
 Corallo-Fungus 451, 455, 482; croceus, ornithopodioides 473.  
 Corallofungus 456.  
 Coralloidea 456, 484.  
 Coralloides 456, 465, 482; sp. 465.  
 Cordyceps militaris 451.  
 Corella 437; brasiliensis 437.  
 Cornicularia 450, 456, 457, 460, 463, 482-484.  
 Corticium 474.  
 Corynoides 457.  
 Cotylelobium 371.

- Courtoisia cyperoides 112.  
 Craibiodendron 179; stellatum 179.  
 Craterellus taxophila ("taxophilus") 454.  
 Crepidotus 496.  
 Crinium 457.  
 Crinula 457; caliciiiformis 457.  
 Crypsis 128.  
 Cupressus sp. 392.  
 Curatella 188.  
 Cyperus 107, 120, 126; *subgen.* Cyperus *sect.* Dichostylis 119; *subgen.* Kyllinga 128; *subgen.* Kyllinga *sect.* Kyllinga *subsect.* Pingues 128; *subgen.* Pycreus 105; *sect.* Dichostylis 120; *sect.* Diffusi 114; *sect.* Haspani 114; *sect.* Pycreus 105; alopecuroides 101; alopecuroides *var.* digynus 101; alternifolius 124; altus 101; aristatus 122; babakans 104\*, 105, 129; babakensis 105; benghalensis 105; bulbosus 107; calcilola 122; calcicolus 122; castaneus 117, 119; compactus 126; compactus *var.* pauciflorus 126; corymbosus *var.* scariosus 103; crypsoides 127\*, 128, 130; cuspidatus 117, 119; cyperoides 126; digitatus 99, 101; digitatus *var.* hookeri 99; diphyllus 103; distans 111; elatus 99; elatus *var.* macronux 97, 99; eleusinoides 111; exaltatus 99, 100\*, 101, 129; flavidus 116; haspan 116; haspan *var.* indicus 116; haspan *f.* flavidus 116; hyalinus 107, 122; imbricatus 99, 101; iwasakii 101; lanceus 122, 124; latespicatus 124, 125\*, 126, 130; latespicatus *var.* graciliscens 126; latespicatus *var.* laxus 126; luzulae 112; michelianus 120; michelianus *subsp.* pygmaeus *f.* filifolius 120; multispicatus 114, 115\*, 129; nipponicus 120; nutans 111; nutans *var.* eleusinoides 111; ohwii 97, 98\*, 99, 129; pachycephalus 118\*, 119, 120, 129; pilosus 105; pilosus *var.* babakensis 105; pilosus *f.* babakensis 105; platystylis 112, 113\*, 129; pseudokyllingoides 112; pygmaeus 120; ramosii 109, 110\*, 111, 129; rubroviridis 109, 111; scariosus 120\*, 103, 129; spectabilis 99; sphacelatus 106\*, 107, 129; sphacelatus *var.* tenuior 109; subtrigonus 120; tenerifae 121\*, 122, 129; tenuispica 116; unioloides 122, 123\*, 124, 126, 129; venustus 101; zollingeri 103, 107, 108\*, 109, 129.  
 Cyrtococcum, *see* Index on p. 347; trigonum *var.* celebicum 258.  
 Dacryopsella 457.  
 Dacryopsis 457; typhae 441, 457.  
 Damara alba 54; selanica 50, 53, 54, 56.  
 Dansera 351-353; procera 355.  
 Danthonia, *see* Index on p. 347; macgregorii 262, 263\*; mamberamensis 259\*; pilosa *var.* bonthainica 258.  
 Davilla 188; hirsuta 194.  
 Deflexula 457.  
 Delima, *see* Index on p. 223.  
 Delimopsis 187; hirsuta 187, 193-195.  
 Dendroceros 412.  
 Dendrocladium 441, 453, 457-459; hirsutum 459; peckholtii 453, 459.  
 Dendroglossa 406, 407.  
 Derminus 497.  
 Deyeuxia epileuca 325.  
 Dialium 351-354; *subgen.* Arouna 351-355; *subgen.* Arouna *subsect.* Indumentosa 352-355; *subgen.* Arouna *subsect.* Pirula 352-355; *subgen.* Arouna *subsect.* Recta 352-353; *subgen.* Dansera 351, 354, 355; *subgen.* Dialium 351-355; *subgen.* Uittienia 351, 354, 355; divaricatum 69; hexasepalum 351, 354; modestum 355; procerum 355.  
 Dichanthium 265, 295; erectum 265; ramosum 266; simplex 265.  
 Dichonema 437, 439; aeruginosa 437; sericeum 439.  
 Dichonemia 437.  
 Dictyonema 437-439; excentricum 438; membranaceum 438; sericeum 437, 438.  
 Dimeria, *see* Index on p. 347; ornithopoda *var.* glabra 266; ornithopoda *var.* ramosa 266.  
 Dimorphocystis 459; laevis 459.  
 Diploschistes centrifugus 394.  
 Dipterocarpus 167, 371; gracilis 9; grandiflorus 9; hasseltii 9; parallelus 11; tampoerau 11; tampurau 11; tuberculatus 167.



- Doliocarpus 188.  
 Donkella 450, 459.  
 Drymoglossum 403, 404, 407; abbreviatum 404; carnosum 403, 405, 406; microphyllum 406; nobukoanum 406; obovatum 406; spatulatum 403.  
 Drymotaenium 404.  
 Dryobalanops 371; hallii 11.  
 Duval-Jouvea babakensis 105.  
  
 Ectrosia 267.  
 Ectrosiopsis 267, 268; aruensis 268, 269; carvifolia 267; lasioclada 268; subaristata 268, 269; subtriflora 268.  
 Eichhornia crassipes 99.  
 Eleiastis 187; laevis 187, 205.  
 Engelhardtia selanica 50, 56.  
 Eoagaricus 460, 470.  
 Eocronartium 470.  
 Eragrostis, *see* Index on p. 347; cumingii *var.* kisarensis 272; cumingii *var.* novoguineensis 271; cumingii *var.* rindjaniensis 272.  
 Erianthus 275; *sect.* Ripidium 275; beccarii 275; chrysotrix 275; decus-sylvae 275; fallax 275, 276; longifolius 276; sumatranus 275.  
 Eriochloa 276; annulata 276; polystachya 276; procera 276; provera *var.* involu-crata 276; punctata 276; ramosa 276; ramosa *var.* involucrata 276; sundaica 276.  
 Eriocladium 461.  
 Eriocladus 442, 443, 460-463, 465, 489; brasiliensis 443, 460-462, 465.  
 Erioderma 388.  
 Erythrina lithosperma 391.  
 Eucalyptus 302.  
 Eugenia malaccensis 206.  
 Eulalia, *see* Index on p. 348.  
 Eulaliopsis binata 278.  
 Euryandra 187; scandens 187, 212, 213.  
  
 Festuca 325; papuana 328; pauciflora 242; pusilla *monstr.* uncinata 326; re-motiflorus 242.  
 Fimbristylis 122.  
 Fructus indicus sarmentosus 193, 196.  
 Fucus caespitosus 452; clavatus 452.  
  
 Funis urens aspera 190; Funis urens glabra 205.  
  
 Ganua ligulata 363.  
 Gelidium 452; corneum 452; pusillum 452.  
 Geoglossum sp. 451.  
 Gliocoryne 463.  
 Gleditschia 69.  
 Gomphidius 496.  
 Goniophlebium 407.  
 Gordonia, *see* Index on p. 182.  
 Graphis psidii 393.  
 Gyalecta misera 395; recedens 395.  
 Gymnoderma 386, 388; coccocarpum 385, 387\*.  
 Gymnopoda 386.  
 Gymnosporangium 465.  
 Gyrolophium 437, 438; elegans 438; mauritianum 438.  
  
 Hansemannia 71; aruensis 75; arvensis 75; brevipes 79; gawadensis 79; glabra 84; gogolense 69; gogolensis 84; mollis 88; oblonga 90; pachycarpa 91.  
 Harmandia kunstleri 360.  
 Hausemannia 71; brevipes 79; glabra 84; mollis 88.  
 Heppia pulchra 396; scabra 396.  
 Hericium 463; hystrix 466.  
 Heringia 463.  
 Hernandia kunstleri 360.  
 Herpotherallon 435, 438, 439.  
 Hierochloë 277, 278; angusta 227; horsfieldii 227, 278.  
 Hirsutella 463, 466; entomophila 463.  
 Holcus ciliatus 231; nitidus 343, 344; parviflorus 246, 247, 249.  
 Holocoryne 443, 454-456, 463, 464, 482, 483, 490.  
 Holwaya 457.  
 Hopea, *see* Index on p. 68; dolosa 18, 19\*; gregaria 21, 22\*; iriana 28, 29\*; nabirensis 27, 28\*; nodosa 25\*; parvifolia 37; similis 30.  
 Hoppea 50.  
 Hormomitaria 464; sulphurea 464.  
 Hydnum echinus 466; hystrix 463.

- Hymenolepis 404, 406.  
 Hypochnus 435, 438, 439; nigrocinctus 438, rubrocinctus 438, 439.  
 Hypolyssus 464.
- Ionaspis badia* 394.
- Isachne*, see Index on p. 348; *albens* var. *elatuscula* 280; *albens* var. *magna* 280; *albens* var. *sylvestris* 280; *albomarginata* 279; *albomarginata* var. *hirsuta* 279; *clementis* var. *vulcanica* 282; *globosa* var. *daviumbuense* 282; *langkawiensis* 284; *miliacea* var. *madurensis* 285; *miliacea* var. *ovalifolia* 285; *myosotis* var. *micrantha* 286; *pangerangensis* var. *marginata* 288, 289\*; *pauciflora* var. *depauperata* 288; *surgens* 281.
- Isaria farinosa* 477, 478.
- Ischaemum*, see Index on p. 348; *barbatum* var. *arundinaceum* 295; *barbatum* var. *gibbum* 294; *barbatum* var. *imbricatum* 294; *barbatum* var. *lodiculare* 294; *celebicum* 296, 297\*; *nativitatis* 298.
- Iseilema* 302; *maculatum* 302, 303\*.
- Juncellus* 103; *alopecuroides* 101.
- Kordera* 456.  
*Korosvel* 187.
- Lachnobolus* 465; *brasiliensis* 465.  
*Lachnocladium* 441-443, 458, 459, 461, 462, 464, 465, 489; *sect.* *Coniocladium* 458, 459; *sect.* *Dendrocladium* 457-459; *sect.* *Lachnocladium* 458; *brasiliense* 458, 460-462; *giganteum* 441, 457-459; *gudalupense* 458; *guyanense* 458; *hookeri* 465; *insigne* 458; *pallescens* 488; *rubiginosum* 453; *tubulosum* 458.
- Laplacea* 178.  
*Laudatea* 437-439; *caespitosa* 439.  
*Lecanactis albida* 393.  
*Lecidea pertusarioides* 397.  
*Lemmaphyllum* 403-409; *sect.* *Eulemma-phyllum* 404; *sect.* *Phlebodiopsis* 403, 409; *sect.* *Pseudolepisorus* 404, 408, 409; *sect.* *Pseudovittaria* 404; *accedens* 403, 409; *carnosum* 405-407, 409; *microphyllum* 407-409; *microphyllum* var. *obovatum* 407; *spatulatum* 403, 405, 406.
- Lentaria* 462, 465, 477; *surculus* 461.  
*Lentinus lepideus* 477.  
*Leontoglossum* 187; *scabrum* 187, 193-195.  
*Lepidogrammitis* 403-409; *drymoglossoides* 404.  
*Lepisorus* 404-407.  
*Leprocaulon* 386; *arbuscula* 386.  
*Leptaspis* 302, 305; *angustifolia* 302, 304, 305; *banksii* 304, 305; *cochleata* 305; *tararaensis* 304; 305; *urceolata* 304, 305; *wightiana* 310.  
*Leptochilus* 407, 409; *minutulum* 406.  
*Lumnitzera* 370, 371.
- Madhuca neriifolia* 363.  
*Manina* 465.  
*Mariscus microcephalus* 126; *microcephalus* var. *pauciflorus* 126.  
*Martela* 465, 466.  
*Martella* 465, 466.  
*Masseola* 442, 466, 489.  
*Matruchotia* 466; *varians* 466.  
*Megaceros* 412.  
*Megalospora flavidula* 398.  
*Meoschium imbricatum* 294.  
*Merisma* 442, 460, 466, 467, 482; *cristatum* 466, 467; *foetidum* 466, 467; *tuberosum* 471.  
*Microcera* 467; *clavariella* 451; *coccophila* 467.  
*Microglaena javanica* 391; *marginata* 391.  
*Microstegium*, see Index on p. 348; *hirsutum* 306; *steenii* 306.  
*Milium ramosum* 276.  
*Mimosa* 359; *invisa* 359, 360; *invisa* var. *inermis* 359.  
*Mitrulea inflata* 470.  
*Mnesithea* 307, 308; *subgen.* *Coelorachis* 308; *geminata* 308, 309; *laevis* 307, 308, 309; *laevis* var. *hirta* 308; *mollicoma* 308, 309; *pubescens* 308, 309; *rupincola* 256; *triflora* 308, 309.  
*Mucronella* 467.  
*Mucronia* 467.  
*Mycoblastus griseomarginatus* 397.  
*Myriostachya* 310; *wightiana* 310; *wightiana* var. *longispicula* 310.

- Myuropteris 406, 407; cordata 406.  
 Myxomycidium 467; pendulum 467.
- Narenga 276; fallax 276.  
 Neurodium sinense 404.  
 Noronhia emarginata 400.  
 Notothyas 412.
- Opegrapha aggregata 392.  
 Ophiurus perforatus 308.  
 Oplismenus, see Index on p. 349.  
 Orthopogon burmanni var. lanatus 312;  
   imbecillis 310.  
 Oryza 312; longiglumis 312\*; ridleyi 312\*,  
   313; stenothyrsa 313.  
 Ottochloa 313; grandiflora 313\*.
- Palaquium 363.  
 Panicum, see Index on p. 349; brevifolium  
   var. hirtifolium 315; montanum  
   var. pubescens 316; sarmentosum var.  
   parvispiculatum 318.  
 Parapterulicium 467; subarbusculum 467.  
 Parashorea 371.  
 Paspalidium 318; punctatum 318; punctatum  
   var. longiglume 319.  
 Paspalum, see Index on p. 349; commersonii  
   var. hirsutum 319; commersonii  
   var. turgidum 320; pseudo-orbiculare  
   321.  
 Penicillaria 456, 467, 468, 474.  
 Perona 468.  
 Pertusaria 397.  
 Peziza scleropyxis 487.  
 Phacorrhiza 468, 470, 471, 492; sclerotioi-  
   des 468.  
 Phacoriza 468.  
 Phacorrhiza 468.  
 Phaeoaphelaria 468; australiensis 468.  
 Phaeoceros 411, 412.  
 Phaeoclavulina 468; macrospora 468.  
 Phaeocoryne 468, 469.  
 Phaeographis ramulicola 393.  
 Phaeopterula 458, 469; juruensis 458,  
   459, 469.  
 Phaeotyphula 469.  
 Phaioclavulina 468, 469.  
 Pholiota 497.  
 Phyllopsora 388.
- Phymatodes 407.  
 Physalacria 444, 460, 470; inflata 460,  
   470.  
 Pistillaria 443, 470, 471, 487, 488, 491;  
   aculeata 446; acuminata 445; coccinea  
   471, 488; diaphana 470; micans 443,  
   470, 471, 488; muscicola 470; ovata  
   470; pusilla 470, 471; queletii 444-446;  
   quisquiliaris 443, 470, 471, 491; sclero-  
   tioides 468, 470, 487, 488, 492; todei  
   491.  
 Pistillina 457, 471; hyalina 471; typhae  
   441, 457.  
 Pithecellobium 69, 70, 74, 75, 96; kuba-  
   ryanum 96; sp. 96; splendens 359.  
 Pithecolobium sessile 93; vaillantii 95.  
 Pleopeltis 405; sect. Phlebodiopsis 403,  
   409.  
 Pleuropus 496, 497.  
 Pleurotus 497.  
 Poa, see Index on p. 349; borneensis 322;  
   erectifolia var. scabrifolia 326; lamii  
   326; turfosa 329, 320\*; wisselii 330.  
 Podocarpus 362, 363; palembanica 362.  
 Podostrombium 469, 471.  
 Podostromium 471.  
 Podostrumbium 471.  
 Pogonatherum 333; crinitum 333, 334;  
   panicum 33, 334; rufibarbatum 334;  
   saccharoideum 333; saccharoideum var.  
   monandrum 333.  
 Pollinia, see Index on p. 349.  
 Polyporus 471, 472.  
 Polyosus 472.  
 Polyozus 441, 471.  
 Polypodium sect. Lepisorus 408; sect.  
   Pleopeltis 409; accedens 403, 405, 408,  
   409; damuense 405; drymoglossoides  
   404, 405, 407-409; subrostratum 407-  
   409.  
 Polyporus 463.  
 Pongamia glabra 94.  
 Porina isidiosa 392.  
 Pseudopogonatherum 335; egregia 335;  
   irritans 335.  
 Pseudotyphula 472; ochracea 472.  
 Psidium cattleyanum 393, 398, 399.  
 Psoroma 388.

- Pteridium 472, 474.  
 Pteris piloselloides 403.  
 Pterocymbium 364; beccarii 365; splendens 363, 364\*, 365; stipitatum 365; tinctorium 365.  
 Pterula 441, 443, 446, 456, 458, 462, 467-469, 472-474; *subgen.* Phaeopterula 469; capillaris 473; dendroides 473; fascicularis 457; hirsuta 458, 459, 469; merismatoides 473; multifida 468, 474; penicillata 473; plumosa 472-474; subulata 473, 474; tenax 473.  
 Pterulicium 467, 474.  
 Pterygota 366; alata 365\*, 366; forbesii 366; horsfieldii 365, 366; thwaitesii 366; trinervia 366.  
 Pycreus 103; latespicatus 124.  
 Pyrrosia 404.  
 Quercus 164; sp. 164.  
 Ramaria 441-443, 447, 448, 450-452, 454-457, 459, 462, 463, 465, 475-486, 490; amethystina 448-450, 454, 477; capitata 444; ceratoides 477; coralloides 477, 485; coralloides *var.* apicibus purpureis 484, 486; cristata 477, 485; cyanocephala 457; farinosa 477, 478; fastigiata 477; fimbriata 477, 478; gelatinosa 477; medullaris 477; muscoides 476, 477, 484; nigrescens 468; ornithopodioides 477; palmata 448, 477; zippelii 457.  
 Ramariopsis 486.  
 Rhipidonema 439.  
 Roehlingia 187; suaveolens 187, 208.  
 Rottboellia, *see* Index on p. 350.  
 Russula 496.  
 Saccharum 335; fallax 275; longifolium 276; panicum 333; robustum 335, 336; spontaneum 336.  
 Schellolepis 407.  
 Schima, *see* Index on p. 182; wallichii *subsp.* bancana 150\*, 169, 170\*, 173\*; wallichii *subsp.* brevifolia 150\*, 155\*, 177\*; wallichii *subsp.* crenata 172; wallichii *subsp.* liukiensis 150\*; wallichii *subsp.* mertensiana 149, 150\*; wallichii *subsp.* monticola 150\*, 155\*, 176; wallichii *subsp.* noronhae 151; wallichii *subsp.* oblata 150\*, 163\*, 165, 166\*, 170\*; wallichii *var.* crenata 150\*, 172, 173\*; wallichii *var.* khasiana 150\*, 164; wallichii *var.* noronhae 150\*, 151, 152\*, 155\*, 156\*, 161\*, 166\*; wallichii *var.* pulgarensis 150\*, 175, 173\*; wallichii *var.* superba 150\*, 159, 161\*, 163\*.  
 Schizachyrium, *see* Index on p. 350; fragile *var.* luzoniense 338; fragile *var.* malayanum 337; fragile *var.* sinense 338.  
 Schrebera kusnotoi 360, 361\*; swietenoides 362.  
 Scleroglossum 441, 487, 488, 492; acrospermum 487, 488; bulbosum 488; conicum 487; lanceolatum 487.  
 Scleromitra 471, 488.  
 Sclerotium 492.  
 Scytinopogon 442, 488.  
 Sebacina 467; incrustans 467.  
 Segueria asiatica 193-195, 198.  
 Sehima 179; galpinii 179.  
 Septobasidium 435.  
 Serianthes givalobium 357, 358\*.  
 Setaria, *see* Index on p. 350; laxa *var.* navitatis 343; pallide-fusca *var.* brevifolia 340; pallide-fusca *var.* ictura 340; roemerii 340.  
 Shorea, *see* Index on p. 68; 164, 369; montigena 57, 58\*; robusta 164.  
 Siphula ceratites *var.* simplex 387\*, 389; dichotoma 387\*, 390.  
 Sorghum, *see* Index on p. 350; timorense *var.* villosissimum 345.  
 Sparassis 442, 466, 488; crispa 466, 488.  
 Sparaxis 466, 488.  
 Sphaerula 489; capitata 489.  
 Sphyridium 386.  
 Spodiopogon 275; beccarii 275.  
 Spondias sp. 392.  
 Sporotrichum 439.  
 Staurothele australis 390.  
 Stelligera 462, 489; membranacea 489.  
 Sterculia 365, 366; alata 366; blancoi 366; foetida 365.

- Stichoclavaria 490; falcata 490.  
Stichoramaria 454, 455, 464, 490; cristata 490.  
Styrax 10.  
Swartzia 69.
- Taenitis 403.
- Tetracera, *see* Index on p. 224; asiatica *subsp.* andamanica 195, 197\*; asiatica *subsp.* zeylanica 196, 197\*; nordtiana *var.* celebica 203, 204\*; nordtiana *var.* louisianica 202, 204\*.
- Tetradia 365, 366; horsfieldii 365, 366.
- Thelephora 436, 439, 458, 461, 467; *subdiv.* Himantia 439; *trib.* Merisma 467; con-torta 471, 472; palmata 466, 467, 477; sericea 436; textilis 435, 436.
- Thelocarpon 401; algicola 400.
- Thysanothecium 389; casuarinum 387\*, 388, 389; hookeri 389; hyalinum 389; indicum 389.
- Tigarea aspera 222.
- Trachytella 223; actaea 223; aspera 223; calligonum 223.
- Tragia scandens 190, 191.
- Traxilisa 223; aspera 223.
- Tremellodendron 472.
- Trentepohlia 392-395.
- Tricholoma 497, 498.
- Trisetum 234; involutum 242; latifolium 233, 234.
- Typha 490.
- Typhidium 490, 491.
- Typhula 446, 447, 470-472, 490, 491; can-dida 455; grevillei 455; hyalina 447; phacorhiza 490; phaeosperma 469; sclerotioides 463, 491, 492; todei 491.
- Uittienia 351-353; modesta 355.
- Ulva pavonia 437.
- Unona selanica 50.
- Valbomia 187.
- Vararia 489.
- Vatica, *see* Index on p. 68; 369, 371.
- Viburnum 131, 132; clemensae 131\*, 132.
- Wahlbomia 187; indica 187, 205, 206.
- Wainiocora 437, 438, 440; ciferrii 440.
- Weatherbya 403, 405, 406, 408, 409; acce-dens 406.
- Wistaria 69.
- Xyloglossum 441, 487, 488, 491, 492; herbarum 488; sclerotioides 488.
-