




A JOURNAL ON TAXONOMIC BOTANY,
PLANT SOCIOLOGY AND ECOLOGY



REINWARDTIA

12(4)

REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY,
PLANT SOCIOLOGY AND ECOLOGY

Vol. 12(4): 261 - 337, 31 March 2008

Editors

ELIZABETH A. WIDJAJA, MIEN A. RIFAI, SOEDARSONO RISWAN, JOHANIS P. MOGEA

Correspondence on The Reinwardtia journal and subscriptions should be addressed to
HERBARIUM BOGORIENSE, BIDANG BOTANI, PUSAT PENELITIAN BIOLOGI - LIPI,
BOGOR, INDONESIA

FLORISTICS AND STRUCTURE OF A LOWLAND DIPTEROCARP FOREST AT WANARISET SAMBOJA, EAST KALIMANTAN, INDONESIA

Received November 3, 2007; accepted January 20, 2008.

KUSWATA KARTAWINATA

Herbarium Bogoriense, Research Center for Biology - LIPI, Cibinong, Bogor, Indonesia; UNESCO Jakarta Office, Jakarta, Indonesia; Botany Department, Field Museum, Chicago, Illinois 60605-2496, USA. E-mail: kkjak@indo.net.id (author for correspondence).

PURWANINGSIH, TUKIRIN PARTOMIHARDJO, RAZALI YUSUF, ROCHADI ABDULHADI & SOEDARSONO RISWAN

Herbarium Bogoriense, Research Center for Biology - LIPI, Cibinong, Bogor, Indonesia

ABSTRACT.

KARTAWINATA, K., PURWANINGSIH, PARTOMIHARDJO, T., YUSUF, R., ABDULHADI, R. & RISWAN, S. 2008. Floristics and structure of a lowland dipterocarp forest at Wanariset Samboja, East Kalimantan, Indonesia. *Reinwardtia* 12(4): 301–323. — The results of a floristic inventory of trees with DBH \leq 10 cm in a lowland dipterocarp forest in East Kalimantan show that 553 species of 192 genera in 62 families, represented by 5847 individuals, with the total basal area of 350.01 m² occurred in the plot of 10.5 hectare sampled. The two leading families in terms of number of species were *Myrtaceae* and *Lauraceae* while according to the total sum of importance values for families were *Dipterocarpaceae* and *Euphorbiaceae*. The forest had the second highest species richness in Indonesia. We recorded 25 species of dipterocarps, constituting 4.53 % of total species with basal area of 85.53 m² or 24.44 % of the total basal area in the plot. *Shorea laevis* (a dipterocarp) and *Pholidocarpus majadum*, (a palm) were the most prominent species occurring here and were two of the ten leading species. The species-area curve rose steadily even up to an area of 10.5 hectare, with a very slight indication of levelling off at about five hectares, indicating high heterogeneity of the forest. Three largest trees were *Shorea pauciflora* (DBH = 196.50 cm) *Dipterocarpus cornutus* (DBH = 170.90 cm), and *Alstonia scholaris* (DBH = 170.00 cm). Some species could be identified as fruit trees and medicinal plants.

Key words: Composition, structure, dipterocarp forest, species richness, East Kalimantan

KARTAWINATA, K., PURWANINGSIH, PARTOMIHARDJO, T., YUSUF, R., ABDULHADI, R. & RISWAN, S. 2008. Floristik dan struktur hutan pamah dipterocarpa di Wanariset Samboja, Kalimantan, Indonesia. *Reinwardtia* 12(4): 301–323. — Hasil inventarisasi pohon dengan DBH (diameter setinggi dada) \geq 10 cm menunjukkan bahwa 553 spesies dari 192 marga dalam 62 suku, yang diwakili oleh 5847 batang pohon dengan luas bidang dasar total 350.01 m² terdapat dalam plot cuplikan 10.5 hektare. Berdasarkan jumlah spesies dua suku utama adalah *Myrtaceae* dan *Lauraceae*, sedangkan menurut nilai penting suku adalah *Dipterocarpaceae* dan *Euphorbiaceae*. Hutan di sini mempunyai kekayaan spesies pohon tertinggi kedua di Indonesia. Spesies Dipterocarpaceae tercatat 25 spesies atau 4.53 % dari jumlah total spesies dengan luas bidang dasar 85.53 m² atau 24.44 % dari luas bidang dasar seluruh pohon dalam petak *Shorea laevis* (dipterokarpa) dan *Pholidocarpus majadum* (palem) adalah spesies pohon paling menonjol di sini dan merupakan dua dari sepuluh jenis pohon utama. Kurva spesies-luas menanjak tajam bahkan sampai 10.5 hektare, dan agak mendatar pada luasan lima hektare, yang menunjukkan heterogenitas hutan yang tinggi. Tiga pohon terbesar adalah *Shorea pauciflora* (DBH = 196.50 cm) *Dipterocarpus cornutus* (DBH = 170.90 cm), dan *Alstonia scholaris* (DBH = 170.00 cm). Beberapa jenis dapat diidentifikasi sebagai pohon buah-buahan dan tumbuhan obat

Kata kunci: Komposisi, struktur, hutan dipterokarpa, kekayaan jenis, Kalimantan Timur.

INTRODUCTION

Borneo is widely acknowledged as one of the most important centers of plant diversity in the world as well as the center of distribution and species diversity for a large number of families and genera within the Malesian archipelago (Whitmore, 1986; Soepadmo, 1995). The most widespread forest ecosystem in Borneo is the mixed dipterocarp forest, mainly characterised by a 40-60 m tall canopy dominated by an association of species of the *Dipterocarpaceae* family. This forest shows the greatest number of species of any rain forest ecosystem in Malesia (Whitmore, 1986; Philips *et al.*, 1994). Mixed dipterocarp forests are also one of the most productive in the tropics and have been extensively logged during the last 30 years. Harvesting rates range from 80 to 100 m³ ha⁻¹ whereas in other parts of the tropics they do not exceed 30 to 50 m³ ha⁻¹ (Sist, 2000). In Kalimantan, most of the lowland dipterocarp forests have been heavily logged and now the hill forests of the interior constitute the remaining primary forest and the main source of timber. Current knowledge of the ecology, floristics, structure and species richness of the lowland mixed dipterocarps forests is mainly based on studies carried out in Sabah, Brunei and Sarawak (*e.g.* Burgess, 1961; Ashton, 1964; Nicholson, 1965; Bruenig, 1969, 1970, 1973; Proctor *et al.*, 1983; Baillie *et al.*, 1987; Ashton *et al.*, 1992; Newberry *et al.*, 1992; Davies & Becker, 1996). For Kalimantan, the studies remain few (Kartawinata *et al.*, 1981; Riswan, 1982; Guhardja *et al.*, 2000; Riswan, 1987a & b; Suselo & Riswan, 1987; Partomihardjo *et al.*, 1987; Soekardjo *et al.*, 1990; Setiadi *et al.*, 1996; Soedjito, 1990; Soedjito & Kartawinata, 1995; Sist & Saridan, 1999; Tanuwijaya *et al.*, 1996). A floristic analysis of the lowland dipterocarp forests of Borneo found that on a regional scale, diversity is highest in south-east Borneo and central Sarawak with *Dipterocarpaceae* as the most common family followed by *Euphorbiaceae* and several geographically distinct floristic regions could be detected. (Slik *et al.* 2003). Based on data from Northern Borneo, mainly Sarawak, Brunei and Sabah, it was suggested that forests of western Borneo were significantly richer in dipterocarp species than those of eastern Borneo (Ashton, 1989; Davies & Becker, 1996). The richness of the lowland rainforest of Sarawak and Brunei might be linked to the higher climatic stability of North-western Borneo which experiences less dramatic and severe drought periods attributed to El Niño-

Southern Oscillation (ENSO) events than the eastern part of the island, especially Sabah (Ashton, 1989; Goldammer *et al.* 1996; Walsh; 1996,) and East Kalimantan (Guhardja *et al.* 2000; Leighton & Wirawan 1986;). However, Sist & Saridan (1999) recently reported that the species richness of the mixed dipterocarp forest of Berau in East Kalimantan, in spite of its eastern location and its proximity to Sabah was much higher than that of the forest of Sabah and similar to that recorded in Sarawak. Similarly, the forest at Wanariset Samboja, was reported to be the richest in Borneo and even in the world (Kartawinata *et al.* 1981, Whitmore 1986). These facts clearly show that our knowledge of the floristic richness and variability of the mixed dipterocarp forest of Kalimantan is still limited. During the last two decades, the forest of Kalimantan has been depleted and is disappearing at an alarming rate. The main causes are intensive and uncontrolled logging, and conversion into industrial plantations. Successive fires following land clearing affected million of hectares during El Nino events (Sunderlin and Resosudarmo, 1996; Dennis, 1999). Regeneration after fires in East Kalimantan was mainly through seedbank germination in lowland dipterocarp forest and through repouting in *kerangas* (Riswan 1982; Riswan & Kartawinata 1988a, 1989) and dipterocarp forests. The regeneration developed better in the twice-burnt area than that in the area burnt once and the density of young trees was higher and even exceeded that of primary forest (Eichhorn 2006). Furthermore, in the burnt forest, the survival and sprouting capacity of primary forest trees and seedling establishment of pioneer trees and shrubs suppressed the establishment of non-forest species and post fire vegetation was found to be less resilient than it was presumed (Nieuwstadt 2002; Nieuwstadts *et al.* 2001,). Fires resulted in changes of forest structure and composition, loss of tree species diversity and invasion of pioneer species. In sum, forest recovery was not only affected by burning but also by environmental changes resulting from fire (Simbolon *et al.* 2005)

A better knowledge of basic ecological information, including floristic composition and structure of the forest, are necessary for development of a sustainable forest management scheme. To date, there are still very few detailed descriptions and quantitative assessments of forest floristics and structure from a huge area of the Malesian rain forests (Whitmore & Sidiyasa 1986) and for Indonesia. Kartawinata (2005) reviewed the state of quantitative vegetation studies from 1960's

onwards and recommended a list of future actions on the subject. The *Lembaga Biologi Nasional* (National Biological Institute), now known as *Pusat Penelitian Biologi* (Research Center for Biology) noted such needs and in the mid 1970s initiated and integrated the vegetation analysis project into its overall biological research program (Kartawinata 2005). The present study was a part of this program presenting a basic descriptive account of the structure and floristic composition of a 10.5 ha permanent plot set up in a lowland mixed dipterocarp forest in Wanariset Samboja in East Kalimantan. It was intended for use by various future studies in order to provide a permanent basis for long-term study of forest dynamics and floristic changes. In this paper, analysis of the species inventory data collected in the plots will be limited to the description of the forest in terms of the main structural parameters (basal area and density), species richness, pattern of relative abundance and family composition, integrating also data from a 1.6 ha section of this plot reported earlier by Kartawinata *et al.* (1981) and on forest gaps by Partomihardjo *et al.* (1987).

STUDY SITE AND METHODS

The study area is located within a 500-ha research forest managed by the Wanariset (Field Research Station) of the Forest Research and Development Agency (FORDA) of the Ministry of Forestry of Indonesia at Samboja, District of Kutai Kertanegara, East Kalimantan, Indonesia at $0^{\circ} 59'$ Lat. and $116^{\circ} 57'$ Long, about 38 km north of Balikpapan (Figure 1). The physiography is undulating to flat. The forest is lowland dipterocarp forest on dryland with small patches of seasonally swampy ground. The elevation varies between 3 to about 50 m above sea level. The climate is everwet and belongs to the rainfall type A with the ratio between dry and wet months (Q) of 4.4 (Schmidt and Ferguson, 1951). The mean annual rainfall recorded at the nearest meteorological station (Balikpapan BP (Woods and Bower, 1982)) was 2425 mm for the period of 1927-1980. The mean monthly rainfall ranged from 126 mm in October to 236 mm in March (Figure 1) and the mean annual number of raindays was 145 with the mean number of monthly raindays ranging from 9 to 14. The forest is situated on the Red Yellow Podzolic Soil and occurs on an alluvial plain of upper Miocene sedimentary rocks (Soepraptohardjo, 1972).

A block of 150m x 700 m was set up in 1979 in an undisturbed location within a lowland dipterocarp forest about 50 m to the north of Km 1.6 of the now Semboja-Semoi-Sepaku road. It was constructed by sequentially placing a series of 10 m x 10 m plots, first along the width of the block, thus forming a 150 m x 10 m transect consisting of 15 plots; with the surface of each plot was parallel to the ground. The second, third and 70th transects were laid down adjacent to one another along the 700-m length of the block. Thus 1050 plots were established. The 700-m length of the plot stretched roughly South to North and the 150-m width from East to West. The habitat of each plot with reference to topography, whether it is located on a swampy site, a flat dry land, a slope or a ridge was noted qualitatively. In each plot, all the trees with DBH (Diameter at Breast Height) ≥ 10 cm were mapped, numbered with aluminum tags, identified, and measured for DBH. Important features of the forest were also recorded qualitatively. The gaps within the plot were mapped and a profile diagram of a 15 x 60 m² subplot was made as reported by Partomihardjo (1987). Voucher specimens or fallen leaves (if a leafy twig was not accessible) from each tree was collected for identification at the Herbarium Bogoriense, Bogor. The authority of botanical names of plants in the plot followed Whitmore *et al.* (1989, 1990), Keßler & Sidiyasa (1994) and Keßler *et al.* (2000). The inventory was carried out between 1979 and 1981.

RESULTS AND DISCUSSION

Floristic characteristics and diversity

The results of the inventory of trees with DBH ≥ 10 cm showed that 553 species of 192 genera in 62 families, represented by 5847 individuals, with the total basal area of 350.01 m² (Table 1) occurred in the 10.5 hectare plot. Of 550 species recorded, we were able to name 425 species. Appendix 1 lists all species of trees with DBH ≥ 10 cm by family and reports absolute densities, relative densities, frequencies, relative frequencies, basal areas, relative basal areas and Importance Values (IV). These parameters were calculated following the standard procedure as discussed by Mueller-Dombois & Ellenberg (1974). The Total Species Importance Values for a Family (TSIVF) indicates the family importance value based on the sum of IVs of all species in a family (Kartawinata *et al.*




Figure 1. The location of the study area at Wanariset Samboja, East Kalimantan

2004). It is evident that the forest had a high species richness compared to other areas within the region. No single species was dominant, instead dominance was shared by several common species. The forest was characterized by uneven species composition, certain species were represented by large number of individuals, while the majority of species were represented by only a few individuals and often only by one specimen. (Appendix 1).

The ten richest families in terms of number of species are shown in Table 3. Ranked by number of species present, Myrtaceae, Lauraceae, Euphorbiaceae and Myristicaceae were the top four families and Dipterocarpaceae occupied the 5th ranking. In small plots in Borneo and Sumatra,

Lauraceae constitutes one of the top three most common families (Whitmore & Sidiyasa 1986). In the lowland forests of West Malesia, Euphorbiaceae is by far the richest family (cf Abdulhadi 1991, Davies & Becker 1996, Poore 1968, Riswan 1982, Sist & Saridan 1999, Suselo & Riswan 1987, Whitmore 1986, Whitmore & Sidiyasa 1986). As in the present study, *Annonaceae*, *Burseraceae*, *Dipterocarpaceae*, *Moraceae*, *Myristicaceae* and *Rubiaceae* are also in the top ten families ranked by species richness in Borneo and Sumatra (Kartawinata et al. 1981, Table 1). In terms of the Total Sum of Important Values for Families (TSIVF), the ten most important families show different order. *Dipterocarpaceae* and *Euphor-*

biaceae had the first and second highest values (Table 3). The high value for *Dipterocarpaceae* was attributed to the presence of large trees in the plots (Annex 1). It is interesting to note that although it contains only three species, *Arecaceae* had a high TSIVF value, resulting from the high number of individuals of *Pholidocarpus majadum* and *Borassodendron borneensis*. In order to determine whether the 550 species recorded in the 10.5-ha plot represent the total number of species in the area studied, a species-area curve was constructed (Figure 2). The 1050 subplots of 10x10 m each were examined to determine the number of additional species recorded each time a subplot was

added. A considerable number of additional and it continued to rise steeply even up to 10.5 hectare. There was a very slight indication of leveling off at about five hectares. This implies that a minimum area can not be determined for this forest. This is similar to lowland tropical rain forests elsewhere in Borneo and the Malay Peninsula as reported by various authors (Kartawinata 2006, Kartawinata *et al.* 1981; Sist & Saridan 1999, Riswan 1982; Whitmore 1986; Wyatt-Smith 1966, etc.), but it is less dramatic compared to lowland forests of Sulawesi (Kartawinata 2005; Whitmore & Sidiyasa 1986).

Table 1. Floristic and structural characteristics

| Stand Characteristics | Dipterocarps | Non- Dipterocarps | Total |
|------------------------------|-----------------|-------------------|--------|
| Number of Species | 25 (4.53 %) | 527 (95.47 %) | 552 |
| Number of trees | 575 (9.83) | 5272 (90.17 %) | 5847 |
| Mean Density (Trees/Ha) | 54.76 | 502.10 | 556.86 |
| Basal area (M^2) | 85.53 (24.44 %) | 264.48 (75.56 %) | 350.01 |
| Mean Basal Area/Ha (M^2) | 8.48 | 25.19 | 33.33 |

Table 2. Density and number of species of trees with DBH \geq 10 cm in selected plots of different forest types in Kalimantan, Sumatra and Sulawesi (Extracted from Kartawinata 2006)

| Locality | Alt. (m) | Plot size (ha) | Mean Density (Trees/ha) | Number of Species | Reference |
|-------------------------------|----------|----------------|-------------------------|-------------------|----------------------------------|
| EAST KALIMANTAN | | | | | |
| Wanariset Samboja | < 100 | 10.5 | 557 | 552 | Present Study |
| Malinau 1 | 100 | 2 x 1.0 | 413 | 240 | Yusuf (2003) |
| Malinau 2 | 100 | 4 x 1.0 | 759 | 404 | Samssoedin (2005) |
| Malinau 3 | <100 | 1.0 | 567 | 225 | Kartawinata (unpublished) |
| Berau | <100 | 3 x 4.0 | 521 | 538 | Sist & Saridan (1999) |
| Lempake | <100 | 1.6 | 445 | 209 | Riswan, (1987a) |
| Bukit Bangkirai | 110 | 1.0 | 445 | 141 | Simbolon <i>et al.</i> (2005) |
| NORTH SUMATRA | | | | | |
| Leuser National Park | 450-670 | 1.6 | 538 | 116 | Abdulhadi <i>et al.</i> (1989) |
| Ketambe 1 | 350-450 | 1.6 | 420 | 94 | Abdulhadi (1991) |
| Ketambe 2 | 350-450 | 1.6 | 475 | 127 | Abdulhadi <i>et al.</i> (1991) |
| Ketambe 3 | | | | | |
| Batang Gadis National Park | 660 | 1.0 | 583 | 182 | Kartawinata <i>et al.</i> (2004) |
| Aek Nauli | | | | | |
| RIAU | | | | | |
| Bukit Tigapuluh National Park | 297 | 1.0 | 453 | 216 | Polosakan (2001) |
| Bukit Lawang | | | | | |

Table 3. Ten important families in terms of the Total Sum of Importance Values (TSIVF) and number of species arranged in descending order indicating the rank

| Order | Family | No of species | Order | Family | TSIVF (%) |
|-------|-------------------------|---------------|-------|-------------------------|-----------|
| 1 | <i>Myrtaceae</i> | 59 | 1 | <i>Dipterocarpaceae</i> | 44.272 |
| 2 | <i>Lauraceae</i> | 51 | 2 | <i>Euphorbiaceae</i> | 29.277 |
| 3 | <i>Euphorbiaceae</i> | 45 | 3 | <i>Myrtaceae</i> | 18.592 |
| 4 | <i>Myristicaceae</i> | 33 | 4 | <i>Lauraceae</i> | 17.999 |
| 5 | <i>Dipterocarpaceae</i> | 25 | 5 | <i>Myristicaceae</i> | 17.982 |
| 6 | <i>Annonaceae</i> | 24 | 6 | <i>Arecaceae</i> | 16.608 |
| 7 | <i>Rubiaceae</i> | 22 | 7 | <i>Sapotaceae</i> | 15.336 |
| 8 | <i>Burseraceae</i> | 21 | 8 | <i>Annonaceae</i> | 13.061 |
| 9 | <i>Moraceae</i> | 21 | 9 | <i>Ebenaceae</i> | 12.164 |
| 10 | <i>Fabaceae</i> | 19 | 10 | <i>Burseraceae</i> | 9.316 |

The forest of the present study is the second most species rich in East Kalimantan after that in Malinau, which constitutes the richest forest in Indonesia (Kartawinata 2005). It can be seen from Table 2 that if the number of species in one hectare is extrapolated from the number in 10.5 ha plot, as is shown also in the species-area curve (Figure 2, and see also data of Abdulhadi *et al.* 1981), the total species richness may be comparable to that of Lempake (E. Kalimantan), Gunung Mulu (Sarawak) Belalong and Andulau (Brunei), Bukit Lawang (Riau), Bukit Lagong (Peninsular Malaysia), and higher than that of Aek Nauli and Ketambe (Sumatra) and Toraut (Sulawesi). The species richness was however lower than at Malinau 3 (East Kalimantan), at least partly due to different plot design.

The calculation of species frequencies was based on the species data from the 70 transects, each consisting of 15 subplots with the total area of 1500 m². The majority of species had frequencies of less than 50 % and species with frequencies greater than 50 % are shown in Table 5. Interestingly, among dipterocarps only *Shorea ovalis* and *S. laevis* showed relatively high frequency, suggesting that most dipterocarps species apparently grow in clumps in this forest.

Simple ordination of the plots did not produce a recognizable pattern of grouping the plots that warrants the separation of swampy habitat on flat lands from dry habitat on slopes and ridges. Field observations concurred that the composition of the forest on swampy sites was not much different from that on dryland, although the following species were




Figure 2. Species-area curve in a lowland dipterocarp forest at Wanariset Samboja, East Kalimantan

Table 4. Ten most important species in descending order of Importance Value (IV)

| No. | Species | Family | IV |
|-----|----------------------------------|-------------------------|--------|
| 1 | <i>Shorea laevis</i> | <i>Dipterocarpaceae</i> | 11.941 |
| 2 | <i>Pholidocarpus majadum</i> | <i>Arecaceae</i> | 10.656 |
| 3 | <i>Diospyros borneensis</i> | <i>Ebenaceae</i> | 6.512 |
| 4 | <i>Eusideroxylon zwageri</i> | <i>Lauraceae</i> | 6.139 |
| 5 | <i>Scaphium macropodum</i> | <i>Sterculiaceae</i> | 5.800 |
| 6 | <i>Polyalthia sumatrana</i> | <i>Annonaceae</i> | 5.190 |
| 7 | <i>Ganua motleyana</i> | <i>Sapotaceae</i> | 5.071 |
| 8 | <i>Borassodendron borneensis</i> | <i>Arecaceae</i> | 4.918 |
| 9 | <i>Dipterocarpus cornutus</i> | <i>Dipterocarpaceae</i> | 4.568 |
| 10 | <i>Shorea ovalis</i> | <i>Dipterocarpaceae</i> | 4.155 |

mostly found in or were restricted to the swampy sites: *Actynodaphne procera*, *Cratoxylum laevifolius*, *Dialium hydnocarpoides*, *Diospyros elliptica*, *D. laevifolia*, *D. maingayi*, *D. pentaphylla*, *D. sumatrana*, *Elipanthus beccarii*, *Elmerillia tsiampacca*, *Eugenia accuminatissima*, *E. beccarii*, *E. densiflora*, *E. densinervia*, *E. lineata*, *E. oleosa*, *Ganua motleyana*, *Horsfieldia crassifolia*, *Litsea noronhae*, *Mammea*

obovata, *Urophyllum corymbosum* and *Xanthophyllum adenotum*. Three species of tree palms were prominent in the forest and two of them, *Oncosperma horridum* and *Pholidocarpus majadum*, mainly occurred on swampy habitats, while the other, *Borassodendron borneensis*, was present mainly on dryland. Certain other species which occurred at lower densities were also characteristic of swampy habitats: *Anthocephalus cadamba*, *Knema laurina*, *Neesia altissima* and *Pometia pinnata*. The total number of individuals recorded is indicated in Appendix 1. The area of the swampy sites was difficult to quantify precisely because they were patchy, but they occurred mainly on the northern part of the plot and there was no ambiguity in assigning trees to this habitat type.

Table 5. Ten species with highest frequency

| No. | Species | Family | Frequency (%) |
|-----|----------------------------------|--------|---------------|
| 1 | <i>Diospyros borneensis</i> | Eben | 87.14 |
| 2 | <i>Polyalthia sumatrana</i> | Anno | 87.14 |
| 3 | <i>Eusideroxylon zwageri</i> | Laur | 78.57 |
| 4 | <i>Borassodendron borneensis</i> | Arec | 74.29 |
| 5 | <i>Pholidocarpus majadum</i> | Arec | 67.14 |
| 6 | <i>Madhuca sericea</i> | Sapo | 62.86 |
| 7 | <i>Mallotus leptophyllus</i> | Euph | 61.43 |
| 8 | <i>Shorea laevis</i> | Dipt | 60.00 |
| 9 | <i>Dacryodes rugosa</i> | Burs | 60.00 |
| 10 | <i>Shorea ovalis</i> | Dipt | 57.14 |

Some species could be identified as (potential) fruit trees and medicinal plants, including the followings:

- (1) Fruit trees: Anacardiaceae (*Bouea macrophylla*, *Mangifera caesia*, *M. foetida*, *M. pajang*). Bombacaceae (*Durio acutifolius*, *D. dulcis*, *D. graveolens*, *D. kutejensis*, *D. lanceolatus*, *D. oxleyanus*), Burseraceae (*Canarium dichotomum*, *C. denticulatum*, *D. littorale*, *D. patentinervium*, *D. pilosum*, *D. rugosum*), Clusiaceae (*Garcinia celebica*, *G. littoralis*, *G. nervosa*), Euphorbiaceae (*Baccaurea deflexa*, *B. kunstleri*, *B. racemosa*, *B. rumphii*, *B. sumatrana*), Fabaceae (*Parkia roxburghii*, *P. speciosa*), Meliaceae (*Sandoricum borneensis*) and Sapindaceae (*Nephelium lappaceum*, *Pometia pinnata*), and (2) Medicinal plants: Apocynaceae (*Alstonia angustifolia*, *A. scholaris*, *Dyera costulata*), Thymelaeaceae (*Aquilaria malaccensis*) and Annonaceae (*Cananga odorata*).

Structure

The basal area of each tree recorded in the plot was calculated. Appendix 1 shows these data grouped by family, as well as for individual species, along with other measures. The total basal area of trees recorded in the plot was 350.01 m², resulting in a mean basal area of 33.33 m²/ha (Table 1). Ten species with the highest basal areas are presented in Table 4, in which *Shorea laevis*, *Pholidocarpus majadum* and *Scaphium macropodium* were the most prominent. It was also evident that ten dipterocarp species were prevalent, with a total basal area of 53.86 m² (46.74%). As a whole, the dipterocarps were the largest trees and dominated the forest with 25 species (representing 4.47% of the total species richness) occupying a total basal area of 85.53 m² (24.96%). Most species had basal areas of less than 1.0 m² and only 71 species (12.90%) had basal areas greater than 1.0 m², of which the highest basal areas of 10-31 m² were shared by three species (Figure 3), i.e. *Shorea laevis*, *Pholidocarpus majadum* and *Scaphium macropodium* (Table 6).

Table 6. Ten species with highest BA in a 10.5-ha plot

| No. | Species | Family | BA (M2) |
|-----|-------------------------------|--------|---------|
| 1 | <i>Shorea laevis</i> | Dipt | 30.455 |
| 2 | <i>Pholidocarpus majadum</i> | Arec | 16.534 |
| 3 | <i>Scaphium macropodium</i> | Ster | 16.500 |
| 4 | <i>Anthocephalus cadamba</i> | Rubi | 9.734 |
| 5 | <i>Eusideroxylon zwageri</i> | Laur | 9.379 |
| 6 | <i>Dipterocarpus cornutus</i> | Dipt | 9.315 |
| 7 | <i>Ganua motleyana</i> | Sapo | 9.223 |
| 8 | <i>Shorea parvifolia</i> | Dipt | 7.242 |
| 9 | <i>Shorea ovalis</i> | Dipt | 6.844 |
| 10 | <i>Shorea pauciflora</i> | Dipt | 6.499 |

121.725

Figure 4 shows the diameter class distribution of trees with DBH ≥ 10 cm in the 10.5-ha plot. The data show more or less a typical size class distribution of tropical undisturbed primary forest. This reveals that 79.13% of the total trees had DBH of less than 30 cm and only 20.87% occurred in the diameter class greater than 30 cm. The trees with large DBH were mainly dipterocarps. Three largest tree species were *Shorea pauciflora* (DBH = 196.50 cm), *Dipterocarpus cornutus* (170.90 cm) and *Alstonia scholaris* (170.0 cm). It is interesting to note that a pioneer species, *Anthocephalus cadamba*, developed well in this forest, where trees were distributed in all diameter classes and reached a maximum DBH of 128 cm. In contrast, *Alstonia scholaris*, another pioneer species, demonstrated a disjunct size distribution (Table 7)




Figure 3. Basal area class distribution and number of tree species with DBH ≥ 10 cm in a 10.5-ha plot in a lowland dipterocarp forest at Wanariset Samboja, E. Kalimantan

disjunct size distribution (Table 7) with only two large individuals of DBH 140 cm and 170 cm (Appendix 1). *A. cadamba* is a light-demanding species (Whitmore 1986) and, in this study, had a large number of individuals with big diameters and a low number of small individuals. This pattern contrasts that of shade-demanding species, such as *Dipterocarpus cornutus*, which had a higher number of individuals of small sizes (Figure 5). Other secondary forest species present in a relatively high number of individuals in the plot are shown in Table 7, but none were as large as *A. cadamba* and *Alstonia scholaris*. These species appeared to occur on sites previously occupied by gaps. An earlier study on this plot (Partomihardjo *et al.* 1987) (Figure 6) reported that gaps covered a total area of 17, 399 m² (16.6 % of the canopy) and gap formation was estimated to be 1,187 m² (1.05 % of the canopy opening annually) and the recovery rate was about 16 years. Meanwhile, a man-made gap of 0.5 ha in a lowland dipterocarp forest at Lempake, about 100 km north east of Wanariset Samboja (Riswan 1982; Riswan & Kartawinata 1989) was immediately occupied by secondary forest species after clearing. The primary forest species arrived later and achieved a 50% proportion of all species after 18 months. The primary forest arrivals included dipterocarps, *Hopea crudiformis*, *Shorea parvifolia* and *Shorea leprosula*, which were present also in the present study (Appendix 1) and have been reported to behave like pioneer species (Riswan 1982; Riswan & Kartawinata 1989, 1991). Furthermore, in a 35 year old 0.8-ha gap at Lempake the large trees were dominated by secondary forest species, primarily *Macaranga* spp. (Riswan 1982; Riswan & Kartawinata 1988a).




Figure 4. Diameter class distribution and number of tree species with DBH ≥ 10 cm in a 10.5-ha plot in a lowland dipterocarp forest at Wanariset Samboja, E. Kalimantan.




Figure 5. Number of individuals of a light-demanding (*Anthocephalus cadamba*) and shade-demanding (*Dipterocarpus cornutus*) according to diameter-class in a 10.5-ha plot of a lowland dipterocarp forest at Wanariset Samboja, Kalimantan Timur

Table 7. Number of trees of major secondary forest species according to diameter class in the 10.5-ha plot of lowland dipterocarp forest at Wanariset Samboja, East Kalimantan.

| SPECIES | DIAMETER CLASS | | | | | | | | | | |
|--------------------------------|----------------|---------|---------|---------|---------|---------|---------|---------|---------|-----------|-----------|
| | 10-19.9 | 20-29.9 | 30-39.9 | 40-49.9 | 50-59.9 | 60-69.9 | 70-79.9 | 80-89.9 | 90.99.9 | 100-109.9 | 110-129.9 |
| <i>Anthocephalus cadamba</i> | 8 | 1 | 1 | 3 | 2 | 3 | 3 | 2 | 3 | 2 | |
| <i>Alstonia scholaris</i> | 5 | 3 | 1 | | | | | | | | 2 |
| <i>Alstonia angustifolia</i> | 4 | 2 | 1 | | | | | | | | |
| <i>Artocarpus anisophyllus</i> | 12 | 2 | 2 | 1 | | | | | | | |
| <i>Artocarpus rigidus</i> | 13 | 2 | 1 | | | | | | | | |
| <i>Buchanania sessilifolia</i> | 3 | 12 | 7 | | | | | | | | |
| <i>Dillenia excelsa</i> | 15 | 2 | 3 | 1 | | | | | | | |
| <i>Endospermum diadenum</i> | 2 | 1 | 1 | | | | | | | | |
| <i>Porterandia anisophylla</i> | 8 | 1 | | | | | | | | | |
| <i>Macaranga gigantea</i> | 2 | 3 | 2 | 2 | | | | | | | |
| <i>Macaranga hypoleuca</i> | 4 | 1 | | | | | | | | | |
| <i>Macaranga tanarius</i> | 14 | | 1 | | | | | | | | |
| <i>Parinari oblongifolia</i> | 7 | 1 | 3 | 1 | 1 | 2 | | | | | |
| <i>Schima wallichii</i> | 2 | | 2 | | 1 | | | | | | |




Figure 6. Canopy phases and a profile diagram of a portion of the 10.5-ha plot within a lowland dipterocarp forest at Wanariset Samboja, East Kalimantan. D – *Dysoxylum sp.*, Dl – *Drypetes laevis*, Dr – *Dacryodes rostrata*, Gb – *Gonystylus bancanus*, H – *Horsfieldia sp.*, Kc – *Knema cinerea*, Km – *Koompasia malaccensis*, *Madhuca* sp., Mc – *Microcos crassifolia*, Ml – *Mallotus leptophyllus*, Ms – *Madhuca sericea*, Nk – *Neoscortechinia kingii*, P – *Polyalthia sp.*, Pl – *Polyalthia rumphii*, Sl – *Shorea laevis*, So – *Shorea ovalis*, Sp – *Shorea parvifolia*, Vu – *Vatica umbonata*. (After Partomihardjo *et al.* 1987)

ACKNOWLEDGEMENT

We thank the Forest Research and Development Agency for allowing us to undertake the study at the forest at the Wanariset Samboja. We are also grateful to Mrs. J.J. Afriastini and Mr. Ismail Rahman of the Herbarium Bogoriense for assisting us in field identification and identifying voucher specimens, as well as to others who have in many ways helped us in the field.

REFERENCES

- ABDULHADI, R., 1991. A *Meliaceae* forest in Ketambe, Gunung Leuser National Park, Sumatra with special reference to the status of Dipterocarp species. In Soerianegara, S.S. Tjitrosomo, R.C.Umaly & I. Umboh (Eds.). *Proceedings of the Fourth Round-table Conference on Dipterocarps*. Bogor, Indonesia, 12-15 December 1989. BIOTROP Special Publication No. 41: 307-315,
- ABDULHADI, R., MIRMANO, E. & YUSUF, R. 1989. Struktur dan komposisi petak hutan *Dipterocarpaceae* di Ketambe, Taman Nasional G. Leuser, Aceh. *Ekologi Indonesia* 1: 29-36.
- ABDULHADI, R. , YUSUF, R. & KARTAWINATA, K. 1991. A riverine tropical rain forest in Ketambe, Gunung Leuser National Park, Sumatra, Indonesia. In Soerianegara, S.S. Tjitrosomo, R.C.Umaly & I. Umboh (Eds.). *Proceedings of the Fourth Round-table Conference on Dipterocarps*. Bogor, Indonesia, 12-15 December 1989. BIOTROP Special Publication No. 41: 247-255.
- ASHTON, P.S. 1964. *Ecological studies in the mixed dipterocarp forests of Brunei state*. Clarendon Press, Oxford. 75 pp
- ASHTON, P. S. 1989. Dipterocarp reproductive biology. In Lieth, H. and Werger, M.J.A. (Eds.). *Tropical rain forest ecosystems: biogeographical and ecological studies* : 219-240. Elsevier, Amsterdam.
- ASHTON, P. S. & HALL, P. 1992. Comparison of structure among mix dipterocarp forests of North West Borneo. *Journal of Ecology* 80 : 459-481.
- BAILLIE, I.C. & ASHTON, P.S. 1983. Some soil aspects of the nutrient cycle in mixed dipterocarp forests in Sarawak. In Sutton, S.I., Whitmore, T.C. & Chadwick, A.C. (Eds.), *Tropical Rain Forest: Ecology and Management* .:347-356. Blackwell Scientific Publications, Oxford. .
- BAILLIE, I.C., ASHTON, P.S., COURT, M.N., ANDERSON, J.A.R., FITZPATRICK, E.A. & TINSLEY, J. 1987. Site characteristics and the distribution of tree species in mixed dipterocarp forest on tertiary sediments in central Sarawak, Malaysia. *Journal of Tropical Ecology* 3: 201-220.
- BRUENIG, E.F. 1969. The classification of forests types in Sarawak. *Malayan Forester* 32: 143-179.
- BRUENIG, E.F. 1970. Stand structure, physiognomy and environmental factors in some lowland forests in Sarawak. *Tropical Ecology* 11 (1) : 26-43.
- BRUENIG, E.F. 1973. Species richness and stand diversity in relation to site and succession of forests in Sarawak and Brunei. *Amazoniana* 4 (3): 293-320
- BURGESS, P.F. 1961. The structure and composition of lowland tropical rainforest in North Borneo. *Malaysian Forester* 24: 66-80.
- DAVIES, S.J. & BECKER, P. 1996. Floristic composition and stand structure of mixed dipterocarp and heath forests in Brunei Darussalam. *Journal of Tropical Forest Science* 8 (4): 542-569.
- DENNIS, R. 1999. *A review of fire projects in Indonesia (1982-1998)*. CIFOR Publication, 105 pages.
- EICHHORN, K. (2006). Plant diversity after rain-forest fires in Borneo. *Blumea Suplement* 18: 1-140
- GOLDAMMER, J.G., SEIBERT & SCHINDELE, W. 1996. Fire in dipterocarp forests. In Schulte A. & Schöne D. (Eds.) *Dipterocarp Forest Ecosystems, Towards Sustainable Management* . :155-185. World Scientific Publishing, Singapore.
- GUHARDJA, E., FATAWI, M., SUTISNA, M., MORI, T., OHTA, S. (Eds.), 2000. Rainforest ecosystems of East Kalimantan: El Niño, drought, fire and human impacts. *Ecological Studies* 140:1-330. Springer-Verlag, Tokyo.
- KARTAWINATA, K. 2005. Six decades of natural vegetation studies in Indonesia. In Soemodihardjo, S. & S.D. Sastrapradja (Eds.), *Six Decades of Science and Scientists in Indonesia* . :95-140. Naturindo, Bogor.
- KARTAWINATA, K. ROCHADI, A. & PARTOMIHARDJO, J. 1981. Composition and structure of a lowland dipterocarp forest at Wanariset, East Kalimantan (Indonesia). *Malaysian Forester* 44, 2/3 : 397-406.
- KARTAWINATA, K., SAMSOEDIN, I. HERIYANTO, M. & AFRIASTINI, J.J. 2004. A tree species inventory in a one hectare plot at the Batang Gadis National Park, North Sumatra, Indonesia. *Reinwardtia* 12: 145-157.
- KEßLER, P.J.A. & SIDYASA, K. 1994. *Trees of the Balikpapan-Samarinda area, East Kalimantan, Indonesia: A manual to 280 selected species*. Tropenbos Series 7. The Tropenbos Foundation, Wageningen.
- KEßLER, P.J.A., PELSER, P.B., RIDSDALE, C.E. & SIDYASA, K. 2000. *Secondary forest trees of Kalimantan, Indonesia –A manual to 300 selected species*. Tropenbos-Kalimantan Series 3. MOFFEC-Tropenbos-Kalimantan Project, Wanariset Samboja, Balikpapan.
- LEIGHTON, M. & WIRAWAN, N. 1986. Catastrophic drought and fire in Borneo tropical rain forest associated with the 1982-1983 El Niño Southern Oscillation event. In Prance, G.T. (Ed.). *Tropical Rain Forest and the World Atmosphere* . :75-102 Westview, Bopulder, Colorado..
- MANOKARAN, N. & SWAINE, M.D. 1994. Population dynamics of trees in dipterocarps forests of Malaysia. *Malaysian Forest Record* . 41. Forest

- Research Institute Malaysia, Kepong.
- MUELLER-DOMBOIS, D. & ELLENBERG, H. 1974. *Aims and Methods of Vegetation Ecology*. John Wiley & Sons, New York
- NEWBERY,D (MCC.), CAMPBELL, E.J.F., LEE, Y.F., RIDSDALE, C.E. & STILL, M.J. 1992. Primary lowland dipterocarp forest at Danum Valley, Sabah, Malaysia : structure, relative abundance and family composition. *Philosophical Transactions of the Royal Society B* 335 : 341-356.
- NICHOLSON, D.I. 1965. A study of virgin forest near Sandakan, North Borneo. In *Proceedings of the Symposium on Humid Tropics Vegetation*. :67-87. Kuching, UNESCO, Paris.
- NIEUWSTADT, M. G. L .VAN., D. SHEIL, & K. KARTAWINATA. 2001. The ecological consequences of logging in the burned forest of East Kalimantan, Indonesia. *Conservation Biology* 15:1183-1186
- NIEUWSTADT, M. G L.VAN, 2002. *Trial by fire: Positive development of a tropical dipterocarp forest*. Ph. D. dissertation, Utrecht University, Utrecht
- PARTOMIHARDJO, T., YUSUF, R. SUNARTI, S., PURWANINGSIH, ABDULHADI, R. AND KARTAWINATA , K. 1987. A preliminary notes on gaps in a lowland dipterocarp forest in Wanariset, East Kalimantan. In Kostermanns, A.J.G.H. (Ed): *Proceedings of the Third International Round Table on Dipterocarps*. :241-253. UNESCO, Jakarta, Indonesia.
- PHILIPS, O.L., HALL, P., GENTRY, A.H., SAWYER, S.A. & VASQUEZ, R. 1994. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences* 91: 2805-2809.
- POLOSAKAN, R. 2001. Komposisi jenis pohon di hutan kawasan Taman Nasional Bukit Tigapuluh, Propinsi Riau. In Sunaryo and Six Others (Eds.). Laporan Teknik 2001: 31-40. Proyek Inventarisasi dan Karakterisasi Sumberdaya Hayati, Pusat Penelitian Biologi, LIPI, Bogor.
- POULSEN, A.D., NIELSEN, I.C., TAN, S. & BALSLEV, H. 1996. A quantitative inventory of trees in one hectare of mixed dipterocarp forest in Temburong, Brunei Darussalam. In Edwards, D.S., Booth, W.E., & Choy, S.C. (Eds). *Tropical Rain Forest Research - Current Issues*. Dordrecht, the Netherland's Kluwer Academic Press.(In Press cited by Davies & Becker 1996)
- POORE, M.E.D. 1968. Studies in Malaysian rainforests. I. The forest on triassic sediments in Jengka forest reserve. *Journal of Ecology* 56 : 143-196.
- PROCTOR, J., ANDERSON, J.M., CHAI, P. & WALLACK, H.W. 1983. Ecological studies in four contrasting lowland rain forests in Gunung Mulu national park. I. Forest environment, structure and floristics. *Journal of Ecology* 71 : 237-260.
- RISWAN, S. 1982. *Ecological studies on primary, secondary and experimentally cleared mixed dipterocarp forest and kerangas forest in East Kalimantan, Indonesia*. Ph.D. thesis, University of Aberdeen. Aberdeen.
- RISWAN, S. 1987a. Structure and floristic composition of mixed dipterocarp forest at Lempake, East Kalimantan In Kostermanns, A.J.G.H. (Ed): *Proceedings of The third International Round Table on Dipterocarps*. :435-457. UNESCO, Jakarta, Indonesia.
- RISWAN, S. 1987b. Kerangas forest at Gunung Pasir, Samboja, E. Kalimantan. In Kostermans, A.J.G.H. (Ed). *Proceedings of The third International Round Table on Dipterocarps*. :471-494. UNESCO, Jakarta, Indonesia.
- RISWAN, S. & KARTAWINATA, K. 1988a. Regeneration after disturbance in a *kerangas* (heath) forest in East Kalimantan, Indonesia. In Soemodihardjo, S. (Ed.). Some ecological aspects of tropical forest of east kalimantan: a collection of research reports. *MAB Indonesia Contribution No. 48*: 61-85.
- RISWAN, S. & KARTAWINATA, K. 1988b. A lowland dipterocarp forest 35 years after pepper plantation in East Kalimantan, Indonesia. In Soemodihardjo, S. (Ed.). Some ecological aspects of tropical forest of East Kalimantan: A collection of Research Reports. *MAB Indonesia Contribution No. 48*: 1-39.
- RISWAN, S. & KARTAWINATA, K. 1989. Regeneration after disturbance in a lowland mixed dipterocarp forest in East Kalimantan, Indonesia. *Ekologi Indonesia* 1: 9 28.
- RISWAN, S. & KARTAWINATA, K. 1991. Regeneration after disturbance in a lowland mixed dipterocarp forest in East Kalimantan, Indonesia. In Gomez Pompa, A., Whitmore, T.C. & Hadley, M. (Ed.). *Rain Forest Regeneration and Management. Man and the Biosphere Series Vol. 6*: 295-301. UNESCO, Paris and Parthenon Publishing Group Park Ridge..
- SAMSOEDIN, I. 2005. *Biodiversity and sustainability in the Bulungan Research Forest, East Kalimantan, Indonesia: The response of plant species to logging*. Ph. D. Thesis, University of Stirling. Stirling.
- SCHMIDT, F.H. & J.H. FERGUSON (1951). *Rainfall types based on wet and dry period ratios for Indonesia with Western New Guinea.. Verhandelingen Djawatan Meteorolgi dan Geofisika*, Djakarta 42.
- SETIADI, Y. , PERT, D.R., WEBB, C.O. & LEIGHTON, M. 1996. Abundance and spatial distribution of seedling recruitment around adult treess of five *Shorea* species in the Gunung Palung National Park, West Kalimantan, Indonesia. *Tropical Biodiversity* 3: 169-179.
- SIMBOLON, H., SIREGAR, M., WAKIYAMA, S., SUKIGARA, N., ABE, Y. & SHIMIZU, H. 2005. Impacts of forest fires on tree diversity in tropical rain forest of East Kalimantan, Indonesia. *Phyton* 45: 551-559.

- SIST, P. 2000. Reduced impact logging in the tropics: objectives, principles and impacts. *International Forestry Review* 2: 3-10.
- SIST, P. & SARIDAN, A. 1999. Stand structure and floristic composition of a primary lowland dipterocarp forest in East Kalimantan. *Journal of Tropical Forest Science*, 11: 704-722.
- SLIK, J.F.F. and 16 others, 2003. A floristic analysis of the lowland dipterocarp forests of Borneo. *Journal of Biogeography* 30: 1517-1531.
- SOEDJITO, H. 1990. *Root systems of successional and old growth forest species and its role on nutrient dynamics within a tropical rainforest in Indonesia*. Ph.D. dissertation, Rutgers, University, New Brunswick.
- SOEDJITO, H. & KARTAWINATA, K. 1995. Long-term ecological research in Indonesia : achieving sustainable forest management. In Primack R.B. & Lovejoy, T.E. (Eds) "Ecology, Conservation and Management of South East Asian Rainforests". :129-139. Yale University Press, USA.
- SOEKARDJO, S., HAGIHARA, A., YAMAKURA, T., OGAWA, H. 1990. Floristic composition of a tropical rainforest in Indonesian Borneo. *Bull. Nagoya University For.* 10: 1- 43. (Quoted by Ruslim, Y, Matius, P. & Sutisna, M. 2000. In Guhardja *et al.* (Eds.). Rainforest Ecosystems of East Kalimantan: El Niño, Drought, Fire and Human Impacts. *Ecological Studies* Vol. 49. Springer-Verlag, Tokyo.
- SOEPADMO, E. 1995. Plant diversity of the Malesian tropical rainforest and its phytogeographical and economic significance. In Primack R.B. & Lovejoy, T.E. (Eds). *Ecology, Conservation and Management of South East Asian Rainforests*. :19-40. Yale University Press, USA.
- SOEPRAPTOHARDJO, M. 1972. *Generalized soil map, Indonesia*. Scale 1: 2,500,000. 3rd Ed. Soil Research Institute, Bogor
- SUNDERLIN, W.D., RESOSUDARMO, I.A.P. 1996. *Rate and causes of deforestation In indonesia: towards a resolution of the ambiguities*. CIFOR occasional paper n° 9, 19 pp.
- SUSELO, T.B, & RISWAN, S. 1987. Compositional and structural pattern of lowland mixed dipterocarp forest in the Kutai National park, East Kalimantan. In KOSTERMANS, A.J.G.H. (Ed). *Proceedings of the Third International Round Table on Dipterocarps*. :459-470. UNESCO, Jakarta, Indonesia.
- TANUWIJAYA, S.M., ALIMUDIN, R.H., PERT, D.R.A., WEBB, C.O. & LEIGHTON, M. 1996. Population structure and regeneration in two potentially valuable leguminous rain forest tree species, *Sindora coriacea* and *Dialium platysepalum* in Gunung Palung National Park, Indonesia. *Tropical biodiversity* 3: 157-168.
- WALSH, R.P.D. 1996. Drought frequency changes in Sabah and adjacent parts of northern Borneo since the late nineteenth century and possible implications for tropical rain forest dynamics. *J. Trop. Ecol.* 12: 385-407.
- WHITMORE, T. C. 1986. *Tropical rain forests of the Far East*. English Language Book Society/Oxford University Press, Oxford, 1st edition.
- WHITMORE, T.C. & K. SIDIYASA. 1986. Composition and structure of a lowland rain forest at Toraut, northern Sulawesi. *Kew Bulletin* 41: 747-756.
- WHITMORE, T.C., TANTRA, I.G.M. AND SUTISNA, U. 1989. *Tree flora of Indonesia. Check List for Kalimantan. Part I*. Forest Research and Development Centre, Bogor.
- WHITMORE, T.C., TANTRA, I.G.M. AND SUTISNA, U. 1990. *Tree flora of Indonesia. Check List for Kalimantan, Part II*. Forest Research and Development Centre, Bogor.
- WOODS, T.N. & BOWER, R.P. 1982. *Rainfall records, East Kalimantan* - Catatan curah hujan Kalimantan Timur. Report: Analysis summaries and histograms. Transmigration Area Development Project, Dinas Pekerjaan Umum Propinsi Daerah Tingkat I, Kalimantan Timur. Samarinda.

Appendix 1. Number of Species in a Family (in parenthesis), Number of occurrence, Frequency (%), Number of trees, Basal Area (cm²), Relative Basal Area (%), Relative Frequency (%), Relative Density (%), Importance Value in a 10.5 ha plot of lowland dipterocarp forest at Wanariset Semboja, East Kalimantan, Indonesia. The Family Importance Value is the Total Species Importance Values for a Family (TSIVF).

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|----------|-----------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 1 | Actinidiaceae (2 spp.) | | | | | | | | |
| 1 | <i>Saurauia</i> sp. 1 | 1 | 1.43 | 1 | 132.73 | 0.004 | 0.027 | 0.017 | 0.048 |
| 2 | <i>Saurauia</i> sp. 2 | 1 | 1.43 | 1 | 196.07 | 0.006 | 0.027 | 0.017 | 0.050 |
| | Family Importance Value | | | | | | | | 0.098 |
| 2 | Alangiaceae (1 sp.) | | | | | | | | |
| 3 | <i>Alangium ebenaceum</i> | 20 | 28.57 | 26 | 6,899.50 | 0.201 | 0.538 | 0.446 | 1.186 |
| | Family Importance Value | | | | | | | | 1.186 |
| 3 | Anacardiaceae (15) | | | | | | | | |
| 4 | <i>Bouea oppositifolia</i> | 7 | 10.00 | 8 | 2,910.34 | 0.085 | 0.188 | 0.137 | 0.411 |
| 5 | <i>Buchanania sessifolia</i> | 17 | 24.29 | 23 | 6,344.42 | 0.185 | 0.457 | 0.395 | 1.037 |
| 6 | <i>Campnosperma coriaceum</i> | 2 | 2.86 | 2 | 1,005.31 | 0.029 | 0.054 | 0.034 | 0.117 |
| 7 | <i>Koordersiodendron pinnatum</i> | 1 | 1.43 | 1 | 426.38 | 0.012 | 0.027 | 0.017 | 0.057 |
| 8 | <i>Mangifera caesia</i> | 10 | 14.29 | 10 | 9,984.09 | 0.291 | 0.269 | 0.172 | 0.732 |
| 9 | <i>Mangifera foetida</i> | 9 | 12.86 | 10 | 2,646.81 | 0.077 | 0.242 | 0.172 | 0.491 |
| 10 | <i>Mangifera indica</i> | 1 | 1.43 | 1 | 84.95 | 0.002 | 0.027 | 0.017 | 0.047 |
| 11 | <i>Mangifera macrocarpa</i> | 1 | 1.43 | 1 | 343.07 | 0.010 | 0.027 | 0.017 | 0.054 |
| 12 | <i>Mangifera pajang</i> | 4 | 5.71 | 4 | 4,616.66 | 0.135 | 0.108 | 0.069 | 0.311 |
| 13 | <i>Melanochyla bracteata</i> | 4 | 5.71 | 4 | 716.21 | 0.021 | 0.108 | 0.069 | 0.197 |
| 14 | <i>Melanochyla fulvinervis</i> | 7 | 10.00 | 8 | 1,873.00 | 0.055 | 0.188 | 0.137 | 0.380 |
| 15 | <i>Parishia maingayi</i> | 2 | 2.86 | 2 | 7,853.05 | 0.229 | 0.054 | 0.034 | 0.317 |
| 16 | <i>Semecarpus burburyanus</i> | 1 | 1.43 | 1 | 271.72 | 0.008 | 0.027 | 0.017 | 0.052 |
| 17 | <i>Semecarpus forstenii</i> | 3 | 4.29 | 3 | 1,740.21 | 0.051 | 0.081 | 0.051 | 0.183 |
| 18 | <i>Semecarpus glauca</i> | 1 | 1.43 | 1 | 4,185.39 | 0.122 | 0.027 | 0.017 | 0.166 |
| | Family Importance Value | | | | | | | | 4.552 |
| 4 | Annonaceae (24) | | | | | | | | |
| 19 | <i>Cananga odorata</i> | 5 | 7.14 | 6 | 8,079.85 | 0.236 | 0.134 | 0.103 | 0.473 |
| 20 | <i>Cyathocalyx sumatrana</i> | 1 | 1.43 | 3 | 1,359.32 | 0.040 | 0.027 | 0.051 | 0.118 |
| 21 | <i>Goniothalamus macrophylla</i> | 4 | 5.71 | 4 | 637.27 | 0.019 | 0.108 | 0.069 | 0.195 |
| 22 | <i>Meiogyne virgata</i> | 5 | 7.14 | 5 | 1,111.89 | 0.032 | 0.134 | 0.086 | 0.253 |
| 23 | <i>Mitrophora polypyrena</i> | 1 | 1.43 | 1 | 132.73 | 0.004 | 0.027 | 0.017 | 0.048 |
| 24 | <i>Monocardia marginalis</i> | 9 | 12.86 | 13 | 7,056.34 | 0.206 | 0.242 | 0.223 | 0.671 |
| 25 | <i>Oxymitra</i> sp. | 1 | 1.43 | 1 | 122.72 | 0.004 | 0.027 | 0.017 | 0.048 |
| 26 | <i>Polyalthia lateritia</i> | 2 | 2.86 | 2 | 753.94 | 0.022 | 0.054 | 0.034 | 0.110 |
| 27 | <i>Polyalthia glauca</i> | 1 | 1.43 | 1 | 109.36 | 0.003 | 0.027 | 0.017 | 0.047 |
| 28 | <i>Polyalthia lateriflora</i> | 31 | 44.29 | 78 | 29,839.03 | 0.871 | 0.834 | 1.339 | 3.043 |
| 29 | <i>Polyalthia rumphii</i> | 19 | 27.14 | 26 | 6,745.95 | 0.197 | 0.511 | 0.446 | 1.154 |
| 30 | <i>Polyalthia</i> sp. 1 | 1 | 1.43 | 1 | 426.38 | 0.012 | 0.027 | 0.017 | 0.057 |
| 31 | <i>Polyalthia</i> sp. 2 | 1 | 1.43 | 1 | 124.69 | 0.004 | 0.027 | 0.017 | 0.048 |
| 32 | <i>Polyalthia</i> sp. 3 | 1 | 1.43 | 1 | 116.90 | 0.003 | 0.027 | 0.017 | 0.047 |
| 33 | <i>Polyalthia</i> sp. 4 | 1 | 1.43 | 1 | 692.79 | 0.020 | 0.027 | 0.017 | 0.064 |
| 34 | <i>Polyalthia</i> sp. 5 | 1 | 1.43 | 1 | 829.58 | 0.024 | 0.027 | 0.017 | 0.068 |
| 35 | <i>Polyalthia sumatrana</i> | 61 | 87.14 | 143 | 37,494.99 | 1.094 | 1.641 | 2.455 | 5.189 |
| 36 | <i>Popowia hirta</i> | 6 | 8.57 | 6 | 1,025.05 | 0.030 | 0.161 | 0.103 | 0.294 |
| 37 | <i>Popowia</i> sp. 1 | 1 | 1.43 | 1 | 162.86 | 0.005 | 0.027 | 0.017 | 0.049 |
| 38 | <i>Popowia</i> sp. 2 | 1 | 1.43 | 1 | 95.03 | 0.003 | 0.027 | 0.017 | 0.047 |
| 39 | <i>Popowia tomentosa</i> | 2 | 2.86 | 2 | 276.78 | 0.008 | 0.054 | 0.034 | 0.096 |
| 40 | <i>Xylopia ferruginea</i> | 2 | 2.86 | 2 | 979.93 | 0.029 | 0.054 | 0.034 | 0.117 |
| 41 | <i>Xylopia malayana</i> | 13 | 18.57 | 16 | 5,242.90 | 0.153 | 0.350 | 0.275 | 0.777 |
| 42 | <i>Xylopia</i> sp. 1 | 1 | 1.43 | 1 | 86.59 | 0.003 | 0.027 | 0.017 | 0.047 |
| | Family Importance Value | | | | | | | | 13.061 |
| 5 | Apocynaceae (5) | | | | | | | | |
| 43 | <i>Alstonia angustifolia</i> | 6 | 8.57 | 7 | 2,778.12 | 0.081 | 0.161 | 0.120 | 0.363 |
| 44 | <i>Alstonia scholaris</i> | 11 | 15.71 | 11 | 41,046.72 | 1.198 | 0.296 | 0.189 | 1.683 |
| 45 | <i>Dyera</i> sp. | 1 | 1.43 | 1 | 1,231.63 | 0.036 | 0.027 | 0.017 | 0.080 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|-----------|---|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 46 | <i>Dyera lowii</i> | 1 | 1.43 | 1 | 1,063.62 | 0.031 | 0.027 | 0.017 | 0.075 |
| 47 | <i>Willughbeia firma</i> | 3 | 4.29 | 4 | 3,277.67 | 0.096 | 0.081 | 0.069 | 0.245 |
| | Family Importance Value | | | | | | | | 2.445 |
| 6 | Aquifoliaceae (1) | | | | | | | | |
| 48 | <i>Ilex cymosa</i> | 4 | 5.71 | 4 | 17,262.70 | 0.504 | 0.108 | 0.069 | 0.680 |
| | Family Importance Value | | | | | | | | 0.680 |
| 7 | Arecaceae (3) | | | | | | | | |
| 49 | <i>Borassodendron borneensis</i> | 52 | 74.29 | 120 | 49,993.30 | 1.459 | 1.399 | 2.060 | 4.917 |
| 50 | <i>Oncosperma horridum</i> | 13 | 18.57 | 25 | 8,783.28 | 0.256 | 0.350 | 0.429 | 1.035 |
| 51 | <i>Pholidocarpus majadum</i> | 47 | 67.14 | 266 | 165,336.91 | 4.825 | 1.264 | 4.566 | 10.655 |
| | Family Importance Value | | | | | | | | 16.608 |
| 8 | Bombacaceae (7) | | | | | | | | |
| 52 | <i>Durio acutifolius</i> | 17 | 24.29 | 28 | 13,674.62 | 0.399 | 0.457 | 0.481 | 1.337 |
| 53 | <i>Durio dulcis</i> | 9 | 12.86 | 9 | 27,197.65 | 0.794 | 0.242 | 0.154 | 1.190 |
| 54 | <i>Durio graveolens</i> | 16 | 22.86 | 17 | 7,520.90 | 0.220 | 0.430 | 0.292 | 0.942 |
| 55 | <i>Durio kutejensis</i> | 1 | 1.43 | 1 | 314.16 | 0.009 | 0.027 | 0.017 | 0.053 |
| 56 | <i>Durio lanceolatus</i> | 12 | 17.14 | 16 | 3,490.79 | 0.102 | 0.323 | 0.275 | 0.699 |
| 57 | <i>Durio oxleyanus</i> | 17 | 24.29 | 22 | 11,462.90 | 0.335 | 0.457 | 0.378 | 1.169 |
| 58 | <i>Neesia synandra</i> | 12 | 17.14 | 13 | 12,072.47 | 0.352 | 0.323 | 0.223 | 0.898 |
| | Family Importance Value | | | | | | | | 6.289 |
| 9 | Burseraceae (21) | | | | | | | | |
| 59 | <i>Canarium decumanum</i> | 1 | 1.43 | 1 | 143.14 | 0.004 | 0.027 | 0.017 | 0.048 |
| 60 | <i>Canarium denticulatum</i> | 1 | 1.43 | 1 | 224.32 | 0.007 | 0.027 | 0.017 | 0.051 |
| 61 | <i>Canarium dichotomum</i> | 3 | 4.29 | 4 | 765.21 | 0.022 | 0.081 | 0.069 | 0.172 |
| 62 | <i>Canarium hirsutum</i> | 6 | 8.57 | 6 | 1,248.98 | 0.036 | 0.161 | 0.103 | 0.301 |
| 63 | <i>Canarium littorale</i> | 6 | 8.57 | 6 | 965.46 | 0.028 | 0.161 | 0.103 | 0.293 |
| 64 | <i>Canarium patentinervum</i> | 1 | 1.43 | 1 | 232.35 | 0.007 | 0.027 | 0.017 | 0.051 |
| 65 | <i>Canarium pilosum</i> | 3 | 4.29 | 5 | 1,618.41 | 0.047 | 0.081 | 0.086 | 0.214 |
| 66 | <i>Dacryodes laxa</i> | 1 | 1.43 | 1 | 193.59 | 0.006 | 0.027 | 0.017 | 0.050 |
| 67 | <i>Dacryodes rostrata</i> | 39 | 55.71 | 62 | 21,663.10 | 0.632 | 1.049 | 1.064 | 2.745 |
| 68 | <i>Dacryodes rubiginosa</i> | 1 | 1.43 | 1 | 535.02 | 0.016 | 0.027 | 0.017 | 0.060 |
| 69 | <i>Dacryodes rugosa</i> | 42 | 60.00 | 66 | 17,259.69 | 0.504 | 1.130 | 1.133 | 2.766 |
| 70 | <i>Haplolobus moluccanus</i> | 7 | 10.00 | 7 | 1,833.09 | 0.053 | 0.188 | 0.120 | 0.362 |
| 71 | <i>Santiria griffithii</i> | 14 | 20.00 | 19 | 5,495.28 | 0.160 | 0.377 | 0.326 | 0.863 |
| 72 | <i>Santiria laevigata</i> | 1 | 1.43 | 1 | 3,067.96 | 0.090 | 0.027 | 0.017 | 0.134 |
| 73 | <i>Santiria megaphylla</i> | 6 | 8.57 | 6 | 1,003.10 | 0.029 | 0.161 | 0.103 | 0.294 |
| 74 | <i>Santiria</i> sp. 1 | 1 | 1.43 | 1 | 502.73 | 0.015 | 0.027 | 0.017 | 0.059 |
| 75 | <i>Santiria</i> sp. 2 | 1 | 1.43 | 1 | 730.62 | 0.021 | 0.027 | 0.017 | 0.065 |
| 76 | <i>Santiria</i> sp. 3 | 1 | 1.43 | 1 | 165.13 | 0.005 | 0.027 | 0.017 | 0.049 |
| 77 | <i>Santiria</i> sp. 4 | 1 | 1.43 | 1 | 103.87 | 0.003 | 0.027 | 0.017 | 0.047 |
| 78 | <i>Santiria tomentosa</i> | 7 | 10.00 | 7 | 2,042.89 | 0.060 | 0.188 | 0.120 | 0.368 |
| 79 | <i>Trioma malaccensis</i> | 6 | 8.57 | 7 | 1,540.18 | 0.045 | 0.161 | 0.120 | 0.326 |
| | Family IV | | | | | | | | 9.316 |
| 10 | Celastraceae (3) | | | | | | | | |
| 80 | <i>Bhesa paniculata</i> | 16 | 22.86 | 22 | 6,563.37 | 0.192 | 0.430 | 0.378 | 1.000 |
| 81 | <i>Euonymus javanicus</i> | 2 | 2.86 | 2 | 460.45 | 0.013 | 0.054 | 0.034 | 0.102 |
| 82 | <i>Lophopetalum javanicum</i> | 1 | 1.43 | 1 | 91.61 | 0.003 | 0.027 | 0.017 | 0.047 |
| | Family Importance Value | | | | | | | | 1.148 |
| 11 | Chrysobalanaceae (6) | | | | | | | | |
| 83 | <i>Atuna racemosa</i> ssp. <i>excelsa</i> | 7 | 10.00 | 7 | 7,044.01 | 0.206 | 0.188 | 0.120 | 0.514 |
| 84 | <i>Licania splendens</i> | 19 | 27.14 | 27 | 27,448.19 | 0.801 | 0.511 | 0.463 | 1.776 |
| 85 | <i>Maranthes corymbosa</i> | 2 | 2.86 | 3 | 1,596.52 | 0.047 | 0.054 | 0.051 | 0.152 |
| 86 | <i>Parastemon urophyllus</i> | 4 | 5.71 | 4 | 4,844.15 | 0.141 | 0.108 | 0.069 | 0.318 |
| 87 | <i>Parinari oblongifolia</i> | 13 | 18.57 | 15 | 15,568.71 | 0.454 | 0.350 | 0.257 | 1.061 |
| 88 | <i>Parinari</i> sp. | 1 | 1.43 | 1 | 84.95 | 0.002 | 0.027 | 0.017 | 0.047 |
| | Family Importance Value | | | | | | | | 3.867 |
| 12 | Clusiaceae (19) | | | | | | | | |
| 89 | <i>Calophyllum pulcherrimum</i> | 3 | 4.29 | 3 | 1,347.71 | 0.039 | 0.081 | 0.051 | 0.172 |
| 90 | <i>Calophyllum soulattri</i> | 11 | 15.71 | 11 | 42,461.00 | 1.239 | 0.296 | 0.189 | 1.724 |
| 91 | <i>Garcinia celebica</i> | 8 | 11.43 | 10 | 6,005.72 | 0.175 | 0.215 | 0.172 | 0.562 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|-----------|------------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 18 | Ebenaceae (12) | | | | | | | | |
| 142 | <i>Diospyros borneensis</i> | 61 | 87.14 | 195 | 52,219.17 | 1.524 | 1.641 | 3.347 | 6.512 |
| 143 | <i>Diospyros buxifolia</i> | 2 | 2.86 | 2 | 1,755.56 | 0.051 | 0.054 | 0.034 | 0.139 |
| 144 | <i>Diospyros elliptica</i> | 5 | 7.14 | 8 | 11,884.09 | 0.347 | 0.134 | 0.137 | 0.619 |
| 145 | <i>Diospyros macrocarpa</i> | 12 | 17.14 | 18 | 7,307.95 | 0.213 | 0.323 | 0.309 | 0.845 |
| 146 | <i>Diospyros maingayi</i> | 19 | 27.14 | 48 | 20,940.24 | 0.611 | 0.511 | 0.824 | 1.946 |
| 147 | <i>Diospyros malayana</i> | 1 | 1.43 | 1 | 510.71 | 0.015 | 0.027 | 0.017 | 0.059 |
| 148 | <i>Diospyros oblonga</i> | 15 | 21.43 | 26 | 5,646.10 | 0.165 | 0.403 | 0.446 | 1.015 |
| 149 | <i>Diospyros</i> sp 1 | 1 | 1.43 | 1 | 111.22 | 0.003 | 0.027 | 0.017 | 0.047 |
| 150 | <i>Diospyros</i> sp. 2 | 1 | 1.43 | 1 | 191.18 | 0.006 | 0.027 | 0.017 | 0.050 |
| 151 | <i>Diospyros</i> sp. 3 | 2 | 2.86 | 3 | 287.15 | 0.008 | 0.054 | 0.051 | 0.114 |
| 152 | <i>Diospyros sumatrana</i> | 5 | 7.14 | 6 | 677.98 | 0.020 | 0.134 | 0.103 | 0.257 |
| 153 | <i>Diospyros wallichii</i> | 8 | 11.43 | 12 | 4,804.86 | 0.140 | 0.215 | 0.206 | 0.561 |
| | Family Importance Value | | | | | | | | 12.164 |
| 19 | Elaeocarpaceae (5) | | | | | | | | |
| 154 | <i>Elaeocarpus glaber</i> | 12 | 17.14 | 12 | 6,224.44 | 0.182 | 0.323 | 0.206 | 0.710 |
| 155 | <i>Elaeocarpus obtusus</i> | 2 | 2.86 | 2 | 701.87 | 0.020 | 0.054 | 0.034 | 0.109 |
| 156 | <i>Elaeocarpus oxypyren</i> | 1 | 1.43 | 2 | 545.17 | 0.016 | 0.027 | 0.034 | 0.077 |
| 157 | <i>Elaeocarpus polystachyus</i> | 9 | 12.86 | 11 | 6,417.47 | 0.187 | 0.242 | 0.189 | 0.618 |
| 158 | <i>Sloanea javanica</i> | 1 | 1.43 | 1 | 346.36 | 0.010 | 0.027 | 0.017 | 0.054 |
| | Family Importance Value | | | | | | | | 1.568 |
| 20 | Euphorbiaceae (45) | | | | | | | | |
| 159 | <i>Antidesma neurocarpum</i> | 1 | 1.43 | 1 | 706.86 | 0.021 | 0.030 | 0.017 | 0.068 |
| 160 | <i>Aporusa elmeri</i> | 27 | 38.57 | 29 | 5,017.06 | 0.146 | 0.726 | 0.498 | 1.370 |
| 161 | <i>Aporusa falcifera</i> | 16 | 22.86 | 24 | 5,683.79 | 0.166 | 0.430 | 0.412 | 1.008 |
| 162 | <i>Aporusa lucida</i> | 5 | 7.14 | 6 | 2,450.46 | 0.072 | 0.134 | 0.103 | 0.309 |
| 163 | <i>Aporusa lunata</i> | 2 | 2.86 | 2 | 478.95 | 0.014 | 0.054 | 0.034 | 0.102 |
| 164 | <i>Aporusa maingayi</i> | 4 | 5.71 | 5 | 2,271.45 | 0.066 | 0.108 | 0.086 | 0.260 |
| 165 | <i>Aporusa nitida</i> | 1 | 1.43 | 2 | 184.92 | 0.005 | 0.027 | 0.034 | 0.067 |
| 166 | <i>Aporusa</i> sp. | 8 | 11.43 | 8 | 1,368.64 | 0.040 | 0.215 | 0.137 | 0.392 |
| 167 | <i>Aporusa sphaeridophora</i> | 8 | 11.43 | 8 | 1,368.64 | 0.040 | 0.215 | 0.137 | 0.392 |
| 168 | <i>Baccaurea costulata</i> | 18 | 25.71 | 23 | 5,928.13 | 0.173 | 0.484 | 0.395 | 1.052 |
| 169 | <i>Baccaurea kunstleri</i> | 3 | 4.29 | 3 | 493.87 | 0.014 | 0.081 | 0.051 | 0.147 |
| 170 | <i>Baccaurea macrocarpa</i> | 28 | 40.00 | 42 | 13,486.93 | 0.394 | 0.753 | 0.721 | 1.868 |
| 171 | <i>Baccaurea minor</i> | 4 | 5.71 | 4 | 553.61 | 0.016 | 0.108 | 0.069 | 0.192 |
| 172 | <i>Baccaurea parviflora</i> | 1 | 1.43 | 1 | 113.10 | 0.003 | 0.027 | 0.017 | 0.047 |
| 173 | <i>Baccaurea racemosa</i> | 7 | 10.00 | 7 | 12,883.86 | 0.376 | 0.188 | 0.120 | 0.684 |
| 174 | <i>Baccaurea</i> sp. 1 | 1 | 1.43 | 1 | 615.75 | 0.018 | 0.027 | 0.017 | 0.062 |
| 175 | <i>Baccaurea</i> sp. 2 | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| 176 | <i>Baccaurea</i> sp. 3 | 1 | 1.43 | 1 | 188.69 | 0.006 | 0.027 | 0.017 | 0.050 |
| 177 | <i>Baccaurea</i> sp. 4 | 1 | 1.43 | 1 | 408.28 | 0.012 | 0.027 | 0.017 | 0.056 |
| 178 | <i>Baccaurea</i> sp. 5 | 1 | 1.43 | 3 | 436.76 | 0.013 | 0.027 | 0.051 | 0.091 |
| 179 | <i>Baccaurea sumatrana</i> | 9 | 12.86 | 9 | 1,814.82 | 0.053 | 0.242 | 0.154 | 0.450 |
| 180 | <i>Blumeodendron elatiospermum</i> | 3 | 4.29 | 3 | 488.67 | 0.014 | 0.081 | 0.051 | 0.146 |
| 181 | <i>Blumeodendron tokbrai</i> | 3 | 4.29 | 3 | 1,188.70 | 0.035 | 0.081 | 0.051 | 0.167 |
| 182 | <i>Chaetocarpus castanocarpus</i> | 21 | 30.00 | 25 | 14,811.69 | 0.432 | 0.565 | 0.429 | 1.426 |
| 183 | <i>Cleistanthus maingayi</i> | 1 | 1.43 | 1 | 95.03 | 0.003 | 0.027 | 0.017 | 0.047 |
| 184 | <i>Croton oblongus</i> | 16 | 22.86 | 24 | 4,227.28 | 0.123 | 0.430 | 0.412 | 0.966 |
| 185 | <i>Drypetes crassipes</i> | 3 | 4.29 | 1 | 900.26 | 0.026 | 0.081 | 0.017 | 0.124 |
| 186 | <i>Drypetes laevis</i> | 33 | 47.14 | 78 | 56,120.60 | 1.638 | 0.888 | 1.339 | 3.864 |
| 187 | <i>Drypetes littoralis</i> | 1 | 1.43 | 1 | 95.03 | 0.003 | 0.027 | 0.017 | 0.047 |
| 188 | <i>Drypetes longifolia</i> | 8 | 11.43 | 11 | 3,770.41 | 0.110 | 0.215 | 0.189 | 0.514 |
| 189 | <i>Drypetes minahasae</i> | 22 | 31.43 | 38 | 17,926.66 | 0.523 | 0.592 | 0.652 | 1.767 |
| 190 | <i>Drypetes</i> sp. | 1 | 1.43 | 1 | 201.06 | 0.006 | 0.027 | 0.017 | 0.050 |
| 191 | <i>Endospermum diadenum</i> | 3 | 4.29 | 4 | 1,701.44 | 0.050 | 0.081 | 0.069 | 0.199 |
| 192 | <i>Fahrenheitia pendula</i> | 6 | 8.57 | 6 | 1,317.20 | 0.038 | 0.161 | 0.103 | 0.303 |
| 193 | <i>Glochidion philippicum</i> | 7 | 10.00 | 7 | 1,510.91 | 0.044 | 0.188 | 0.120 | 0.353 |
| 194 | <i>Glochidion rubrum</i> | 4 | 5.71 | 5 | 1,294.68 | 0.038 | 0.108 | 0.086 | 0.231 |
| 195 | <i>Macaranga gigantea</i> | 8 | 11.43 | 9 | 6,069.00 | 0.177 | 0.215 | 0.154 | 0.547 |
| 196 | <i>Macaranga hypoleuca</i> | 5 | 7.14 | 5 | 1,181.77 | 0.034 | 0.134 | 0.086 | 0.255 |
| 197 | <i>Macaranga lowii</i> | 3 | 4.29 | 5 | 669.15 | 0.020 | 0.081 | 0.086 | 0.186 |
| 198 | <i>Macaranga tanarius</i> | 12 | 17.14 | 15 | 2,836.33 | 0.083 | 0.323 | 0.257 | 0.663 |
| 199 | <i>Macaranga winkleri</i> | 2 | 2.86 | 2 | 250.65 | 0.007 | 0.054 | 0.034 | 0.095 |
| 200 | <i>Mallotus penangensis</i> | 38 | 54.29 | 69 | 12,328.46 | 0.360 | 1.022 | 1.184 | 2.566 |
| 201 | <i>Mallotus leptophyllus</i> | 43 | 61.43 | 104 | 15,928.22 | 0.465 | 1.157 | 1.785 | 3.407 |
| 202 | <i>Neoscortechinia kingii</i> | 30 | 42.86 | 39 | 10,504.88 | 0.307 | 0.807 | 0.669 | 1.783 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|--------------------------------|-------------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 203 | <i>Pimeleodendron griffithianum</i> | 23 | 32.86 | 28 | 8,699.46 | 0.254 | 0.619 | 0.481 | 1.353 |
| 204 | <i>Ptychopyxis bacciformis</i> | 1 | 1.43 | 1 | 201.06 | 0.006 | 0.027 | 0.017 | 0.050 |
| Family Importance Value | | | | | | | | | |
| 21 | Fabaceae (19) | | | | | | | | |
| 205 | <i>Archidendron clypearia</i> | 5 | 7.14 | 5 | 1,420.93 | 0.041 | 0.134 | 0.086 | 0.262 |
| 206 | <i>Archidendron microcarpum</i> | 5 | 7.14 | 5 | 6,417.90 | 0.187 | 0.134 | 0.086 | 0.408 |
| 207 | <i>Crudia reticulata</i> | 2 | 2.86 | 2 | 168.27 | 0.005 | 0.054 | 0.034 | 0.093 |
| 208 | <i>Crudia ripicola</i> | 1 | 1.43 | 1 | 962.11 | 0.028 | 0.027 | 0.017 | 0.072 |
| 209 | <i>Dialium hydnocarpoides</i> | 1 | 1.43 | 1 | 547.39 | 0.016 | 0.027 | 0.017 | 0.060 |
| 210 | <i>Dialium indum</i> | 7 | 10.00 | 9 | 2,909.06 | 0.085 | 0.188 | 0.154 | 0.428 |
| 211 | <i>Dialium platysepalum</i> | 5 | 7.14 | 5 | 3,675.24 | 0.107 | 0.134 | 0.086 | 0.328 |
| 212 | <i>Dialium</i> sp. | 1 | 1.43 | 1 | 2,436.69 | 0.071 | 0.027 | 0.017 | 0.115 |
| 213 | <i>Koompassia excelsa</i> | 13 | 18.57 | 15 | 14,019.13 | 0.409 | 0.350 | 0.257 | 1.016 |
| 214 | <i>Koompassia malaccensis</i> | 32 | 45.71 | 76 | 42,828.62 | 1.250 | 0.861 | 1.304 | 3.415 |
| 215 | <i>Milletia sericea</i> | 2 | 2.86 | 2 | 193.22 | 0.006 | 0.054 | 0.034 | 0.094 |
| 216 | <i>Parkia speciosa</i> | 1 | 1.43 | 2 | 4,964.70 | 0.145 | 0.027 | 0.034 | 0.206 |
| 217 | <i>Parkia timoriana</i> . | 5 | 7.14 | 5 | 29,722.22 | 0.867 | 0.134 | 0.086 | 1.088 |
| 218 | <i>Phitecellobium</i> sp. | 1 | 1.43 | 1 | 141.03 | 0.004 | 0.027 | 0.017 | 0.048 |
| 219 | <i>Pithecellobium microcarpum</i> | 3 | 4.29 | 3 | 988.22 | 0.029 | 0.081 | 0.051 | 0.161 |
| 220 | <i>Saraca declinata</i> | 3 | 4.29 | 3 | 431.36 | 0.013 | 0.081 | 0.051 | 0.145 |
| 221 | <i>Sindora leiocarpa</i> | 2 | 2.86 | 2 | 254.63 | 0.007 | 0.054 | 0.034 | 0.096 |
| 222 | <i>Sindora velutina</i> | 12 | 17.14 | 17 | 8,645.59 | 0.252 | 0.323 | 0.292 | 0.867 |
| 223 | <i>Uitteinia modesta</i> | 2 | 2.86 | 2 | 228.97 | 0.007 | 0.054 | 0.034 | 0.095 |
| Family Importance Value | | | | | | | | | |
| 22 | Fagaceae 12) | | | | | | | | |
| 224 | <i>Castanopsis costata</i> | 2 | 2.86 | 2 | 995.29 | 0.029 | 0.054 | 0.034 | 0.117 |
| 225 | <i>Castanopsis javanica</i> | 1 | 1.43 | 1 | 1,116.28 | 0.033 | 0.027 | 0.017 | 0.077 |
| 226 | <i>Castanopsis lucida</i> | 1 | 1.43 | 2 | 732.57 | 0.021 | 0.027 | 0.034 | 0.083 |
| 227 | <i>Castanopsis</i> sp. | 1 | 1.43 | 1 | 452.39 | 0.013 | 0.027 | 0.017 | 0.057 |
| 228 | <i>Lithocarpus blumeanus</i> | 18 | 25.71 | 29 | 15,863.89 | 0.463 | 0.484 | 0.498 | 1.445 |
| 229 | <i>Lithocarpus conocarpus</i> | 5 | 7.14 | 5 | 1,545.08 | 0.045 | 0.134 | 0.086 | 0.265 |
| 230 | <i>Lithocarpus hystrix</i> | 2 | 2.86 | 2 | 2,133.34 | 0.062 | 0.054 | 0.034 | 0.150 |
| 231 | <i>Lithocarpus</i> sp. 1 | 1 | 1.43 | 1 | 186.27 | 0.005 | 0.027 | 0.017 | 0.049 |
| 232 | <i>Lithocarpus</i> sp. 2 | 1 | 1.43 | 1 | 769.45 | 0.022 | 0.027 | 0.017 | 0.067 |
| 233 | <i>Lithocarpus sundaicus</i> | 4 | 5.71 | 4 | 3,680.77 | 0.107 | 0.108 | 0.069 | 0.284 |
| 234 | <i>Quercus argenteata</i> | 8 | 11.43 | 10 | 7,462.50 | 0.218 | 0.215 | 0.172 | 0.605 |
| 235 | <i>Quercus gemmiflora</i> | 2 | 2.86 | 2 | 475.36 | 0.014 | 0.054 | 0.034 | 0.102 |
| Family Importance Value | | | | | | | | | |
| 23 | Flacourtiaceae (1) | | | | | | | | |
| 236 | <i>Hydnocarpus polypetala</i> | 23 | 32.86 | 32 | 5,820.27 | 0.170 | 0.619 | 0.549 | 1.338 |
| Family Importance Value | | | | | | | | | |
| 24 | Hypericaceae (2) | | | | | | | | |
| 237 | <i>Cratoxylon cochinchinense</i> | 1 | 1.43 | 1 | 1,075.21 | 0.031 | 0.027 | 0.017 | 0.075 |
| 238 | <i>Cratoxylon hypericinum</i> | 5 | 7.14 | 5 | 4,156.45 | 0.121 | 0.134 | 0.086 | 0.342 |
| Family Importance Value | | | | | | | | | |
| 26 | Iacacinaceae (1) | | | | | | | | |
| 239 | <i>Stemonurus</i> sp. | 1 | 1.43 | 1 | 229.66 | 0.007 | 0.027 | 0.017 | 0.051 |
| Family Importance Value | | | | | | | | | |
| 27 | Lauraceae (51) | | | | | | | | |
| 240 | <i>Actinodaphne glomerata</i> | 1 | 1.43 | 1 | 283.53 | 0.008 | 0.027 | 0.017 | 0.052 |
| 241 | <i>Actinodaphne procera</i> | 4 | 5.71 | 6 | 1,347.67 | 0.039 | 0.108 | 0.103 | 0.250 |
| 242 | <i>Alseodaphne</i> sp. 1 | 1 | 1.43 | 1 | 138.93 | 0.004 | 0.027 | 0.017 | 0.048 |
| 243 | <i>Alseodaphne</i> sp. 2 | 8 | 11.43 | 9 | 4,093.17 | 0.119 | 0.215 | 0.154 | 0.489 |
| 244 | <i>Alseodaphne ob lanceolata</i> | 9 | 12.86 | 11 | 4,027.09 | 0.118 | 0.242 | 0.189 | 0.548 |
| 245 | <i>Alseodaphne umbelliflora</i> | 9 | 12.86 | 9 | 8,616.86 | 0.251 | 0.242 | 0.154 | 0.648 |
| 246 | <i>Beilschmiedia glabra</i> | 7 | 10.00 | 7 | 1,734.66 | 0.051 | 0.188 | 0.120 | 0.359 |
| 247 | <i>Beilschmiedia maingayi</i> | 20 | 28.57 | 24 | 5,837.79 | 0.170 | 0.538 | 0.412 | 1.120 |
| 248 | <i>Beilschmiedia</i> sp. 1 | 1 | 1.43 | 1 | 349.67 | 0.010 | 0.027 | 0.017 | 0.054 |
| 249 | <i>Beilschmiedia</i> sp. 2 | 1 | 1.43 | 1 | 206.12 | 0.006 | 0.027 | 0.017 | 0.050 |
| 250 | <i>Cryptocarya crassifolia</i> | 1 | 1.43 | 1 | 314.16 | 0.009 | 0.027 | 0.017 | 0.053 |
| 251 | <i>Cryptocarya crassinervia</i> | 3 | 4.29 | 3 | 892.88 | 0.026 | 0.081 | 0.051 | 0.158 |
| 252 | <i>Cryptocarya cumingii</i> | 1 | 1.43 | 1 | 193.59 | 0.006 | 0.027 | 0.017 | 0.050 |
| 253 | <i>Cryptocarya ferrea</i> | 1 | 1.43 | 1 | 219.04 | 0.006 | 0.027 | 0.017 | 0.050 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|-----|----------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 254 | <i>Dehaasia borneensis</i> | 1 | 1.43 | 1 | 444.88 | 0.013 | 0.027 | 0.017 | 0.057 |
| 255 | <i>Dehaasia caesia</i> | 5 | 7.14 | 5 | 3,377.01 | 0.099 | 0.134 | 0.086 | 0.319 |
| 256 | <i>Dehaasia firma</i> | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| 257 | <i>Dehaasia incrassata</i> | 2 | 2.86 | 2 | 255.64 | 0.007 | 0.054 | 0.034 | 0.096 |
| 258 | <i>Dehaasia</i> sp. | 1 | 1.43 | 1 | 404.71 | 0.012 | 0.027 | 0.017 | 0.056 |
| 259 | <i>Endiandra beccariana</i> | 5 | 7.14 | 6 | 1,140.00 | 0.033 | 0.134 | 0.103 | 0.271 |
| 260 | <i>Endiandra rubescens</i> | 18 | 25.71 | 19 | 7,708.87 | 0.225 | 0.484 | 0.326 | 1.035 |
| 261 | <i>Eusideroxylon zwageri</i> | 55 | 78.57 | 112 | 93,791.11 | 2.737 | 1.479 | 1.922 | 6.139 |
| 262 | <i>Litsea accendens</i> | 2 | 2.86 | 2 | 487.39 | 0.014 | 0.054 | 0.034 | 0.102 |
| 263 | <i>Litsea angulata</i> | 1 | 1.43 | 1 | 397.61 | 0.012 | 0.027 | 0.017 | 0.056 |
| 264 | <i>Litsea crassifolia</i> | 1 | 1.43 | 1 | 459.96 | 0.013 | 0.027 | 0.017 | 0.057 |
| 265 | <i>Litsea elliptica</i> | 11 | 15.71 | 13 | 6,147.83 | 0.179 | 0.296 | 0.223 | 0.698 |
| 266 | <i>Litsea ferruginea</i> | 10 | 14.29 | 12 | 8,534.30 | 0.249 | 0.269 | 0.206 | 0.724 |
| 267 | <i>Litsea firma</i> | 2 | 2.86 | 2 | 712.37 | 0.021 | 0.054 | 0.034 | 0.109 |
| 268 | <i>Litsea grandis</i> | 9 | 12.86 | 10 | 3,329.13 | 0.097 | 0.242 | 0.172 | 0.511 |
| 269 | <i>Litsea lancifolia</i> | 2 | 2.86 | 2 | 308.14 | 0.009 | 0.054 | 0.034 | 0.097 |
| 270 | <i>Litsea mappacea</i> | 1 | 1.43 | 1 | 390.57 | 0.011 | 0.027 | 0.017 | 0.055 |
| 271 | <i>Litsea noronhae</i> | 4 | 5.71 | 4 | 1,672.54 | 0.049 | 0.108 | 0.069 | 0.225 |
| 272 | <i>Litsea resinosa</i> | 11 | 15.71 | 16 | 10,845.73 | 0.317 | 0.296 | 0.275 | 0.887 |
| 273 | <i>Litsea robusta</i> | 1 | 1.43 | 1 | 314.16 | 0.009 | 0.027 | 0.017 | 0.053 |
| 274 | <i>Litsea</i> sp. 1 | 1 | 1.43 | 1 | 346.36 | 0.010 | 0.027 | 0.017 | 0.054 |
| 275 | <i>Litsea</i> sp. 2 | 1 | 1.43 | 1 | 637.94 | 0.019 | 0.027 | 0.017 | 0.063 |
| 276 | <i>Litsea</i> sp. 3 | 1 | 1.43 | 1 | 89.92 | 0.003 | 0.027 | 0.017 | 0.047 |
| 277 | <i>Litsea</i> sp. 4 | 1 | 1.43 | 1 | 102.07 | 0.003 | 0.027 | 0.017 | 0.047 |
| 278 | <i>Litsea</i> sp. 5 | 1 | 1.43 | 1 | 118.82 | 0.003 | 0.027 | 0.017 | 0.048 |
| 279 | <i>Litsea</i> sp. 6 | 1 | 1.43 | 1 | 183.85 | 0.005 | 0.027 | 0.017 | 0.049 |
| 280 | <i>Litsea</i> sp. 7 | 1 | 1.43 | 1 | 452.39 | 0.013 | 0.027 | 0.017 | 0.057 |
| 281 | <i>Litsea</i> sp. 8 | 1 | 1.43 | 1 | 471.44 | 0.014 | 0.027 | 0.017 | 0.058 |
| 282 | <i>Litsea</i> sp. 9 | 1 | 1.43 | 1 | 539.13 | 0.016 | 0.027 | 0.017 | 0.060 |
| 283 | <i>Litsea</i> sp. 10 | 1 | 1.43 | 1 | 725.83 | 0.021 | 0.027 | 0.017 | 0.065 |
| 284 | <i>Litsea</i> sp. 11 | 1 | 1.43 | 1 | 1,152.09 | 0.034 | 0.027 | 0.017 | 0.078 |
| 285 | <i>Litsea tomentosa</i> | 1 | 1.43 | 1 | 98.52 | 0.003 | 0.027 | 0.017 | 0.047 |
| 286 | <i>Litsea wallichii</i> | 1 | 1.43 | 1 | 84.95 | 0.002 | 0.027 | 0.017 | 0.047 |
| 287 | <i>Neolitsea cesiaefolia</i> | 4 | 5.71 | 4 | 505.50 | 0.015 | 0.108 | 0.069 | 0.191 |
| 288 | <i>Notaphoebe</i> sp. | 1 | 1.43 | 1 | 237.79 | 0.007 | 0.027 | 0.017 | 0.051 |
| 289 | <i>Notaphoebe umbelliflora</i> | 8 | 11.43 | 10 | 10,307.36 | 0.301 | 0.215 | 0.172 | 0.688 |
| 290 | <i>Phoebe laevis</i> | 4 | 5.71 | 4 | 1,922.09 | 0.056 | 0.108 | 0.069 | 0.232 |
| | Family Importance Value | | | | | | | | 17.408 |
| 28 | Lecythidaceae (5) | | | | | | | | |
| 291 | <i>Barringtonia acutangula</i> | 1 | 1.43 | 1 | 286.52 | 0.008 | 0.027 | 0.017 | 0.052 |
| 292 | <i>Barringtonia lanceolata</i> | 7 | 10.00 | 9 | 4,867.72 | 0.142 | 0.188 | 0.154 | 0.485 |
| 293 | <i>Barringtonia macrostachya</i> | 15 | 21.43 | 19 | 5,246.79 | 0.153 | 0.403 | 0.326 | 0.883 |
| 294 | <i>Barringtonia</i> sp. | 1 | 1.43 | 1 | 95.03 | 0.003 | 0.027 | 0.017 | 0.047 |
| 295 | <i>Planchonia valida</i> | 3 | 4.29 | 3 | 1,047.29 | 0.031 | 0.081 | 0.051 | 0.163 |
| | Family Importance Value | | | | | | | | 1.630 |
| 29 | Loganiaceae (1) | | | | | | | | |
| 296 | <i>Strychnos lucida</i> | 1 | 1.43 | 1 | 188.69 | 0.006 | 0.027 | 0.017 | 0.050 |
| | Family Importance Value | | | | | | | | 0.050 |
| 30 | Lythraceae (1) | | | | | | | | |
| 297 | <i>Lagerstroemia floribunda</i> | 6 | 8.57 | 9 | 6,505.05 | 0.190 | 0.161 | 0.154 | 0.506 |
| | Family Importance Value | | | | | | | | 0.506 |
| 31 | Magnoliaceae (4) | | | | | | | | |
| 298 | <i>Elmerillia tsianampacca</i> | 2 | 2.86 | 3 | 430.96 | 0.013 | 0.054 | 0.051 | 0.118 |
| 299 | <i>Magnolia candollii</i> | 1 | 1.43 | 1 | 122.72 | 0.004 | 0.027 | 0.017 | 0.048 |
| 300 | <i>Magnolia elegans</i> | 2 | 2.86 | 2 | 367.57 | 0.011 | 0.054 | 0.034 | 0.099 |
| 301 | <i>Magnolia</i> sp. | 4 | 5.71 | 4 | 663.21 | 0.019 | 0.108 | 0.069 | 0.196 |
| | Family Importance Value | | | | | | | | 0.460 |
| 32 | Melastomataceae (14) | | | | | | | | |
| 302 | <i>Memecylon beccarianum</i> | 6 | 8.57 | 6 | 2,593.09 | 0.076 | 0.161 | 0.103 | 0.340 |
| 303 | <i>Memecylon costatum</i> | 2 | 2.86 | 2 | 320.27 | 0.009 | 0.054 | 0.034 | 0.097 |
| 304 | <i>Memecylon lilacinum</i> | 7 | 10.00 | 8 | 2,081.88 | 0.061 | 0.188 | 0.137 | 0.386 |
| 305 | <i>Memecylon ovatum</i> | 6 | 8.57 | 8 | 1,263.34 | 0.037 | 0.161 | 0.137 | 0.336 |
| 306 | <i>Pternandra azurea</i> | 3 | 4.29 | 3 | 357.79 | 0.010 | 0.081 | 0.051 | 0.143 |
| 307 | <i>Pternandra caeruleascens</i> | 32 | 45.71 | 60 | 11,172.08 | 0.326 | 0.861 | 1.030 | 2.217 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|--------------------------------|----------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 308 | <i>Pternandra cordata</i> | 26 | 37.14 | 57 | 10,460.75 | 0.305 | 0.699 | 0.978 | 1.983 |
| 309 | <i>Pternandra galeata</i> | 21 | 30.00 | 25 | 4,376.10 | 0.128 | 0.565 | 0.429 | 1.122 |
| 310 | <i>Pternandra latifolia</i> | 2 | 2.86 | 2 | 254.01 | 0.007 | 0.054 | 0.034 | 0.096 |
| 311 | <i>Pternandra rostrata</i> | 14 | 20.00 | 21 | 2,786.40 | 0.081 | 0.377 | 0.360 | 0.818 |
| 312 | <i>Pternandra</i> sp. 1 | 1 | 1.43 | 1 | 130.70 | 0.004 | 0.027 | 0.017 | 0.048 |
| 313 | <i>Pternandra</i> sp. 2 | 1 | 1.43 | 1 | 86.59 | 0.003 | 0.027 | 0.017 | 0.047 |
| 314 | <i>Pternandra</i> sp. 3 | 1 | 1.43 | 1 | 186.27 | 0.005 | 0.027 | 0.017 | 0.049 |
| 315 | <i>Pternandra</i> sp. 4 | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| <i>Family Importance Value</i> | | | | | | | | | |
| 33 | Meliaceae (17) | | | | | | | | |
| 316 | <i>Aglaiia macrocarpa</i> | 6 | 8.57 | 13 | 2,514.29 | 0.073 | 0.161 | 0.223 | 0.458 |
| 317 | <i>Aglaiia odorata</i> | 2 | 2.86 | 2 | 221.24 | 0.006 | 0.054 | 0.034 | 0.095 |
| 318 | <i>Aglaiia silvestris</i> | 26 | 37.14 | 32 | 6,969.66 | 0.203 | 0.699 | 0.549 | 1.452 |
| 319 | <i>Aglaiia</i> sp. | 1 | 1.43 | 1 | 109.36 | 0.003 | 0.027 | 0.017 | 0.047 |
| 320 | <i>Aphananixis polystachya</i> | 12 | 17.14 | 15 | 4,866.39 | 0.142 | 0.323 | 0.257 | 0.722 |
| 321 | <i>Chisocheton cumingianus</i> | 1 | 1.43 | 1 | 143.14 | 0.004 | 0.027 | 0.017 | 0.048 |
| 322 | <i>Chisocheton medusae</i> | 5 | 7.14 | 6 | 2,331.40 | 0.068 | 0.134 | 0.103 | 0.306 |
| 323 | <i>Chisocheton</i> sp. | 5 | 7.14 | 1 | 246.06 | 0.007 | 0.134 | 0.017 | 0.159 |
| 324 | <i>Chisocheton patens</i> | 1 | 1.43 | 1 | 86.59 | 0.003 | 0.027 | 0.017 | 0.047 |
| 325 | <i>Dysoxylum arborescens</i> | 1 | 1.43 | 1 | 881.41 | 0.026 | 0.027 | 0.017 | 0.070 |
| 326 | <i>Dysoxylum densiflorum</i> | 2 | 2.86 | 2 | 364.57 | 0.011 | 0.054 | 0.034 | 0.099 |
| 327 | <i>Dysoxylum excelsum</i> | 16 | 22.86 | 25 | 5,843.87 | 0.171 | 0.430 | 0.429 | 1.030 |
| 328 | <i>Dysoxylum macrocarpum</i> | 4 | 5.71 | 4 | 3,081.34 | 0.090 | 0.108 | 0.069 | 0.266 |
| 329 | <i>Dysoxylum pachyrache</i> | 4 | 5.71 | 4 | 598.12 | 0.017 | 0.108 | 0.069 | 0.194 |
| 330 | <i>Kokoona littoralis</i> | 3 | 4.29 | 3 | 1,641.02 | 0.048 | 0.081 | 0.051 | 0.180 |
| 331 | <i>Lansium domesticum</i> | 21 | 30.00 | 21 | 4,071.20 | 0.119 | 0.565 | 0.360 | 1.044 |
| 332 | <i>Sandoricum borneense</i> | 6 | 8.57 | 7 | 2,706.12 | 0.079 | 0.161 | 0.120 | 0.361 |
| <i>Family Importance Value</i> | | | | | | | | | |
| 34 | Monimiaceae (1) | | | | | | | | |
| 333 | <i>Kibara coriacea</i> | 4 | 5.71 | 4 | 670.89 | 0.020 | 0.108 | 0.069 | 0.196 |
| <i>Family Importance Value</i> | | | | | | | | | |
| 35 | Moraceae (21) | | | | | | | | |
| 334 | <i>Artocarpus altilis</i> | 6 | 8.57 | 6 | 2,939.32 | 0.086 | 0.161 | 0.103 | 0.350 |
| 335 | <i>Artocarpus anisophyllus</i> | 13 | 18.57 | 17 | 6,330.02 | 0.185 | 0.350 | 0.292 | 0.826 |
| 336 | <i>Artocarpus champeden</i> | 1 | 1.43 | 2 | 1,719.72 | 0.050 | 0.027 | 0.034 | 0.111 |
| 337 | <i>Artocarpus dadah</i> | 5 | 7.14 | 14 | 5,363.16 | 0.157 | 0.134 | 0.240 | 0.531 |
| 338 | <i>Artocarpus kemando</i> | 6 | 8.57 | 9 | 3,150.37 | 0.092 | 0.161 | 0.154 | 0.408 |
| 339 | <i>Artocarpus longifolius</i> | 1 | 1.43 | 1 | 235.06 | 0.007 | 0.027 | 0.017 | 0.051 |
| 340 | <i>Artocarpus nitidus</i> | 27 | 38.57 | 38 | 11,924.51 | 0.348 | 0.726 | 0.652 | 1.726 |
| 341 | <i>Artocarpus rigidus</i> | 12 | 17.14 | 16 | 3,729.70 | 0.109 | 0.323 | 0.275 | 0.706 |
| 342 | <i>Artocarpus</i> sp. 1 | 8 | 11.43 | 9 | 7,243.06 | 0.211 | 0.215 | 0.154 | 0.581 |
| 343 | <i>Artocarpus</i> sp.2 | 1 | 1.43 | 1 | 174.37 | 0.005 | 0.027 | 0.017 | 0.049 |
| 344 | <i>Artocarpus tamara</i> | 1 | 1.43 | 1 | 664.10 | 0.019 | 0.027 | 0.017 | 0.063 |
| 345 | <i>Ficus ampelas</i> | 1 | 1.43 | 1 | 130.70 | 0.004 | 0.027 | 0.017 | 0.048 |
| 346 | <i>Ficus aurata</i> | 2 | 2.86 | 2 | 211.76 | 0.006 | 0.054 | 0.034 | 0.094 |
| 347 | <i>Ficus crassiramica</i> | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| 348 | <i>Ficus</i> sp. 1 | 2 | 2.86 | 2 | 295.98 | 0.009 | 0.054 | 0.034 | 0.097 |
| 349 | <i>Ficus</i> sp. 2 | 2 | 2.86 | 2 | 463.40 | 0.014 | 0.054 | 0.034 | 0.102 |
| 350 | <i>Ficus sumatrana</i> | 3 | 4.29 | 3 | 327.63 | 0.010 | 0.081 | 0.051 | 0.142 |
| 351 | <i>Ficus sundaica</i> | 2 | 2.86 | 2 | 335.08 | 0.010 | 0.054 | 0.034 | 0.098 |
| 352 | <i>Ficus variegata</i> | 1 | 1.43 | 1 | 160.61 | 0.005 | 0.027 | 0.017 | 0.049 |
| 353 | <i>Ficus xanthophylla</i> | 4 | 5.71 | 4 | 3,609.92 | 0.105 | 0.108 | 0.069 | 0.282 |
| <i>Family Importance Value</i> | | | | | | | | | |
| 36 | Myristicaceae (33) | | | | | | | | |
| 354 | <i>Gymnanthera contracta</i> | 36 | 51.43 | 69 | 21,616.58 | 0.631 | 0.968 | 1.184 | 2.784 |
| 355 | <i>Gymnanthera</i> sp. 1 | 1 | 1.43 | 1 | 363.05 | 0.011 | 0.027 | 0.017 | 0.055 |
| 356 | <i>Gymnanthera</i> sp. 2 | 1 | 1.43 | 1 | 498.76 | 0.015 | 0.027 | 0.017 | 0.059 |
| 357 | <i>Gymnanthera</i> sp. 3 | 1 | 1.43 | 1 | 257.30 | 0.008 | 0.027 | 0.017 | 0.052 |
| 358 | <i>Horsfieldia bracteosa</i> | 18 | 25.71 | 27 | 10,393.08 | 0.303 | 0.484 | 0.463 | 1.251 |
| 359 | <i>Horsfieldia crassifolia</i> | 1 | 1.43 | 1 | 2,463.01 | 0.072 | 0.027 | 0.017 | 0.116 |
| 360 | <i>Horsfieldia glabra</i> | 20 | 28.57 | 32 | 14,414.24 | 0.421 | 0.538 | 0.549 | 1.508 |
| 361 | <i>Horsfieldia grandis</i> | 10 | 14.29 | 12 | 1,783.58 | 0.052 | 0.269 | 0.206 | 0.527 |
| 362 | <i>Horsfieldia irya</i> | 2 | 2.86 | 4 | 1,863.20 | 0.054 | 0.054 | 0.069 | 0.177 |
| 363 | <i>Horsfieldia punctatifolia</i> | 13 | 18.57 | 25 | 9,126.34 | 0.266 | 0.350 | 0.429 | 1.045 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm ²) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|-----|--------------------------------------|-------------------------------|---------------|------------------------------|-------------------------------|-------------------------|------------------------|----------------------|------------------|
| 364 | <i>Horsfieldia</i> sp. 1 | 1 | 1.43 | 1 | 254.47 | 0.007 | 0.027 | 0.017 | 0.051 |
| 365 | <i>Horsfieldia</i> sp. 2 | 1 | 1.43 | 1 | 206.12 | 0.006 | 0.027 | 0.017 | 0.050 |
| 366 | <i>Horsfieldia</i> sp. 3 | 1 | 1.43 | 1 | 153.94 | 0.004 | 0.027 | 0.017 | 0.049 |
| 367 | <i>Horsfieldia</i> sp. 4 | 1 | 1.43 | 1 | 176.71 | 0.005 | 0.027 | 0.017 | 0.049 |
| 368 | <i>Knema cinerea</i> | 28 | 40.00 | 72 | 21,642.47 | 0.632 | 0.753 | 1.236 | 2.621 |
| 369 | <i>Knema conferta</i> | 24 | 34.29 | 42 | 10,594.10 | 0.309 | 0.646 | 0.721 | 1.676 |
| 370 | <i>Knema latericia</i> | 19 | 27.14 | 24 | 5,754.04 | 0.168 | 0.511 | 0.412 | 1.091 |
| 371 | <i>Knema latifolia</i> | 4 | 5.71 | 6 | 1,248.47 | 0.036 | 0.108 | 0.103 | 0.247 |
| 372 | <i>Knema laurina</i> | 9 | 12.86 | 13 | 2,643.36 | 0.077 | 0.242 | 0.223 | 0.542 |
| 373 | <i>Knema lunduensis</i> | 23 | 32.86 | 29 | 7,643.17 | 0.223 | 0.619 | 0.498 | 1.339 |
| 374 | <i>Knema percociacea</i> | 8 | 11.43 | 13 | 2,975.00 | 0.087 | 0.215 | 0.223 | 0.525 |
| 375 | <i>Knema</i> sp. 1 | 1 | 1.43 | 1 | 283.53 | 0.008 | 0.027 | 0.017 | 0.052 |
| 376 | <i>Knema</i> sp. 2 | 1 | 1.43 | 1 | 467.59 | 0.014 | 0.027 | 0.017 | 0.058 |
| 377 | <i>Knema</i> sp. 3 | 1 | 1.43 | 1 | 514.72 | 0.015 | 0.027 | 0.017 | 0.059 |
| 378 | <i>Knema</i> sp. 4 | 1 | 1.43 | 1 | 376.68 | 0.011 | 0.027 | 0.017 | 0.055 |
| 379 | <i>Knema</i> sp. 5 | 1 | 1.43 | 1 | 452.39 | 0.013 | 0.027 | 0.017 | 0.057 |
| 380 | <i>Knema</i> sp. 6 | 1 | 1.43 | 1 | 174.37 | 0.005 | 0.027 | 0.017 | 0.049 |
| 381 | <i>Myristica gaulterifolia</i> | 19 | 27.14 | 25 | 10,447.70 | 0.305 | 0.511 | 0.429 | 1.245 |
| 382 | <i>Myristica iners</i> | 2 | 2.86 | 2 | 286.07 | 0.008 | 0.054 | 0.034 | 0.096 |
| 383 | <i>Myristica lanceifolia</i> | 1 | 1.43 | 1 | 339.79 | 0.010 | 0.027 | 0.017 | 0.054 |
| 384 | <i>Myristica maxima</i> | 4 | 5.71 | 5 | 1,198.73 | 0.035 | 0.108 | 0.086 | 0.228 |
| 385 | <i>Myristica</i> sp. | 1 | 1.43 | 1 | 176.71 | 0.005 | 0.027 | 0.017 | 0.049 |
| 386 | <i>Myristica villosa</i> | 3 | 4.29 | 4 | 583.19 | 0.017 | 0.081 | 0.069 | 0.166 |
| | Family Importance Value | | | | | | | | 7.572 |
| 37 | Myrsinaceae (1) | | | | | | | | |
| 387 | <i>Ardisia</i> sp. | 1 | 1.43 | 1 | 105.68 | 0.003 | 0.027 | 0.017 | 0.047 |
| | Family Importance Value | | | | | | | | 0.047 |
| 38 | Myrtaceae (59) | | | | | | | | |
| 388 | <i>Cleistocalyx operculata</i> | 1 | 1.43 | 1 | 441.15 | 0.013 | 0.027 | 0.017 | 0.057 |
| 389 | <i>Eugenia acuminatissima</i> | 7 | 10.00 | 12 | 3,825.07 | 0.112 | 0.188 | 0.206 | 0.506 |
| 390 | <i>Eugenia acutangula</i> | 35 | 50.00 | 62 | 22,772.72 | 0.665 | 0.941 | 1.064 | 2.670 |
| 391 | <i>Eugenia albidiiramea</i> | 10 | 14.29 | 15 | 9,549.71 | 0.279 | 0.269 | 0.257 | 0.805 |
| 392 | <i>Eugenia aquea</i> | 3 | 4.29 | 4 | 884.95 | 0.026 | 0.081 | 0.069 | 0.175 |
| 393 | <i>Eugenia beccarii</i> | 2 | 2.86 | 2 | 2,140.21 | 0.062 | 0.054 | 0.034 | 0.151 |
| 394 | <i>Eugenia boerlagei</i> | 6 | 8.57 | 9 | 4,743.91 | 0.138 | 0.161 | 0.154 | 0.454 |
| 395 | <i>Eugenia bruneorhamea</i> | 5 | 7.14 | 6 | 1,066.18 | 0.031 | 0.134 | 0.103 | 0.269 |
| 396 | <i>Eugenia corymbosa</i> | 21 | 30.00 | 38 | 15,267.54 | 0.446 | 0.565 | 0.652 | 1.663 |
| 397 | <i>Eugenia decipiens</i> | 4 | 5.71 | 8 | 2,846.91 | 0.083 | 0.108 | 0.137 | 0.328 |
| 398 | <i>Eugenia densiflora</i> | 1 | 1.43 | 1 | 176.71 | 0.005 | 0.027 | 0.017 | 0.049 |
| 399 | <i>Eugenia densinervia</i> | 2 | 2.86 | 2 | 745.47 | 0.022 | 0.054 | 0.034 | 0.110 |
| 400 | <i>Eugenia excelsa</i> | 1 | 1.43 | 1 | 706.86 | 0.021 | 0.027 | 0.017 | 0.065 |
| 401 | <i>Eugenia fastigiata</i> | 25 | 35.71 | 40 | 17,213.92 | 0.502 | 0.672 | 0.687 | 1.861 |
| 402 | <i>Eugenia jambooloides</i> | 8 | 11.43 | 9 | 4,955.50 | 0.145 | 0.215 | 0.154 | 0.514 |
| 403 | <i>Eugenia lanceolata</i> | 1 | 1.43 | 1 | 191.18 | 0.006 | 0.027 | 0.017 | 0.050 |
| 404 | <i>Eugenia lineata</i> | 7 | 10.00 | 7 | 1,151.49 | 0.034 | 0.188 | 0.120 | 0.342 |
| 405 | <i>Eugenia ochneocarpa</i> | 17 | 24.29 | 25 | 10,150.70 | 0.296 | 0.457 | 0.429 | 1.183 |
| 406 | <i>Eugenia oleosa</i> F. Muell. | 9 | 12.86 | 20 | 11,454.01 | 0.334 | 0.242 | 0.343 | 0.920 |
| 407 | <i>Eugenia opaca</i> K. & V. | 13 | 18.57 | 16 | 5,807.38 | 0.169 | 0.350 | 0.275 | 0.794 |
| 408 | <i>Eugenia operculata</i> | 1 | 1.43 | 1 | 441.15 | 0.013 | 0.027 | 0.017 | 0.057 |
| 409 | <i>Eugenia rostrata</i> Bedd. ex Dut | 2 | 2.86 | 2 | 617.52 | 0.018 | 0.054 | 0.034 | 0.106 |
| 410 | <i>Eugenia</i> sp. 1 | 4 | 5.71 | 4 | 854.31 | 0.025 | 0.108 | 0.069 | 0.201 |
| 411 | <i>Eugenia</i> sp. 2 | 3 | 4.29 | 6 | 2,111.62 | 0.062 | 0.081 | 0.103 | 0.245 |
| 412 | <i>Eugenia</i> sp. 3 | 7 | 10.00 | 10 | 5,842.20 | 0.171 | 0.188 | 0.172 | 0.530 |
| 413 | <i>Eugenia</i> sp. 4 | 1 | 1.43 | 2 | 465.31 | 0.014 | 0.027 | 0.034 | 0.075 |
| 414 | <i>Eugenia</i> sp. 5 | 4 | 5.71 | 1 | 394.28 | 0.012 | 0.108 | 0.017 | 0.136 |
| 415 | <i>Eugenia</i> sp. 6 | 1 | 1.43 | 1 | 83.32 | 0.002 | 0.027 | 0.017 | 0.046 |
| 416 | <i>Eugenia</i> sp. 7 | 1 | 1.43 | 1 | 98.52 | 0.003 | 0.027 | 0.017 | 0.047 |
| 417 | <i>Eugenia</i> sp. 8 | 1 | 1.43 | 1 | 109.36 | 0.003 | 0.027 | 0.017 | 0.047 |
| 418 | <i>Eugenia</i> sp. 9 | 1 | 1.43 | 1 | 124.69 | 0.004 | 0.027 | 0.017 | 0.048 |
| 419 | <i>Eugenia</i> sp. 10 | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| 420 | <i>Eugenia</i> sp. 11 | 1 | 1.43 | 1 | 143.14 | 0.004 | 0.027 | 0.017 | 0.048 |
| 421 | <i>Eugenia</i> sp. 12 | 1 | 1.43 | 1 | 151.75 | 0.004 | 0.027 | 0.017 | 0.048 |
| 422 | <i>Eugenia</i> sp. 13 | 1 | 1.43 | 1 | 206.12 | 0.006 | 0.027 | 0.017 | 0.050 |
| 423 | <i>Eugenia</i> sp. 14 | 1 | 1.43 | 1 | 268.80 | 0.008 | 0.027 | 0.017 | 0.052 |
| 424 | <i>Eugenia</i> sp. 15 | 1 | 1.43 | 1 | 352.99 | 0.010 | 0.027 | 0.017 | 0.054 |
| 425 | <i>Eugenia</i> sp. 16 | 1 | 1.43 | 1 | 397.61 | 0.012 | 0.027 | 0.017 | 0.056 |
| 426 | <i>Eugenia</i> sp. 17 | 1 | 1.43 | 1 | 452.39 | 0.013 | 0.027 | 0.017 | 0.057 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm2) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|-----------|----------------------------------|-------------------------------|---------------|------------------------------|------------------|-------------------------|------------------------|----------------------|------------------|
| 46 | Rosaceae (4) | | | | | | | | |
| 469 | <i>Licania splendens</i> | 27 | 38.57 | 27 | 27,448.19 | 0.801 | 0.726 | 0.463 | 1.991 |
| 470 | <i>Prunus arborea</i> | 10 | 14.29 | 10 | 4,280.74 | 0.125 | 0.269 | 0.172 | 0.566 |
| 471 | <i>Prunus beccarii</i> | 13 | 18.57 | 15 | 3,070.82 | 0.090 | 0.350 | 0.257 | 0.697 |
| 472 | <i>Prunus</i> sp. 1 | 1 | 1.43 | | 240.53 | 0.007 | 0.027 | 0.000 | 0.034 |
| 473 | <i>Prunus</i> sp. 2 | 1 | 1.43 | 1 | 3,216.99 | 0.094 | 0.027 | 0.017 | 0.138 |
| | <i>Family Importance Value</i> | | | | | | - | | 1.434 |
| 47 | Rubiaceae (22) | | | | | | | | |
| 474 | <i>Adina minutiflora</i> | 5 | 7.14 | 5 | 6,614.45 | 0.193 | 0.134 | 0.086 | 0.413 |
| 475 | <i>Anthocephalus cadamba</i> | 20 | 28.57 | 28 | 98,044.19 | 2.861 | 0.538 | 0.481 | 3.880 |
| 476 | <i>Gardenia forsteniana</i> | 4 | 5.71 | 4 | 1,163.60 | 0.034 | 0.108 | 0.069 | 0.210 |
| 477 | <i>Gardenia</i> sp. | 1 | 1.43 | 1 | 349.67 | 0.010 | 0.027 | 0.017 | 0.054 |
| 478 | <i>Gardenia tubifera</i> . | 1 | 1.43 | 1 | 268.80 | 0.008 | 0.027 | 0.017 | 0.052 |
| 479 | <i>Ixora grandifolia</i> | 3 | 4.29 | 4 | 395.95 | 0.012 | 0.081 | 0.069 | 0.161 |
| 480 | <i>Lasianthus</i> sp. | 1 | 1.43 | 1 | 145.27 | 0.004 | 0.027 | 0.017 | 0.048 |
| 481 | <i>Nauclea jinghuhnii</i> | 2 | 2.86 | 2 | 516.50 | 0.015 | 0.054 | 0.034 | 0.103 |
| 482 | <i>Nauclea</i> sp. | 1 | 1.43 | 1 | 201.06 | 0.006 | 0.027 | 0.017 | 0.050 |
| 483 | <i>Nauclea subdita</i> | 1 | 1.43 | 2 | 714.43 | 0.021 | 0.027 | 0.034 | 0.082 |
| 484 | <i>Petunga</i> sp. | 1 | 1.43 | 1 | 196.07 | 0.006 | 0.027 | 0.017 | 0.050 |
| 485 | <i>Porterandia anisophylla</i> . | 10 | 14.29 | 10 | 2,952.47 | 0.086 | 0.269 | 0.172 | 0.527 |
| 486 | <i>Randia</i> sp. | 1 | 1.43 | 1 | 314.16 | 0.009 | 0.027 | 0.017 | 0.053 |
| 487 | <i>Tarenna winkleri</i> | 3 | 4.29 | 3 | 1,075.01 | 0.031 | 0.081 | 0.051 | 0.164 |
| 488 | <i>Timonius flavescentis</i> | 7 | 10.00 | 7 | 1,664.09 | 0.049 | 0.188 | 0.120 | 0.357 |
| 489 | <i>Timonius sericeus</i> | 2 | 2.86 | 2 | 1,082.48 | 0.032 | 0.054 | 0.034 | 0.120 |
| 490 | <i>Tricalysia malaccensis</i> | 1 | 1.43 | 1 | 93.31 | 0.003 | 0.027 | 0.017 | 0.047 |
| 491 | <i>Tricalysia singularis</i> | 3 | 4.29 | 3 | 320.17 | 0.009 | 0.081 | 0.051 | 0.142 |
| 492 | <i>Urophyllum arboreum</i> | 5 | 7.14 | 5 | 1,119.94 | 0.033 | 0.134 | 0.086 | 0.253 |
| 493 | <i>Urophyllum borneensis</i> | 2 | 2.86 | 2 | 282.03 | 0.008 | 0.054 | 0.034 | 0.096 |
| 494 | <i>Urophyllum corymbosum</i> | 2 | 2.86 | 2 | 268.05 | 0.008 | 0.054 | 0.034 | 0.096 |
| 495 | <i>Urophyllum polyneurum</i> | 12 | 17.14 | 23 | 2,867.25 | 0.084 | 0.323 | 0.395 | 0.801 |
| | <i>Family Importance Value</i> | | | | | | | | 7.759 |
| 48 | Sabaiaceae (1) | | | | | | | | |
| 496 | <i>Meliosma sumatrana</i> | 5 | 7.14 | 5 | 2,767.26 | 0.081 | 0.134 | 0.086 | 0.301 |
| | <i>Family Importance Value</i> | | | | | | | | 0.301 |
| 49 | Santalaceae (1) | | | | | | | | |
| 497 | <i>Scleropyrum wallichianum</i> | 5 | 7.14 | 5 | 1,237.81 | 0.036 | 0.134 | 0.086 | 0.256 |
| | <i>Family Importance Value</i> | | | | | | | | 0.256 |
| 50 | Sapindaceae (13) | | | | | | | | |
| 498 | <i>Allophylus cobe</i> | 1 | 1.43 | 1 | 162.86 | 0.005 | 0.027 | 0.017 | 0.049 |
| 499 | <i>Didymocarpus</i> sp. | 1 | 1.43 | 1 | 226.98 | 0.007 | 0.027 | 0.017 | 0.051 |
| 500 | <i>Didymocarpus longan</i> | 7 | 10.00 | 7 | 930.20 | 0.027 | 0.188 | 0.120 | 0.336 |
| 501 | <i>Guioa</i> sp. | 1 | 1.43 | 1 | 183.85 | 0.005 | 0.027 | 0.017 | 0.049 |
| 502 | <i>Nephelium cuspidatum</i> | 4 | 5.71 | 4 | 539.34 | 0.016 | 0.108 | 0.069 | 0.192 |
| 503 | <i>Nephelium lappaceum</i> | 4 | 5.71 | 4 | 635.56 | 0.019 | 0.108 | 0.069 | 0.195 |
| 504 | <i>Nephelium maingayi</i> | 1 | 1.43 | 1 | 122.72 | 0.004 | 0.027 | 0.017 | 0.048 |
| 505 | <i>Nephelium ramboutan-ake</i> | 5 | 7.14 | 5 | 2,089.81 | 0.061 | 0.134 | 0.086 | 0.281 |
| 506 | <i>Paranephelium</i> sp. | 1 | 1.43 | 1 | 141.03 | 0.004 | 0.027 | 0.017 | 0.048 |
| 507 | <i>Pometia pinnata</i> | 14 | 20.00 | 17 | 3,697.80 | 0.108 | 0.377 | 0.292 | 0.776 |
| 508 | <i>Rhysotoechia acuminata</i> | 1 | 1.43 | 1 | 543.25 | 0.016 | 0.027 | 0.017 | 0.060 |
| 509 | <i>Xerospermum laevigatum</i> | 2 | 2.86 | 2 | 613.26 | 0.018 | 0.054 | 0.034 | 0.106 |
| 510 | <i>Xerospermum xanthophyllum</i> | 3 | 4.29 | 4 | 1,460.11 | 0.043 | 0.081 | 0.069 | 0.192 |
| | <i>Family Importance Value</i> | | | | | | | | 2.383 |
| 51 | Sapotaceae (16) | | | | | | | | |
| 511 | <i>Chrysophyllum lanceolatum</i> | 1 | 1.43 | 1 | 576.80 | 0.017 | 0.027 | 0.017 | 0.061 |
| 512 | <i>Ganua motleyana</i> | 24 | 34.29 | 101 | 92,233.50 | 2.692 | 0.646 | 1.734 | 5.071 |
| 513 | <i>Madhuca ligulata</i> | 1 | 1.43 | 2 | 2,116.49 | 0.062 | 0.027 | 0.034 | 0.123 |
| 514 | <i>Madhuca magnifolia</i> | 6 | 8.57 | 7 | 8,087.12 | 0.236 | 0.161 | 0.120 | 0.518 |
| 515 | <i>Madhuca malaccensis</i> (| 1 | 1.43 | 3 | 1,620.98 | 0.047 | 0.027 | 0.051 | 0.126 |
| 516 | <i>Madhuca motleyana</i> | 3 | 4.29 | 4 | 3,721.56 | 0.109 | 0.081 | 0.069 | 0.258 |
| 517 | <i>Madhuca sericea</i> | 44 | 62.86 | 80 | 21,007.02 | 0.613 | 1.183 | 1.373 | 3.170 |
| 518 | <i>Madhuca sessiliflora</i> | 1 | 1.43 | 1 | 130.70 | 0.004 | 0.027 | 0.017 | 0.048 |
| 519 | <i>Palaquium calophyllum</i> | 10 | 14.29 | 14 | 6,978.71 | 0.204 | 0.269 | 0.240 | 0.713 |
| 520 | <i>Palaquium dasyphyllum</i> | 22 | 31.43 | 32 | 9,954.55 | 0.291 | 0.592 | 0.549 | 1.432 |
| 521 | <i>Palaquium ferox</i> | 10 | 14.29 | 12 | 9,806.05 | 0.286 | 0.269 | 0.206 | 0.761 |

| No. | Family and Species | Number of Occurrence in plots | Frequency (%) | Number of Trees (in 10.5 Ha) | Basal Area (Cm2) | Relative Basal Area (%) | Relative Frequency (%) | Relative Density (%) | Importance Value |
|--------------------------------|--------------------------------------|-------------------------------|---------------|------------------------------|------------------|-------------------------|------------------------|----------------------|------------------|
| 522 | <i>Palaquium rostratum</i> | 26 | 37.14 | 41 | 12,565.50 | 0.367 | 0.699 | 0.704 | 1.770 |
| 523 | <i>Palaquium sericeum</i> | 3 | 4.29 | 3 | 413.71 | 0.012 | 0.081 | 0.051 | 0.144 |
| 524 | <i>Palaquium</i> sp. | 1 | 1.43 | 1 | 1,590.43 | 0.046 | 0.027 | 0.017 | 0.090 |
| 525 | <i>Payena lucida</i> | 15 | 21.43 | 20 | 8,807.83 | 0.257 | 0.403 | 0.343 | 1.004 |
| 526 | <i>Payena sericea</i> | 1 | 1.43 | 1 | 107.51 | 0.003 | 0.027 | 0.017 | 0.047 |
| Family Importance Value | | | | | | | | | |
| 52 | Simarubaceae (1) | | | | | | | | |
| 527 | <i>Irvingia malayana</i> | 5 | 7.14 | 5 | 4,917.83 | 0.144 | 0.134 | 0.086 | 0.364 |
| Family Importance Value | | | | | | | | | |
| 53 | Sonneratiaceae (1) | | | | | | | | |
| 528 | <i>Duabanga moluccana</i> | 1 | 1.43 | 1 | 122.72 | 0.004 | 0.027 | 0.017 | 0.048 |
| Family Importance Value | | | | | | | | | |
| 54 | Sterculiaceae (9) | | | | | | | | |
| 529 | <i>Heritiera javanica</i> | 1 | 1.43 | 1 | 176.71 | 0.005 | 0.027 | 0.017 | 0.049 |
| 530 | <i>Heritiera littoralis</i> | 8 | 11.43 | 8 | 3,001.48 | 0.088 | 0.215 | 0.137 | 0.440 |
| 531 | <i>Heritiera simplicifolia</i> | 6 | 8.57 | 6 | 3,826.99 | 0.112 | 0.161 | 0.103 | 0.376 |
| 532 | <i>Pterocymbium tubulatum</i> | 8 | 11.43 | 8 | 1,943.91 | 0.057 | 0.215 | 0.137 | 0.409 |
| 533 | <i>Pterygota</i> sp. | 2 | 2.86 | 2 | 3,079.08 | 0.090 | 0.054 | 0.034 | 0.178 |
| 534 | <i>Scaphium macropodum</i> | 20 | 28.57 | 26 | 165,001.32 | 4.816 | 0.538 | 0.446 | 5.800 |
| 535 | <i>Sterculia gilva</i> | 4 | 5.71 | 5 | 2,058.76 | 0.060 | 0.108 | 0.086 | 0.253 |
| 536 | <i>Sterculia rubiginosa</i> | 7 | 10.00 | 7 | 2,062.43 | 0.060 | 0.188 | 0.120 | 0.369 |
| 537 | <i>Sterculia</i> sp.1 | 1 | 1.43 | 1 | 136.85 | 0.004 | 0.027 | 0.017 | 0.048 |
| Family Importance Value | | | | | | | | | |
| 55 | Symplocaceae (1) | | | | | | | | |
| 538 | <i>Symplocos odorartissima</i> | 1 | 1.43 | 1 | 100.29 | 0.003 | 0.027 | 0.017 | 0.047 |
| Family Importance Value | | | | | | | | | |
| 56 | Theaceae (3) | | | | | | | | |
| 539 | <i>Schima wallichii</i> | 5 | 7.14 | 5 | 4,723.370 | 0.138 | 0.134 | 0.086 | 0.358 |
| 540 | <i>Tetramerista glabra</i> | 8 | 11.43 | 11 | 18,722.63 | 0.546 | 0.215 | 0.189 | 0.950 |
| 541 | <i>Thea</i> sp. | 1 | 1.43 | 1 | 80.12 | 0.002 | 0.027 | 0.017 | 0.046 |
| Family Importance Value | | | | | | | | | |
| 57 | Thymelaeaceae (4) | | | | | | | | |
| 542 | <i>Aquilaria malaccensis</i> | 22 | 31.43 | 30 | 18,030.69 | 0.526 | 0.592 | 0.515 | 1.633 |
| 543 | <i>Gonostylus forbesii</i> | 1 | 1.43 | 1 | 113.10 | 0.003 | 0.027 | 0.017 | 0.047 |
| 544 | <i>Gonostylus macrophyllus</i> | 7 | 10.00 | 7 | 1,530.18 | 0.045 | 0.188 | 0.120 | 0.353 |
| 545 | <i>Gonostylus velutinus</i> | 4 | 5.71 | 4 | 1,242.89 | 0.036 | 0.108 | 0.069 | 0.213 |
| Family Importance Value | | | | | | | | | |
| 58 | Tiliaceae (4) | | | | | | | | |
| 546 | <i>Microcos crassifolia</i> | 3 | 4.29 | 3 | 403.18 | 0.012 | 0.081 | 0.051 | 0.144 |
| 547 | <i>Microcos hirsuta</i> | 2 | 2.86 | 2 | 202.39 | 0.006 | 0.054 | 0.034 | 0.094 |
| 548 | <i>Pentace laxiflora</i> | 21 | 30.00 | 30 | 9,843.85 | 0.287 | 0.565 | 0.515 | 1.367 |
| 549 | <i>Pentace triptera</i> | 1 | 1.43 | 1 | 174.37 | 0.005 | 0.027 | 0.017 | 0.049 |
| Family Importance Value | | | | | | | | | |
| 59 | Ulmaceae (2) | | | | | | | | |
| 550 | <i>Gironniera nervosa</i> | 26 | 37.14 | 54 | 19,101.88 | 0.557 | 0.699 | 0.927 | 2.184 |
| 551 | <i>Gironniera subaequalis</i> | 1 | 1.43 | 1 | 237.79 | 0.007 | 0.027 | 0.017 | 0.051 |
| Family Importance Value | | | | | | | | | |
| 60 | Urticaceae (1) | | | | | | | | |
| 552 | <i>Poikilospermum suaveolens</i> | 1 | 143.00 | 1 | 298.65 | 0.009 | 0.027 | 0.017 | 0.053 |
| Family Importance Value | | | | | | | | | |
| 61 | Verbenaceae (2) | | | | | | | | |
| 553 | <i>Teijsmanniodendron bogoriense</i> | 2 | 2.86 | 2 | 4,183.23 | 0.122 | 0.054 | 0.034 | 0.210 |
| 554 | <i>Teijsmanniodendron coriaceum</i> | 1 | 1.43 | 1 | 81.71 | 0.002 | 0.027 | 0.017 | 0.046 |
| Family Importance Value | | | | | | | | | |
| 62 | Violaceae (1) | | | | | | | | |
| 555 | <i>Rinorea benghalensis</i> | 17 | 24.29 | 28 | 13,946.90 | 0.407 | 0.457 | 0.481 | 1.345 |
| Family Importance Value | | | | | | | | | |
| TOTAL | | 3,762 | 5,515.86 | 5,848 | 3,500,424.54 | 102.162 | 101.187 | 100.378 | 303.726 |

324

REINWARDTIA

[VOL.12]

INSTRUCTION TO AUTHORS

Manuscripts intended for publication in *Reinwardtia* should be written either in English, French or German, and represent articles which have not been published in any other journal or proceedings. Each manuscript received will be considered and processed further if it is accompanied by signed statements given independently by two reviewers chosen by the author(s) attesting to its merits as well as its scientific suitability for publication in *Reinwardtia*.

Two printed copies (on A4 paper) of the manuscript of not more than 200 pages should be sent to Editors, together with an electronic copy prepared on Word Processor computer programme using Times New Roman letter type and saved as Rich Text File must be submitted.

For the style of presentation authors should follow the latest issue of *Reinwardtia* very closely. Title of the article should be followed by author's name and mailing address and one-paragraphed abstract in English (with French or German abstract for paper in French or German) of not more than 250 words. Keywords should be given below each abstract, on a separate page author(s) should the preferred running title of the article submitted.

Taxonomic keys should be prepared using the aligned-couplet type.

Strict adherence to the International Code of Botanica Monenclatural is observed, so that taxonamix and nomenclatural novelties should be clearly shown, Latin description for new taxon proposed should be provided, and the herbaria where type specimens are deposited should be indicated. Synonyms should be presented in the long form [name of taxon, author's name, year of publication, abbreviated journal or book title, volume (number): [page].

Maps, line drawing illustrations or photographs preferably should be prepared in landscape presentation to occupy two columns. Illustrations must be submitted as original art accompanying, but separated from the manuscript. On electronic copy, the illustration should be saved in .jpg or .gif format. Legends for illustrations must be submitted separately at the end of the manuscript.

Bibliography, list of literature cited or references follow the Harvard system.

For each paper published author(s) will receive 25 copies of reprints free of charge, any additional copies should be ordered in advance and the author(s) will be charged accordingly.

CONTENTS

| | Page |
|---|------|
| J.F. VELDKAMP. The correct name for the <i>Tetrastigma</i> (<i>Vitaceae</i>) host of <i>Rafflesia</i> (<i>Rafflesiaeae</i>) in Malesia and a (not so) new species..... | 261 |
| W.J.ODE WILDE & B.E.E. DUYFES. Miscellaneous South East Asian cucurbit news..... | 267 |
| M.A. RIFAI. <i>Endophragmiella bogoriensis</i> Rifai, spec. nov (<i>Hymenomycetes</i>)..... | 275 |
| M.A. RIFAI. Another note on <i>Podoconismegaspemias</i> Boedijn (<i>Hymenomycetes</i>)..... | 277 |
| TOPIK HID A W; M. ITO; T. YUKAWA. The phylogenetic position of the Papuasian genus <i>Sarcochilus</i> R.Br. (<i>Orchidaceae: Aeridinae</i>): evidence from molecular data..... | 281 |
| C.E. RIDSDALE. Notes on <i>MaiesiznNeonaucleea</i> | 285 |
| C.E. RIDSDALE. Thorny problems in the <i>Rubiaceae: Benkara, Fagerlindia and Oxyceros</i> | 289 |
| KUSWATAKARTAWINAIA, PURWANINGSIH, T. PARTOMIHARDJO, R. YUSUF, R. ABDULHADI, S. RISWAN. Floristics and structure of a lowland dipterocarp forest at Wanariset Samboja, East Kalimantan, Indonesia..... | 301 |
| RUGAiAH & S. SUNARTI. Two new wild species of <i>Averrhoa</i> (<i>Oxalidaceae</i>) from Indonesia..... | 325 |
| ATIKRETNOWATI. Anew Javanese species of <i>Marasmius</i> (<i>Trichlomataceae</i>)..... | 334 |

Reinwardtia is a LEPI accredited Journal (80/Akred-LIPI/P2MBI/5/2007)

HERBARIUM BOGORIENSE
BIDANG BOTANI
PUS AT PENELITIAN BIOLOGI - LIPI
BOGOR, INDONESIA