

# ZOO INDONESIA

## Jurnal Fauna Tropika

Akreditasi : 119/AKRED/LIPI/P2MBI/06/2008 (Predikat B)

---

---

|                                                                                                                                                                                                                                           |    |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| AN INVENTORY OF REPTILES AND AMPHIBIANS IN NORTH-WEST OF SIBERUT ISLAND, WEST SUMATERA. <i>Irvan Sidik</i> .....                                                                                                                          | 35 |
| PENGARUH PEMBERIAN HORMON METHYLTESTOSTERONE PADA LARVA IKAN GUPPY ( <i>Poecilia reticulata</i> ) TERHADAP PERUBAHAN JENIS KELAMIN. <i>Gleni Hasan Huwoyon, Rustidja &amp; Rudy Gustiano</i> .....                                        | 47 |
| ASOSIASI KUMBANG KOTORAN (COLEOPTERA : SCARABAEIDAE) DENGAN TUNGAU MACROCHELIDAE DI CAGAR ALAM PANGANDARAN (JAWA BARAT) DAN TAMAN NASIONAL GUNUNG MERAPI (YOGYAKARTA). <i>Dhian Dwibadra, Sri Hartini &amp; Rosichon Ubaidillah</i> ..... | 53 |
| AN OVERVIEW ON THE ECOLOGY OF VARANID LIZARDS. <i>Evy Arida</i> .....                                                                                                                                                                     | 65 |
| FAUNA TANAH PADA STRATIFIKASI LAPISAN TANAH BEKAS PENAMBANGAN EMAS DI JAMPANG, SUKABUMI SELATAN. <i>Erniwati</i> .....                                                                                                                    | 83 |



### **Ketua Redaksi**

Dr. Dede Irving Hartoto (Limnologi)

### **Anggota Redaksi**

Dr. Hagi Yulia Sugeha (Oseanologi)

Dr. Rosichon Ubaidillah (Entomologi)

Dr. Dewi Malia Prawiradilaga (Ornitologi)

Ir. Ike Rachmatika MSc. (Ikhtiologi)

### **Sekretaris Redaksi & Produksi**

Rochmanah S.Kom

Muhamad Ridwan

### **Mitra Bestari**

Drs. Haryono MSi.

Prof. Dr. Woro A. Noerdjito

Dra. Hellen Kurniati

Dr. Sih Kahono

### **Alamat Redaksi**

Zoo Indonesia

Bidang Zoologi, Puslit Biologi LIPI

Gd. Widyasatwaloka

Jl. Raya Bogor-Jakarta KM. 46

Cibinong 16911

Telp. (021) 8765056

Fax. (021) 8765068

[zooindonesia@yahoo.com](mailto:zooindonesia@yahoo.com) ([www.biologi.lipi.go.id](http://www.biologi.lipi.go.id))

**Akreditasi: 119/AKRED/LIPI/P2MBI/06/2008 (Predikat B)**

---

Masyarakat Zoologi Indonesia (MZI) adalah suatu organisasi profesi dengan anggota terdiri dari peneliti, pengajar, pemerhati dan simpatisan kehidupan fauna tropika, khususnya fauna Indonesia. Kegiatan utama MZI adalah memasyarakatkan tentang ilmu kehidupan fauna tropika Indonesia, dalam segala aspeknya, baik dalam bentuk publikasi ilmiah, publikasi popular, pendidikan, penelitian, pameran ataupun pemantauan.

Zoo Indonesia adalah sebuah jurnal ilmiah di bidang fauna tropika yang diterbitkan oleh organisasi profesi keilmianah Masyarakat Zoologi Indonesia (MZI) sejak tahun 1983. Terbit satu tahun satu volume dengan dua nomor (Juni & Nopember). Memuat tulisan hasil penelitian dan tinjauan ilmiah yang berhubungan dengan aspek fauna, khususnya wilayah Indonesia dan Asia. Publikasi ilmiah lain adalah Monograph Zoo Indonesia - Seri Publikasi Ilmiah, terbit tidak menentu.

---

## PETUNJUK PENULISAN

Zoo Indonesia merupakan jurnal ilmiah di bidang zoologi yang diterbitkan oleh organisasi profesi Masyarakat Zoologi Indonesia (MZI) sejak tahun 1983. Terbit setiap tahun satu volume dengan dua nomor (Juni & Nopember). Bentuk naskah terbagi atas naskah utama, berupa hasil penelitian yang utuh dan belum diterbitkan; naskah penunjang, berupa catatan pendek dari hasil penelitian yang dirasakan perlu cepat untuk diinformasikan; dan review, suatu kajian ilmiah yang menyeluruh, lengkap dan cukup mendalam tentang suatu topik berdasarkan rangkuman hasil penelitian beberapa peneliti. Bidang pembahasan dalam Zoo Indonesia meliputi fauna, pada semua aspek keilmuan seperti Biosistematik, Fisiologi, Ekologi, Molekuler, Pemanfaatan, Pengelolaan, Budidaya dll. Tata cara penulisan adalah:

1. Naskah ditulis dalam bahasa Indonesia atau Inggris. Diketik pada format kertas A-4 dengan jarak spasi 1.5, Arial, font 10. Ukuran margin atas & bawah 2.54 cm, kanan & kiri 3.00 cm.
2. Sistematik penulisan :
  - a. **Judul**, singkat dan jelas, penyertaan anak judul sebaiknya dihindari. Diketik dengan huruf besar, dihitamkan, terkecuali pada nama Latin, dengan huruf miring.
  - b. **Nama dan alamat penulis** beserta alamat elektronik, ditulis lengkap tanpa ada singkatan, ditempatkan di bawah judul.
  - c. **Abstrak**, merupakan intisari naskah, ditulis tidak lebih dari 200 kata dan dituangkan dalam satu paragraf. Dibawah abstrak dicantumkan kata kunci maksimal lima kata. Berbahasa Indonesia dan Inggris.
  - d. **Pendahuluan**, ditulis singkat mengenai latar belakang penelitian, permasalahan, hal-hal yang telah diketahui, pendekatan yang dikembangkan dalam memecahkan masalah dan pencapaian tujuan penelitian.
  - e. **Materi & Metode**, menerangkan secara jelas tata cara penelitian, waktu dan tempat penelitian, metode yang digunakan, analisa statistik, sehingga mampu diulang kembali oleh pihak lain atau mengkaji ulang runtutan tata cara penelitian. Data mengenai nomor aksesi spesimen, asal-usul spesimen, lokasi atau hal lain yang dirasa perlu untuk penelusuran kembali, ditempatkan sebagai Lampiran, setelah Daftar Pustaka.
  - f. **Hasil & Pembahasan**, menyajikan hasil penelitian yang diperoleh, sekaligus mengupas dan membahas hasil penelitian, membandingkannya dengan hasil temuan peneliti lain dan penjabaran implikasi dari penelitian yang diperoleh. Penyertaan ilustrasi dalam bentuk Tabel, Gambar atau Sketsa hendaknya berwarna hitam putih. Khusus foto dapat hitam putih atau berwarna, format JPEG. Sitiran untuk menghubungkan nama penulis dan tahun terbitan tidak menggunakan tanda koma. Bila ada beberapa tahun penulisan yang berbeda untuk satu penulis yang sama digunakan tanda penghubung koma, serta tanda gabung bentuk titik koma pada kumpulan sitiran yang mengelompok tetapi berbeda penulis (Hasyim 2005, 2006; Gunawan 2004). Nama penulis yang lebih dari dua orang ditulis *et al.* (jurnal terbitan asing) atau dkk. (jurnal terbitan lokal). Kata penghubung diantara dua penulis menggunakan tanda &.
  - g. **Kesimpulan**, merupakan rangkuman dari keseluruhan hasil penulisan.
  - h. **Daftar Pustaka**, menyajikan semua pustaka yang dipergunakan dalam naskah.

- Flannery, T. 1990. Mammals of New Guinea. Robert Brown & Associates.  
New York.
- Nelson, M.E & L.D Mech. 1987. Demes with a Northeastern Minnesota Deer Population. In: B.D Chepko-Sade & Z Tanghaphin (eds.) Mammalian Dispersal Pattern-The Effect of Social Structure on Population Genetics. University of Chicago Press. 230-243.
- Youngson, R.W. 1970. Rearing red deer calves. Journal of Wildlife Management 34:467-470.

3. **Ucapan Terima Kasih**, sebagai penghargaan atas pihak-pihak yang dirasa layak diberikan.
4. Naskah lengkap dapat dikirim melalui alamat elektronik atau pos. Bila melalui pos dikirim dua rangkap, satu diantaranya tanpa nama dan alamat penulis, disertai disket/compact disk.

Redaksi Zoo Indonesia  
d/a Bidang Zoologi - Puslit Biologi LIPI  
Jl. Raya Bogor-Jakarta Km. 46  
Cibinong 16911  
zooindonesia@yahoo.com

MONOGRAPH ZOO INDONESIA adalah publikasi ilmiah lainnya yang terbit tidak menentu. Berisi bahasan yang sangat mendalam dan holistik mengenai satu aspek pada tingkat jenis (species) ataupun permasalahan.

Terakreditasi berdasarkan SK Kepala LIPI no. 683/D/2008  
No. Akreditasi: 119/AKRED/ LIPI/P2MBI/06/2008 (Predikat B) periode Juni 2008-2011

Penerbitan Volume 17 Nomor 2 tahun 2008 ini didanai oleh DIPA Puslit Biologi LIPI  
T.A 2009



## AN OVERVIEW ON THE ECOLOGY OF VARANID LIZARDS

Evy Arida

Herpetology Laboratory, Museum Zoologicum Bogoriense-LIPI  
Jln. Raya Bogor-Jakarta Km 46, Cibinong 16911  
e-mail: evyarida@lycos.com

### ABSTRACT

**Arida, E. 2008. An overview on the ecology of Varanid lizards. Zoo Indonesia 17(2): 67-82.** Body size is a morphological character that can be useful to estimate the size of home range in varanids. It may also be used to cue for habitat type and mode of life of varanid lizards. Nevertheless, although body size may be a good predictor of home range size, it may not be useful to infer population density, because non-territoriality nature of varanid lizards generates home range overlaps. Although non-territoriality nature may hinder inference of population density, it can signal for high density through the signature of aggression. However, the magnitude of density itself would not be well quantified. An approach to population density estimation is the use of reproductive biology data. Reproductive biology data of wild varanids can be used to base a projection on population trends. Despite the notion, data on reproduction from wild populations seems to be scarce, especially from some regions in Asia. Regular population monitoring for general census and reproductive status is definitely still needed to allow for sound estimates of population density and its dynamics.

**Keywords:** body size, home range, varanids, density, reproduction.

### ABSTRAK

**Arida, E. 2008. Tinjauan ekologi kelompok biawak. Zoo Indonesia 17(2): 67-82.** Daerah jelajah biawak dapat diperkirakan melalui ukuran tubuhnya. Ukuran tubuh biawak dapat pula digunakan untuk mencirikan tipe habitat dan cara hidupnya. Meskipun demikian, ukuran tubuh tidak dapat digunakan untuk memperkirakan kepadatan populasi, yang disebabkan oleh daerah jelajah yang tumpang tindih. Daerah jelajah yang tumpang tindih ini terjadi karena biawak tidak mempunyai teritori, yaitu daerah yang dipertahankan dari kedatangan hewan lain. Dengan keadaan ini, masih ada kemungkinan terjadinya agresi, yang merupakan suatu penanda tingginya kepadatan populasi tersebut, walaupun kepadatan populasi ini tidak dapat diperkirakan dengan pasti. Perkiraan kepadatan populasi dapat dilakukan dengan pendekatan yang berdasarkan pada data perkembangbiakan, yang pada prinsipnya digunakan untuk memproyeksikan dinamika populasi. Namun demikian, data perkembangbiakan dari populasi liar belum banyak tersedia, terutama data dari beberapa wilayah di Asia. Pemantauan populasi yang teratur dalam rangka cacah jiwa dan pemantauan status perkembangbiakan, masih diperlukan untuk memperkirakan kepadatan populasi dan dinamikanya dengan benar.

**Kata kunci:** ukuran tubuh, daerah jelajah, biawak, kepadatan, perkembangbiakan.

## INTRODUCTION

An attempt to characterise varanid lizards' population can possibly be made by taking ecological properties such as home range size and reproductive biology information into account. Morphological characteristics, for example body size, can be used to predict ecological repertoire. Here, a possible relationship between body size, home range, and reproductive biology of varanid lizards will be elaborated to characterise the general trend on the population, based on available published literatures.

### Population

Population ecology is the study of interaction among organisms in relation to environmental aspects with emphasize on demography (*Begon et al.* 1996). It has a tight connection with conservation biology, because assessment of population viability for conservation is based on population size estimates together with the dynamics of this population. Despite commercial exploitation of many species and endemicity of several varanid species, there are probably only a handful population studies on these lizards. This is probably due to logistical difficulties in applying mark-recapture technique especially when involving large individuals. On the other hand, the complexity of integrated population study involving demographic parameters including those requiring long-term continuous observation e.g., reproductive biology, is another factor that hinders data collection.

From a taxonomic perspective, continuous new species discoveries and species complex (Böhme 2003) may confound long-term population monitoring and census. In addition, the broad ecological repertoire and agility of monitor lizards may render difficulties and biases in census.

In spite of the scarce full-length data on monitor lizard populations, a number of data from related studies e.g., spatial ecology and reproductive biology might be useful to predict trends in varanid populations. A range of factors affecting the magnitude and dynamics of monitor lizard populations has been postulated and includes basic demographic parameters such as size, age, and sex. These parameters affect population size indirectly through a mechanism that regulates patterns of individual movement and/or home range (Perry & Garland 2002).

Population density and species distribution can also be affected by environmental parameters such as seasons (Phillips 1995; Guarino 2002) and behavioural parameters such as habitat preference (Bondarenko 1989; Thompson *et al.* 1999) and social system (Stanner & Mendelssohn 1987). In addition, intraspecific competition may directly regulate population size and dynamics of these large lizards (Luiselli *et al.* 1999). Furthermore, energetic-related factors such as dietary shift and mode of life i.e., arboreality, terrestriality, aquatic) may also influence the magnitude and dynamics of varanid populations.

### Ecological & behavioural repertoire

Monitor lizards are generally good runners, climbers, and/or swimmers. This repertoire in movement suggests an ability to move through variable habitats. *V. bengalensis* and *V. salvator* in Asia and *V. gouldii* and *V. tristis* in Australia utilised many different habitats and are among varanid species that are widely distributed (King & Green 1999). The broad diet of some monitor lizard generally suggests an opportunistic foraging nature (Shine 1986). Thus, some species are plastic and may adapt to a wide range of habitat types. However, some species are specialised to live in a particular

habitat type. Species within the *Varanus prasinus* group are specialised to live on trees, which indicates habitat restriction to forested areas and dietary restriction to preys available in this habitat.

The broad ecological repertoire of some varanid lizards may enhance their successful dispersal. Following multiple successful dispersal, broad ecological repertoire may give rise to phenotypic plasticity suggesting invasiveness (Agrawal 2001). Evidence of plastic behaviour in varanid lizards was shown by the Nile monitor (*Varanus niloticus*), a species of African origin, that warrants population monitoring and eradication in south western Florida, USA (Enge *et al.* 2004). Similarly, the Pacific monitor (*Varanus indicus*) is considered as invasive in Guam, on the island of Hawaii, USA (Bergman *et al.* 2000).

Persistence in disturbed area, where habitat are changed and human continuously present, also suggests a signature of plastic adaptation. This is shown by the Asian Water monitor (*V. salvator*), which has a wide distribution in Asia (Gaulke *et al.* 1999). Nevertheless, there is no evidence that *V. salvator* is an invasive species.

### **Body size & home range**

Differential body size, in terms of Snout-Vent Length (SVL), within the extant varanid lizards could have been a signature of evolutionary forces that is reflected on morphological characteristics. The relatively rapid evolution of body size among varanid lizards seems to be a fine-tuning on habitat variation, given three different lineages of dwarf, large, and gigantic species occurring within the boundary of Australian continent (Pianka 1995). Thus, morphological character i.e., variation in species body size (SVL) of varanid lizards may have an association with the type of habitat a species living

in. Further on, species with broad ecological repertoire seem to be associated with a few habitat types or even changed habitats. Such species are presumably of medium SVL, because medium SVL is likely to enable effective distant movement beyond the habitat boundary. An eventual consequence of adaptability to new area (habitat) may give rise to wider foraging area or larger home range size. Nevertheless, in large species that is relatively adaptable to changed habitats such as *V. salvator* (Gaulke *et al.* 1999), probably only medium size individuals that make the cross beyond habitat boundaries. This is true since juveniles tend to live on trees, whereas giant adults are very rarely caught during harvest or captures for study purposes (E. Arida, pers. obs.).

Home range is defined as the area traversed by animals during routine activities. It accounts for the behavioural and physiological demands of the individuals (Perry & Garland 2002). A fraction of an animal's home range that is patrolled and defended from intruders for access to resources is commonly defined as territory (Simon 1975; Kaufmann 1983). Varanids are generally non-territorial and many species within the family have a large home range that overlaps (Stanner & Mendelsohn 1987; Thompson 1994; Lenz 1995; Gaulke *et al.* 1999; King & Green 1999; Sweet 1999; King *et al.* 2002) within and between the sexes (Stanner & Mendelsohn 1987; Thompson 1994; Sweet 1999).

Despite some published ecological data on home range of varanid lizards, there has not been any published trend on home range size plotted with variation in SVL as a measure of body size. This is probably due to the relatively few published data on home range (King & Green 1999) as well as inadequate uniformity in published SVL measurements. However, correlation

between home range size and body mass has been formulated as:

$$H = 0.0307 \times W - 0.5068$$

where H= home range (ha) and W= body weight (g) (Guarino 2002).

Table 1 shows some data on home range size and maximum SVL of 53 *Varanus* from three continents. It is always the case that males of a species have larger home range than conspecific females, presumably because adult males are generally larger than females. Accordingly, larger species are likely to have larger size of home range relative to smaller species. Home range size may have a direct relationship with SVL at adult stage. Within species, this direct relationship may be more obvious in large monitors than in small or medium size monitors. The adult males of large species e.g. *V. komodoensis* and *V. giganteus* are larger than conspecific females (Table 1), whereas SVL of adult males and females of some small to medium size monitors are likely to be similar (monomorphic) e.g. *V. mitchelli* and *V. mertensi*, both of which are semi-aquatic (Shine 1986), and may show no significant distinction in home range size between the sexes.

Direct relationship between SVL and home range size between species can be exemplified in the Pygmy Goanna (*V. brevicauda*), the smallest species among all *Varanus*. *Varanus brevicauda* is relatively sedentary compared to its larger sympatric congeneric i.e., *V. gilleni*, *V. eremius*, and *V. caudolineatus* occurring on the central and western deserts of Australia (James 1996). This suggests that small species travel shorter distance and therefore cover relatively smaller area, making their home range also smaller. Size of activity area in monitor lizards is also associated with mode of life (Thompson 1994; King *et al.* 2002). Many arboreal species are smaller in size compared to some terrestrial

species, which tend to have larger home range. Nonetheless, data on home range size of arboreal monitors are relatively rare and many species have non-strict mode of life.

There are some other factors that directly or indirectly affect the size of home range. For example, variation in home range size of monitor lizards can be related to climatic factors such as rainfall (Auffenberg 1988; Auffenberg 1994) and temperature (King 1980). During wet season that is associated with abundance of prey insect, *V. bengalensis* moved a greater distance (Auffenberg 1994), suggesting larger home range during this period. Thus, rainfall is the indirect factor that indicates prey availability, whereas insect abundance is the direct factor that drives animals to forage. The Sand goanna, *V. gouldii*, were found remain in their burrow during extreme temperatures (King 1980), suggesting no movement during this period. Therefore, one can expect smaller home range for this species during harsh climate than during the period of mild temperature.

Intra and interspecific interaction between individuals may also set boundary to home range size. Since monitor lizards are non-territorial, one can expect that these interactions between individuals within a species or individuals between species would not be easily observed. The degree of interspecific competition among varanid species seems to be minimised by niche segregation (Wikramanayake & Dryden 1993) and this may be corroborated by their broad dietary range (Shine 1986). Niche segregation seems to be also the strategy to minimise intraspecific competition, for example in adult avoidance by juveniles and age-class segregated niche e.g. in *V. komodoensis* (Auffenberg 1981; Imansyah *et al.* 2007). In this regard, niche segregation within and between

species may be used to locate home range or points of species distribution. However, segregated niche might be inadequate to index home range size, if there is no accompanying data on the animals' movement.

Geographic barrier in terms of island may also limit body size, including SVL, and the consequent size of home range. Varanid lizards living on islands are expected to be smaller than those living on mainland, particularly due to depletion of resources and/or increased aggression. This can probably be explained by the absence of predator on island and the non-territorial nature of monitor lizards, which may allow rapid increase of density (Buckley & Jetz 2007) promoting resource overexploitation leading to aggression resulting from severe competition (Case & Schwaner 1993; Pianka 1995). Thus, one can expect overlapping home ranges as a consequence of non-territoriality despite the reduced space on island. Further, home range size is expected to be smaller on island as a function of reduced body size, despite the reduced total area. This means that size of home range in varanids is not directly dependent upon population density, but it depends more directly on individual body size that can be influenced by competition as a consequence of population density.

The smallness of island varanids can be related to differential physiological requirements (Nagy 2005) of individuals surviving on island community relative to their mainland conspecific. During high density on islands, lizards may compensate growth energy for aggression, allowing slow growth that possibly results in reduced body size. Therefore, varanid lizards living on islands might have been selected for their physiology and body size. The Komodo dragon (*V. komodoensis*) is the largest living varanid species

inhabiting islands of several different sizes. The top largest individuals from the larger island of Rinca are relatively larger than the top largest individuals from the much smaller island of Gili Motang (Jessop *et al.* 2007). Although there is no direct evidence of severe competition, dragons on this small island may have been selected for their relatively smaller body size and the related physiological requirement.

### **Home range & population density**

The size of home range in varanid lizards might be useful to estimate population density as in mammals (McNab 1963). However, non-territorial spacing of monitor lizards that is plotted as overlapping convex polygons (Weavers 1993) may bias density estimate. This is because overlapping home ranges represent more than one individual on a given space, allowing bias in the extrapolation to the total distribution area. Nonetheless, non-territoriality in varanids may be useful to cue for change in population density. The absence of territorial maintenance in varanids allows for aggression that is promoted by competition during a rapid increase of density. Unfortunately, little is known about the intraspecific competition among varanid lizards, although cannibalism is known to occur in some species e.g. *V. griseus*, *V. niloticus*, *V. salvator* and *V. komodoensis* (Auffenberg 1981; Lenz 2004).

Cannibalism can be regarded as aggression by which severe intraspecific competition is possibly manifested. Nevertheless, the possible relationship between cannibalism and population density still remains to be tested, since there has not been any direct evidence to justify this hypothesis. Population density of Varanid lizards seems to be also affected by geographical boundary, since most island varanids are likely restricted to

dispersal (Buckley & Jetz 2007). There is probably no significant difference in home range size between island and mainland populations, due to the non-territorial spacing of these lizards. This means, island population density can be limited by the presence of geographical barrier (sea water), without the size of individual home range itself being changed accordingly. Population density estimate based on home range data alone seems to be unreliable, given the possible bias generated from overlapping home ranges. Further, reliable population density estimate is almost impossible to obtain without direct census. Meanwhile, population dynamics data are the series of census data obtained from long-term periodic population monitoring. Differential mode of life in varanid lizards is another factor that hinders generalisation of home range size among all members. In addition, the elusive and secretive nature of a species can also reduce the accuracy of an estimate (Stanner 2004).

### **Body size & reproductive biology**

All species of *Varanus* are egg-layers (Shine 1986). Body size has an influence on clutch sizes, with females of larger species having larger and more variable clutch sizes than those of smaller species (Thompson & Pianka 2001). This implies that larger species have higher reproductive output, although low offspring survival seems to counterbalance. In Komodo monitor, *V. komodoensis*, only about ¼ of clutch size is estimated to reach reproductive age, partly caused by predation of eggs and cannibalism of large adult conspecific (Auffenberg 1981). In Bengal monitor, *V. bengalensis*, about half of neonates dies by the end of their second year (Auffenberg 1994).

There is only a few long-term studies on reproduction of free-ranging monitor lizards but captive animals have provided a relatively large amount of data (Bennett et al. 1998). Meanwhile,

dissected museum specimens can be used to infer reproductive cycle. Monitor lizards appear to be conservative in the length of gestation but highly variable in length of incubation, which may be related to period of prey abundance in the wild and can help maximising offspring survival (Phillips & Millar 1998). Table 2 shows data on reproductive characteristics, some of which are from captive specimens. Small varanids seem to lay multiple clutches over a year (e.g. *V. melinus*, *V. rudicollis*, *V. prasinus*), each with a few to just over a dozen eggs. Large species tend to lay one clutch per year within total deposition time of a few weeks e.g. *V. komodoensis* (Auffenberg 1981).

### **Reproductive biology & population density**

A projection of population density on varanids is very difficult without reproductive success data from free-ranging animals. However, long-term monitoring for reproduction in the wild is logistically expensive, especially for tropical Asian species, since mating or egg deposition may extend to monsoon season (Auffenberg 1994; Shine et al. 1996). However, there are already details on reproduction from species inhabiting tropical Australia. Reproductive status and population density of traded species, for instance from Indonesia, however, can be monitored through the catch rate, although indirect data may not be as reliable as direct observation. Shine et al. (1996, 1998) were able to gather data on reproductive status, size, and food habits of commercially harvested *V. salvator* based on stretched skins in a holding facility in South and North Sumatra, Indonesia. Zoo captives are also useful to estimate reproductive cycle, although proof of reproductive synchrony with wild animals may not always be found (Table 2).

Conservative estimate of surviving individuals of about a quarter of original

clutch size may infer yearly recruitment close to reality, given many negative factors affecting hatching rate and survival of young animals such as predation and cannibalism. Nevertheless, direct census data is still needed repeatedly for a projection on population trend to base on.

## REFERENCES

- Agrawal, A. A. 2001. Phenotypic plasticity in the interactions and evolution of species. *Science* 294: 321.
- Auffenberg, W. 1981. The behavioral ecology of the Komodo monitor. Gainesville, University of Florida Presses.
- Auffenberg, W. 1988. Gray's monitor lizard. Gainesville, Florida, University Presses of Florida.
- Auffenberg, W. 1994. The Bengal monitor. Gainesville, University Presses of Florida.
- Begon, M., M. Mortimer & David J. Thompson. 1996. Population Ecology: A unified study of animals and plants, Blackwell Publishing.
- Bennett, D., T. Wilms & B. Bartholomew. 1998. Monitor lizards: natural history, biology, and husbandry, Chimaira.
- Bergman, D. L., M. D. Chandler & A. Locklear. 2000. The economic impact of invasive species to wildlife services' cooperators. USDA National Wildlife research Symposia Human Conflicts with Wildlife: Economic Considerations. Lincoln, University of Nebraska.
- Böhme, W. 2003. Checklist of the living monitor lizards of the world (family Varanidae). *Zoologische Verhandelingen Leiden*. 341.
- Bondarenko, D. A. 1989. Distribution and population density of the desert monitor *Varanus griseus* in the Karshy steppe zone Uzbek SSR USSR. *Byulleten Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii* 94(3):24-32.
- Buckley, L. B. & W. Jetz. 2007. Insularity and the determinants of lizard population density. *Ecology Letters* 10(6): 481-489.
- Case, T. J. & T. D. Schwaner. 1993. Island/mainland body size differences in Australian varanid lizards. *Oecologia* 94: 102-109.
- Enge, K. M., K. L. Krysko, K.R. Hankins, T.S. Campbell & F.W. King. 2004. Status of the Nile monitor (*Varanus niloticus*) in southwestern Florida. *Southeastern Naturalist* 3(4): 571-582.
- Gaulke, M., W. Erdelen & F. Abel. 1999. A radio-telemetric study of the Water monitor lizard (*Varanus salvator*) in North Sumatra, Indonesia. *Mertensiella (Advances in Monitor Research II)* 11: 63-78.
- Guarino, F. 2002. Spatial ecology of large carnivorous lizard, *Varanus varius* (Squamata: Varanidae). *Journal of Zoology* 258: 449-457.
- Imansyah, M. J., T. S. Jessop, C. Ciofi & Z. Akbar. 2007. Ontogenetic differences in the spatial ecology of immature Komodo dragons. *Journal of Zoology* 1-9.
- James, C. D. 1996. Ecology of the Pygmy Goanna (*Varanus brevicauda*) in Spinifex grasslands of Central Australia. *Australian Journal of Zoology* 44(2): 177-192.
- Jessop, T. S., T. Madsen, C. Ciofi, M.J. Imansyah, D. Purwandana, H. Rudiharto, A. Arifiandy & J.A. Phillips. 2007. Island differences in population size structure and catch per unit effort and their conservation implications for Komodo dragons. *Biological Conservation* 135: 247-255.
- Kaufmann, J. H. 1983. On the definitions and functions of dominance and territoriality. *Biological Review* 58(1): 1-20.
- King, D. 1980. The thermal biology of free-living Sand goanna (*Varanus gouldii*) in southern

- Australia. Copeia 1980(4): 755-767.
- King, D. & B. Green. 1999. Monitors The biology of varanid lizards. Malabar, Krieger Publishing Company.
- King, D. R., E. R. Pianka & B. Green. 2002. Biology, ecology, and evolution. Komodo dragon biology and conservation. J. B. Murphy, C. Ciofi, d. L. Panouse and T. Walsh (edit.). Washington, Smithsonian Institution Press.
- Lenz, S. 1995. Zur Biologie und Ökologie des Nilwarans, *Varanus niloticus* (Linnaeus, 1766) in Gambia, Westafrika. Mertensiella 5.
- Lenz, S. 2004. *Varanus niloticus*. Varanoid lizards of the world. E. R. Pianka, D. R. King & R.A. King (edit.). Bloomington, Indiana University Press.
- Luiselli, L., G. C. Akani & D. Capizzi. 1999. Is there any interspecific competition between dwarf crocodiles (*Osteolaemus tetraspis*) and Nile monitors (*Varanus niloticus ornatus*) in the swamps of central Africa? A study from south-eastern Nigeria. Journal of Zoology 247: 127-131.
- McNab, B. K. 1963. Bioenergetics and the determination of home range size. American Naturalist 97(894): 133-140.
- Nagy, K. A. 2005. Field metabolic rate and body size. Journal of Experimental Biology 208: 1621-1625.
- Perry, G & T. Garland Jr. 2002. Lizard home range revisited: effects of sex, body size, diet, habitat, and phylogeny. Ecology 83(7): 1870-1885.
- Phillips, J. A. 1999. Movement patterns and density of *Varanus albigularis*. Journal of Herpetology 29(3): 407-416.
- Phillips, J. A. & R. P. Millar 1998. Reproductive biology of the White-throated Savanna monitor, *Varanus albigularis*. Journal of Herpetology 32: 366-377.
- Pianka, E. R. 1995. Evolution of body size: varanid lizards as a model system. The American Naturalist 146(3): 398-414.
- Shine, R. 1986. Food habits, habitats, and reproductive biology of four sympatric species of varanid lizards in tropical Australia. Herpetologica 42(3): 346-360.
- Shine, R & P. S. Harlow. 1996. The biology of water monitors *Varanus salvator* in southern Sumatra. Biological Conservation 77(2-3): 125-134.
- Simon, C. A. 1975. The influence of food abundance on territory size in the iguanid lizard *Sceloporus jarrovi*. Ecology 56: 993-998.
- Stanner, M. 2004. *Varanus griseus*. Varanoid lizards of the world. E. R. Pianka, D. R. King & R. A. King. Bloomington (edit.), Indiana University Press.
- Stanner, M. & H. Mendelssohn. 1987. Sex ratio, population density, and home range of the desert monitor *Varanus griseus* in the southern coastal plain of Israel. Amphibia-Reptilia 8(2): 153-164.
- Sweet, S. S. 1999. Spatial ecology of *Varanus glauerti* and *Varanus glebopalma* in Northern Australia. Mertensiella (Advances in Monitor Research II) 11: 317-366.
- Thompson, G. 1994. Activity area during the breeding season of *Varanus gouldii* (Reptilia:Varanidae) in an urban environment. Wildlife Research 21(6): 633-641.
- Thompson, G. & E. R. Pianka. 2001. Allometry of clutch and neonate sizes in monitor lizards (Varanidae: *Varanus*). Copeia 2001(2): 443-458.
- Thompson, G. G., M. De Boer & E.R. Pianka. 1999. Activity areas and daily movements of an arboreal monitor lizard, *Varanus tristis*

(Squamata: Varanidae) during the breeding season. Australian Journal of Ecology 24: 117-122.  
Weavers, B. W. 1993. Home range of male Lace monitors, *Varanus varius* (Reptilia: Varanidae) in sout-eastern Australia. Wildlife Research 20(3): 303-313.

Wikramanayake, E. D. & G. L. Dryden.1993. Thermal ecology of habitat and microhabitat use by sympatric *Varanus bengalensis* and *Varanus salvator* in Sri Lanka. Copeia 3: 709-714.

Table 1. Body sizes, home range sizes and habitat of genus *Varanus* by continental groups.

| No            | Species                  | Home range (ha)                 | Density (per ha)         | Max. adult SVL (cm) | Mode of life                   | Reference                                                |
|---------------|--------------------------|---------------------------------|--------------------------|---------------------|--------------------------------|----------------------------------------------------------|
| <b>Africa</b> |                          |                                 |                          |                     |                                |                                                          |
| 1             | <i>V. albigularis</i>    | 18.3 ± 2.4 (♂)<br>6.1 ± 0.6 (♀) | --                       | ?                   | Terrestrial, arboreal          | Philipps (1995)<br>Branch (1988)                         |
| 2             | <i>V. exanthematicus</i> | --                              | 0.87/ hour<br>3.57 (juv) | 75.0                | Terrestrial, arboreal, aquatic | Bennett (2000)<br>Bennett (2004)                         |
| 3             | <i>V. griseus</i>        | 7.5-1950                        | 0.2                      | 58.3*               | Terrestrial                    | Stanner (2004)                                           |
| 4             | <i>V. niloticus</i>      | 5 (♂)<br>1.5 (♀)                | 1.14                     | 96.0*               | Terrestrial, arboreal, aquatic | Lenz (1995)<br>Lenz (2004)                               |
| 5             | <i>V. ornatus</i>        | --                              | --                       | 92.25*              | Terrestrial, arboreal, aquatic | Böhme & Ziegler (2004)                                   |
| 6             | <i>V. yemenensis</i>     | --                              | --                       | 59.0                | Terrestrial, arboreal,         | Gasperetti in Böhme (1989)<br>Böhme (1989)               |
| <b>Asia</b>   |                          |                                 |                          |                     |                                |                                                          |
| 7             | <i>V. bengalensis</i>    | HR ♂ > HR ♀                     | --                       | 99.17               | Terrestrial, arboreal, aquatic | Pianka (2004)                                            |
| 8             | <i>V. caerulivirens</i>  | --                              | --                       | 40                  | arboreal, semi-aquatic         | Ziegler et al. (2004)                                    |
| 9             | <i>V. ceramponensis</i>  | --                              | --                       | 40.9                | Semi-aquatic?                  | Philipp et al (2004)                                     |
| 10            | <i>V. doreanus</i>       | --                              | 13/600 ? sqm             | 46.0                | Terrestrial                    | Ziegler et al. (1999)                                    |
| 11            | <i>V. dumerili</i>       | --                              | --                       | 50.0                | Terrestrial, arboreal, aquatic | Bennett (2004)                                           |
| 12            | <i>V. finschi</i>        | --                              | --                       | 30.5                | Terrestrial, arboreal, aquatic | Phillip et al. (2004)                                    |
| 13            | <i>V. flavescens</i>     | --                              | --                       | 40.0*               | Terrestrial, arboreal, aquatic | Visser (2004)                                            |
| 14            | <i>V. indicus</i>        | --                              | --                       | 58.0                | Terrestrial, arboreal, aquatic | Wikramanayake & Dryden (1988)<br>Dryden & Ziegler (2004) |
| 15            | <i>V. jobiensis</i>      | --                              | --                       | 45.0                | Terrestrial, arboreal          | Horn (1977)<br>Philipp et al. (2004)                     |

| No               | Species                 | Home range (ha)                    | Density (per ha) | Max. adult SVL (cm)                      | Mode of life                         | Reference                                   |
|------------------|-------------------------|------------------------------------|------------------|------------------------------------------|--------------------------------------|---------------------------------------------|
| 16               | <i>V. juxtindicus</i>   | --                                 | --               | 50.4                                     | Terrestrial, arboreal, aquatic       | Böhme <i>et al.</i> (2004)                  |
| 17               | <i>V. komodoensis</i>   | 258-529<br>HR♂>HR♀                 | --               | 154.0                                    | Terrestrial, arboreal, aquatic       | Ciofi (2004)                                |
| 18               | <i>V. kordensis</i>     | --                                 | --               | 27.0                                     | Arboreal                             | Jacobs (2004)                               |
| 19               | <i>V. mabitang</i>      | --                                 | --               | 64.0                                     | Arboreal                             | Gaulke (2004)                               |
| 20               | <i>V. macraei</i>       | --                                 | --               | 36.0                                     | Arboreal                             | Böhme & Jacobs (2004)                       |
| 21               | <i>V. melinus</i>       | --                                 | --               | 42.0                                     | Arboreal, aquatic                    | Ziegler & Böhme (2004)                      |
| 22               | <i>V. olivaceus</i>     | Small                              | --               | 65.0                                     | Arboreal                             | Pianka (2004)                               |
| 23               | <i>V. prasinus</i>      | --                                 | --               | 29.5                                     | Arboreal                             | Greene (2004)                               |
| 24               | <i>V. rudicolis</i>     | --                                 | --               | 59.0                                     | Arboreal, terrestrial                | Bennett (2004)                              |
| 25               | <i>V. salvadorii</i>    | --                                 | --               | 85.0                                     | Arboreal                             | Horn (2004)                                 |
| 26               | <i>V. salvator</i>      | 1.4-31.7                           | --               | 92.0                                     | Terrestrial, arboreal, aquatic       | Traeholt (1997) a,b<br>Gaulke & Horn (2004) |
| 27               | <i>V. spinulosus</i>    | --                                 | --               | 31.2                                     | Arboreal                             | Philipp <i>et al.</i> (2004)                |
| 28               | <i>V. timorensis</i>    | --                                 | --               | 21.7*                                    | Terrestrial & arboreal               | King & Smith (2004)                         |
| 29               | <i>V. yuwonoi</i>       | --                                 | --               | 57.7*                                    | Terrestrial, arboreal, aquatic       | Philipp <i>et al.</i> (2004)                |
| <b>Australia</b> |                         |                                    |                  |                                          |                                      |                                             |
| 30               | <i>V. acanthurus</i>    | --                                 | --               | 25.0                                     | Terrestrial & arboreal,              | Dryden (2004)                               |
| 31               | <i>V. baritji</i>       | --                                 | --               | 25.2                                     | Terrestrial/saxicolous               | King (2004)                                 |
| 32               | <i>V. brevicauda</i>    | --                                 | 20               | 11.8                                     | Terrestrial                          | James (1994)<br>Pianka (2004)               |
| 33               | <i>V. caudolineatus</i> | --                                 | --               | 12.3♂<br>11.8♀                           | Terrestrial/saxicolous& semiarboreal | Thompson (2004)                             |
| 34               | <i>V. eremius</i>       | large                              | --               | 16.0                                     | Terrestrial                          | Pianka (2004)                               |
| 35               | <i>V. giganteus</i>     | 325.6± 127.0 (♂)<br>47.5 ± 9.1 (♀) | --               | 73.6♂<br>59.1♀                           | Terrestrial                          | Heger (2000)<br>Horn & King (2004)          |
| 36               | <i>V. gilleni</i>       | -                                  | --               | 18.6                                     | Terrestrial, arboreal                | Horn (2004)                                 |
| 37               | <i>V. glauerti</i>      | 1.25-7.36<br>HR♂>HR♀               | --               | <sup>a</sup> 21.5♂<br><sup>a</sup> 18.0♀ | Terrestrial/saxicolous, arboreal     | Sweet (1999)<br>Sweet (2004)                |

| No | Species               | Home range (ha)              | Density (per ha) | Max. adult SVL (cm)                      | Mode of life                                    | Reference                                      |
|----|-----------------------|------------------------------|------------------|------------------------------------------|-------------------------------------------------|------------------------------------------------|
| 38 | <i>V. gleopalma</i>   | 3.5-7.7<br>HR large>HR small | --               | <sup>a</sup> 29.0♂<br><sup>a</sup> 24.5♀ | Terrestrial/saxicolous                          | Sweet (2004)                                   |
| 39 | <i>V. gouldii</i>     | 8.91<br>HRlarge>HRsmall      | --               | <sup>a</sup> 59.0♂<br><sup>a</sup> 36.1♀ | Terrestrial                                     | Thompson (2004)                                |
| 40 | <i>V. keithhornei</i> | --                           | --               | 26.0                                     | Arboreal                                        | Irwin (2004)                                   |
| 41 | <i>V. kingorum</i>    | --                           | --               | 11.4                                     | Terrestrial/saxicolous                          | King (2004)                                    |
| 42 | <i>V. mertensi</i>    | --                           | --               | 48.0                                     | Semi-aquatic, arboreal                          | Christian (2004)                               |
| 43 | <i>V. mitchelli</i>   | limited                      | --               | 32.0                                     | Arboreal, semi-aquatic, terrestrial/ saxicolous | Schultz & Doody (2004)                         |
| 44 | <i>V. panoptes</i>    | --                           | --               | 74.0                                     | Terrestrial, arboreal                           | Christian (2004)                               |
| 45 | <i>V. pilbarensis</i> | --                           | --               | 16.9                                     | Terrestrial/saxicolous                          | King (2004)                                    |
| 46 | <i>V. primordius</i>  | --                           | --               | 12.0                                     | Terrestrial                                     | Husband & Christian (2004)                     |
| 47 | <i>V. rosenbergi</i>  | 19.44±4.58                   | --               | 47.0                                     | Terrestrial, arboreal                           | King & Green (1999)<br>King (unpubl.)          |
| 48 | <i>V. scalaris</i>    | <1.5 (♂)<br><1.0 (♀)         | --               | 25.3                                     | Arboreal                                        | Smith <i>et al.</i> (2004)                     |
| 49 | <i>V. semiremex</i>   | --                           | --               | 27.0                                     | Arboreal, semi-aquatic                          | Pianka (2004)                                  |
| 50 | <i>V. spenceri</i>    | --                           | --               | 55.0                                     | Terrestrial                                     | Vincent & Wilson (1989)                        |
| 51 | <i>V. storri</i>      | Fairly small                 | --               | 13.2                                     | Terrestrial                                     | Eidenmüller (2004)<br>Peters (1973)            |
| 52 | <i>V. tristis</i>     | 40.3 (♂)<br>3.7 (♀)          | --               | 30.5                                     | Arboreal,<br>Terrestrial/saxicolous             | Pianka (2004)<br>Thompson <i>et al.</i> (1999) |
| 53 | <i>V. varius</i>      | 65 ± 34 (♂)<br>25 (♀)        | --               | 76.5♂<br>57.5♀                           | Arboreal, Terrestrial                           | Weavers (1993)<br>Weavers (2004)               |
|    |                       | 184.5                        |                  |                                          |                                                 | Guarino (2002)                                 |

<sup>a</sup> average\*calculated using ratio of tail: SVL and total body length from Pianka *et al.* (2004)

Table 2. Body sizes, clutch sizes, and breeding periods of genus *Varanus* by continental groups.

| No            | Species                  | Clutch size                     | Mating period                                                        | Max. adult SVL (cm) | Sexual maturity                                       | Reference                                                                 |
|---------------|--------------------------|---------------------------------|----------------------------------------------------------------------|---------------------|-------------------------------------------------------|---------------------------------------------------------------------------|
| <b>Africa</b> |                          |                                 |                                                                      |                     |                                                       |                                                                           |
| 1             | <i>V. albigularis</i>    | Max. 50 (larger, larger clutch) | July/August- September (rain season)                                 |                     | 4-5 years                                             | Phillips (2004)                                                           |
| 2             | <i>V. exanthematicus</i> | 6-29 (Max. 41)                  | November                                                             | 75.0                | 27.4 cm SVL (♀)                                       | Bennett (2004)                                                            |
| 3             | <i>V. griseus</i>        | 10-20                           | June                                                                 | 58.3*               | 4-5 years                                             | Stanner (2004)                                                            |
| 4             | <i>V. niloticus</i>      | 5-60                            | End of rainy season (Sept-Nov, West Africa; march-May, South Africa) | 96.0*               | 3-4 years;<br>90-120 cm TL                            | Lenz (2004)                                                               |
| 5             | <i>V. ornatus</i>        | --                              | bimodal                                                              | 92.25*              |                                                       | Böhme & Ziegler (2004)                                                    |
| 6             | <i>V. yemenensis</i>     | --                              | --                                                                   | 59                  |                                                       | Gasperetti in Böhme (1989)                                                |
| <b>Asia</b>   |                          |                                 |                                                                      |                     |                                                       |                                                                           |
| 7             | <i>V. bengalensis</i>    | 8-29 <sup>capt.</sup>           | June-July (Monsoon)                                                  | 99.17               | 3-4 years <sup>capt.</sup><br>≥5 years                | Horn & Visser (1989)<br>Pianka (2004)                                     |
|               |                          | 20.2 <sup>a</sup>               |                                                                      |                     |                                                       | Auffenberg (1994)                                                         |
| 8             | <i>V. caerulivirens</i>  | --                              | --                                                                   | 40                  | --                                                    | Ziegler et al. (2004)                                                     |
| 9             | <i>V. ceramboensis</i>   | --                              | --                                                                   | 40.9                | --                                                    | Philipp et al. (2004)                                                     |
| 10            | <i>V. doreanus</i>       | --                              | --                                                                   | 46.0                | --                                                    | Böhme et al. (2004)                                                       |
| 11            | <i>V. dumerili</i>       | Max. 23 <sup>capt.</sup>        | --                                                                   | 50.0                | --                                                    | Bennett (2004)                                                            |
| 12            | <i>V. finschi</i>        | --                              | --                                                                   | 30.5                | --                                                    | Philipp et al. (2004)                                                     |
| 13            | <i>V. flavescens</i>     | 4-30 (mean: 16)                 | June-July (wet season)                                               | 40.0*               | 26.0 cm SVL (♂)<br>>25.0 cm SVL (♀)<br>3-4 yrs. (♂,♀) | Visser (2004)                                                             |
| 14            | <i>V. indicus</i>        | 10                              | Dry season                                                           | 58.0                | 32.0 cm SVL (♂)<br>27.5 cm SVL (♀)                    | McCoid (1993)<br>McCoid & Hensley (1991)<br>Wikramanayake & Dryden (1988) |
| 15            | <i>V. jobiensis</i>      | 20-30                           | Simulated wet season (capt.)                                         | 45.0                | --                                                    | Bayless & Dwyer (1997)<br>Horn (1977)                                     |
| 16            | <i>V. juxtindicus</i>    | --                              | --                                                                   | 50.4                | -                                                     | Böhme et. al. (2004)                                                      |

| No               | Species               | Clutch size                                      | Mating period                                   | Max. adult SVL (cm) | Sexual maturity                       | Reference                                           |
|------------------|-----------------------|--------------------------------------------------|-------------------------------------------------|---------------------|---------------------------------------|-----------------------------------------------------|
| 17               | <i>V. komodoensis</i> | Max.33,<br>mean 18                               | July-September<br>(dry season) <sup>capt.</sup> | 154.0               | 8-9 years<br>(♂ & ♀) <sup>capt.</sup> | Ciofi (2004)                                        |
|                  |                       | 1-30, mean 19                                    | May-August                                      |                     |                                       | Auffenberg (1981)                                   |
| 18               | <i>V. kordensis</i>   | --                                               | --                                              | 27.0                | --                                    | Jacobs (2004)                                       |
| 19               | <i>V. mabitang</i>    | 6-12                                             | May                                             | 64.0                | --                                    | Gaulke <i>et.al.</i> (2002)<br>Gaulke (2004)        |
| 20               | <i>V. macraei</i>     | Multiple 3                                       | --                                              | 36.0                | --                                    | Jacobs (2002)<br>Böhme & Jacobs (2004)              |
| 21               | <i>V. melinus</i>     | 1.5-3.4x (Max.12)                                | --                                              | 42.0                | --                                    | Ziegler & Böhme (2004)                              |
|                  |                       | 4x (2-7) <sup>capt.</sup>                        |                                                 |                     |                                       | Dedlmar & Böhme (2000)                              |
| 22               | <i>V. olivaceus</i>   | 4-11, mean 7.1                                   | June-September                                  | 65.0                | 45.0 cm SVL (3 years)                 | Auffenberg (1988, 1994)<br>Pianka (2004)            |
| 23               | <i>V. prasinus</i>    | 3x (2-4)                                         | --                                              | 29.5                | 2 years <sup>capt.</sup>              | Greene (2004)                                       |
| 24               | <i>V. rudicollis</i>  | 2-3x (Max.14, mean 8) <sup>capt.</sup>           | --                                              | 59.0                | --                                    | Bennett (2004)                                      |
| 25               | <i>V. salvadorii</i>  | 4-12 <sup>capt.</sup>                            | July & October <sup>capt.</sup>                 | 85.0                | -                                     | Horn (2004)                                         |
| 26               | <i>V. salvator</i>    | Multiple, 5-22, mean 13 (correlated with ♀ size) | All year, peak in August                        | 92.0                | 40 cm SVL (♂)<br>50 cm SVL (♀)        | Shine <i>et. al.</i> (1998)<br>Gaulke & Horn (2004) |
| 27               | <i>V. spinulosus</i>  | --                                               | --                                              | 31.2                | --                                    | Philipp <i>et al.</i> (2004)                        |
| 28               | <i>V. timorensis</i>  | --                                               | May-July                                        | 21.7*               | --                                    | King & Smith (2004)                                 |
| 29               | <i>V. yuwonoi</i>     | --                                               | --                                              | 57.7*               | --                                    | Philipp <i>et al.</i> (2004)                        |
| <b>Australia</b> |                       |                                                  |                                                 |                     |                                       |                                                     |
| 30               | <i>V. acanthurus</i>  | 2-18, mean 7.9                                   | August & November (end dry season)              | 25.0                | 8.9 cm SVL (♂)<br>10.2 cm SVL (♀)     | Dryden (2004)<br>King & Rhodes (1982)               |
| 31               | <i>V. baritji</i>     | 3-9                                              | July                                            | 25.2                | -                                     | King (2004)                                         |

| No | Species                 | Clutch size                        | Mating period                                         | Max. adult SVL (cm)                      | Sexual maturity                    | Reference                                                   |
|----|-------------------------|------------------------------------|-------------------------------------------------------|------------------------------------------|------------------------------------|-------------------------------------------------------------|
| 32 | <i>V. brevicauda</i>    | 2-3                                | September-October (Spring)                            | 11.8                                     | 8.2 cm SVL (♂)<br>9.4 cm SVL (♀)   | Pianka (1994)                                               |
|    |                         |                                    |                                                       |                                          | 7.0 cm SVL (♂)<br>8.3 cm SVL (♀)   | James (1996)                                                |
| 33 | <i>V. caudolineatus</i> | Mean 4.3                           | --                                                    | 12.3♂<br>11.8♀                           | --                                 | Pianka (1994)                                               |
|    |                         |                                    |                                                       |                                          | --                                 | Thompson (2004)                                             |
| 34 | <i>V. eremius</i>       | 2-6, mean 3.6                      | October - November (Spring)                           | 16.0                                     | 11.6 cm SVL (♂)<br>11.0 cm SVL (♀) | Pianka (2004)                                               |
| 35 | <i>V. giganteus</i>     | 13                                 | Spring & early Summer                                 | 73.6♂<br>59.1♀                           | 45.0 cm SVL (♂)<br>49.0 cm SVL (♀) | King (1989) <sup>mus</sup><br>Horn & King (2004)            |
| 36 | <i>V. gilleni</i>       | Mean 4                             | Max. October (♀ reproductive at longer period than ♂) | 18.6                                     | 10.0 cm SVL (♂)<br>9.5 cm SVL (♀)  | James et.al. (1992) <sup>mus</sup><br>Horn & Visser (2004)  |
| 37 | <i>V. glauerti</i>      | 3                                  | mid-May to mid-July                                   | <sup>a</sup> 21.5♂<br><sup>a</sup> 18.0♀ | 15.0 cm SVL (♂, ♀)                 | Sweet (2004)<br>James et.al. (1992)                         |
| 38 | <i>V. glebopalma</i>    | 5-7                                | August-October                                        | <sup>a</sup> 29.0♂<br><sup>a</sup> 24.5♀ | 17.0 cm SVL (♂, ♀)                 | Sweet (2004)<br>James et.al. (1992)<br>Barnett 1977)        |
| 39 | <i>V. gouldii</i>       | Mean 6.2                           | September-November                                    | <sup>a</sup> 59.0♂<br><sup>a</sup> 36.1♀ | 25.0 cm SVL                        | Pianka (1994)<br>Pianka (1970)<br>Thompson (2004)           |
| 40 | <i>V. keithhornei</i>   | 2x (2-4) <sup>capt.</sup>          | September-May <sup>capt.</sup>                        | 26.0                                     | --                                 | Irwin (2004)                                                |
| 41 | <i>V. kingorum</i>      | 5x (3-6) mean 4.5 <sup>capt.</sup> | February (end of wet season) <sup>mus.</sup>          | 11.4                                     | --                                 | James et.al. (1992)<br>Eidenmüller (2001)<br>King (2004)    |
| 42 | <i>V. mertensi</i>      | 3-14                               | February-July                                         | 48.0                                     | --                                 | Vincent & Wilson (1999)<br>Shine (1986)<br>Christian (2004) |

| No | Species               | Clutch size                    | Mating period                                   | Max. adult SVL (cm) | Sexual maturity              | Reference                                                                          |
|----|-----------------------|--------------------------------|-------------------------------------------------|---------------------|------------------------------|------------------------------------------------------------------------------------|
| 43 | <i>V. mitchelli</i>   | 7-12                           | April (Late wet season) & June (mid-dry season) | 32.0                | 22 cm SVL                    | Shine (1986)                                                                       |
|    |                       | 16-20 <sup>capt.</sup>         | April                                           |                     |                              | Schultz & Doody (2004)                                                             |
| 44 | <i>V. panoptes</i>    | 6-14                           | April                                           | 74.0                | --                           | Christian (2004)                                                                   |
| 45 | <i>V. pilbarensis</i> | Max.24 in 6x                   | February-March                                  | 16.9                | --                           | King (2004)                                                                        |
| 46 | <i>V. primordius</i>  | 2-5, mean 3.3 <sup>capt.</sup> | --                                              | 12.0                | 9-12 months <sup>capt.</sup> | Husband (2001)                                                                     |
| 47 | <i>V. rosenbergi</i>  | 10-17                          | January-February (mid-summer)                   | 47.0                | --                           | Green <i>et.al.</i> (1971)<br>King & Green (1979)<br>King (unpubl.)                |
| 48 | <i>V. scalaris</i>    | 3-12, mean 7.7                 | May-mid-June                                    | 25.3                | 12.5 cm SVL (♂)              | Smith <i>et al.</i> (2004)                                                         |
| 49 | <i>V. semiremex</i>   | 2-14, mean 5.9                 | February-April (late wet season)                | 27.0                | 15.0 cm SVL                  | King (2004)                                                                        |
| 50 | <i>V. spenceri</i>    | 1x (11-35)                     | August                                          | 55.0                | 28.0 cm SVL (♂)              | Lemm & Bedford (2004)<br>Greer (1989)<br>Vincent & Wilson (1989)<br>Fyfe (unpubl.) |
| 51 | <i>V. storri</i>      | 1-6, mean 3.9                  | February-March & July-November                  | 13.2                | 9.0 cm SVL (♂,♀)             | Eidenmüller (2004)<br>Peters (1973)<br>James <i>et.al.</i> (1992)                  |
| 52 | <i>V. tristis</i>     | 5-17, mean 10.1                | October-November (Spring)                       | 30.5                | 20.0 cm SVL (♂,♀)            | Pianka (1994)<br>Pianka (2004)                                                     |
| 53 | <i>V. varius</i>      | 5 <sup>capt.</sup>             | mid-November to early January                   | 76.5♂<br>57.5♀      | 3 years <sup>capt.</sup>     | Carter (1990)<br>Weavers (2004)                                                    |
|    |                       |                                | March <sup>capt.</sup>                          |                     |                              | Boylan (1995)                                                                      |

<sup>capt.</sup>= in captivity<sup>mus.</sup>= museum specimens